



**LOS ANGELES COUNTY DEPARTMENT OF  
PUBLIC HEALTH  
ACUTE COMMUNICABLE DISEASE  
CONTROL PROGRAM**

**SCABIES PREVENTION AND  
CONTROL GUIDELINES  
ACUTE AND SUB-ACUTE  
CARE FACILITIES**




## CONTENTS

I.	Introduction.....	3
II.	General Information.....	3
	A. Biology of the Scabies Mite.....	3
	B. Clinical Presentation .....	4
	1. Typical Scabies .....	4
	2. Atypical Scabies.....	5
	C. Epidemiology of Scabies.....	5
	1. Transmission.....	6
	2. Incubation Period .....	6
	3. Period of Communicability .....	6
	D. Diagnosis .....	6
III.	Scabies Prevention and Control Programs.....	6
IV.	Scabies Outbreak Management .....	7
	A. Overview.....	7
	B. Summary of Action Steps .....	8
	C. Management of Symptomatic Cases .....	9
	1. Symptomatic Health Care Workers .....	9
	2. Symptomatic Patients .....	10
	D. Management of Contacts.....	10
	1. HCW Contacts .....	11
	2. Patient Contacts.....	11
	E. Extending Recommendations for Prophylaxis .....	12
	F. Notification of Staff, Visitors and Household Members .....	12
V.	Reporting.....	12
VI.	Appendices.....	13
	A. Diagnosis of Scabies by Skin Scraping .....	14
	B. Treatment and Prophylaxis of Scabies .....	16
	a) Application of Scabicides: General Principles .....	16
	b) Scabicides.....	16
	c) Treatment Regimen: Typical Scabies Infection.....	17
	d) Treatment Regimen: Crusted (Norwegian) or Atypical Scabies ....	18
	e) Treatment Failures.....	19
	C. Treatment Options .....	20
	D. Directions For Scabies Treatment or Prophylaxis With Elimite .....	21
	E. Scabies Fact Sheet.....	22
	F. Isolation and Environmental Control For Patients with Scabies.....	25
	G. Scabies Case/Contact Line-Listing Form: Employees .....	26
	H. Scabies Case/Contact Line-Listing Form: Patients.....	27
	I. Sample Letter to Discharged Patients .....	28
	J. Sample Letter to Physicians of Discharged Patients .....	29
	K. Scabies Outbreak Management Checklist.....	30
VII.	References.....	31

## I. INTRODUCTION

Scabies is an ectoparasitic infestation of the skin caused by the human itch mite, *Sarcoptes scabiei* var. *hominis*. The actual incidence of scabies in Los Angeles County (LAC) is unknown because single occurrences of typical scabies are not reportable to the county health department. Outbreaks of scabies infestations in health care facilities (HCF) and single cases of atypical scabies are reportable to the local health department and present a continuing public health problem in LAC. While regarded as a nuisance disease by many health care professionals, scabies outbreaks in HCFs can be extremely costly in terms of direct and indirect costs of outbreak management, poor public relations, and discomfort and anxiety of affected patients, employees, and their family members. Furthermore, secondary bacterial skin infection is a common complication of scabies infestation that, in elderly or immunocompromised individuals, can lead to sepsis and even death.




A number of factors influence the extent of scabies transmission within a facility, including the mite load and the required level of care of the source case, as well as the duration of the exposure period. Each HCF scabies outbreak is unique and requires an individualized approach. Due to the increase in the number of outbreaks reported by HCF in LAC, these guidelines were developed to provide a rational approach to the prevention and control of scabies. They are intended to assist infection control committees and administrators of HCF in developing a scabies prevention and control program for their facility, and they are based on best practices, state and federal scabies guidelines, current literature and the extensive experience of Acute Communicable Disease Control Program (ACDC) staff.

## II. GENERAL INFORMATION

### A. BIOLOGY OF THE SCABIES MITE

Infestation begins when one or several pregnant female mites are transferred from the skin of an infested person to the skin of an uninfested person. After transfer from the skin of an infested person, or, rarely, from fomites, to the skin of an uninfested person, the adult female mite travels on the skin surface at the rate of about 1 inch per minute seeking a burrow site. After finding a suitable location, she burrows into superficial layers of the skin, forming a slightly elevated narrow tunnel where she deposits 2 to 3 eggs daily during her 4 to 6 week life span. The eggs progress through larval and nymphal stages to form adults in 10 to 17 days. The adults migrate to the skin surface and mate. The males die quickly and the females penetrate the skin and repeat the cycle. The mite requires human skin to complete its life cycle and is unable to survive off the host at room temperature for more than 3 to 4 days.

## Life Cycle:



From the CDC website: <http://www.dpd.cdc.gov/dpdx/HTML/Scabies.htm>

## B. CLINICAL PRESENTATION

Scabies infestations are generally categorized as typical or atypical (crusted, keratotic or Norwegian).

### 1. Typical Scabies



Patients with typical (conventional) scabies usually have only 10 to 15 live adult female mites on the body at any given time. Usually, only one or two mites, and frequently none, are recovered from skin scrapings. Intense pruritis, usually worse at night, and a papular rash with or without burrows occur. The rash and pruritis result from an immune-mediated delayed hypersensitivity reaction to the mite, its eggs, and fecal material.

Areas of the body commonly involved are wrists, finger webs, antecubital fossae, anterior axillary folds, breasts, waistline, lower abdomen, genitals, and buttocks. The scalp and face are rarely involved in adults, but may be observed in young children with scabies.

## 2. Atypical Scabies

When diagnosis and treatment are delayed, scabies can have an unusual or atypical presentation, involving heavy infestation with hundreds to thousands of mites. Atypical clinical presentations are more prevalent in institutionalized or debilitated patients, or



those who are immunosuppressed from underlying disease or drug therapy. When extensive hyperkeratotic skin lesions with crusting and scaling develop, the infestation is called **crusted scabies** or hyperkeratotic (formerly “Norwegian”) scabies. Crusted scabies is highly contagious because thousands of mites are imbedded in the thick crusts and easily shed in scales and flakes from affected skin. Crusted scabies is commonly misdiagnosed by dermatologists, and patients with crusted scabies may develop symptoms of typical scabies in as little as a few days.

## C. EPIDEMIOLOGY OF SCABIES

### 1. *Transmission*

Transfer of the mite is usually from one person to another by direct skin-to-skin contact. Procedures such as bathing a patient, applying body lotions, back rubs, or any extensive hands-on contact can provide an opportunity for mite transmission. Mites may also be transmitted via clothing, bed linen or other fomites. Fomites play a minor role in situations where the infestation in the source case is typical scabies; the inanimate environment of patients with crusted scabies, however, has been shown to be heavily contaminated with infectious mature and immature mites. In HCF, scabies may be introduced into the facility by a newly admitted resident with an unrecognized infestation or by visitors or health care workers as a result of contact with an infested person in the home or community.

### 2. *Incubation Period*

In a previously unexposed healthy individual, the interval between exposure and the onset of itching is usually 4-6 weeks. In persons who have been sensitized to the mite by a previous infestation, re-exposure may produce

symptoms in 48 hours or less (owing to prior sensitization to the mite and its saliva and feces). Following exposure to a source case with crusted scabies involving extremely large numbers of mites, the incubation period may be reduced from the usual time of 4-6 weeks to as little as a few days.

### 3. *Period of Communicability*

Since the scabies mite is an ectoparasite, **an exposed individual is potentially immediately infectious to others, even in the absence of symptoms.** Cases are communicable from the time of infestation until mites and eggs are destroyed by treatment.

## D. DIAGNOSIS

Definitive diagnosis requires microscopic identification of the mite and/or its eggs or fecal pellets on specimens collected by skin scraping, biopsy or other means (Appendix A, "Diagnosis of Scabies by Skin Scraping"). The yield from skin scrapings is highly dependent on the experience of the operator and the severity of the infestation. A negative skin scraping from a person with typical scabies does not rule out scabies infestation; mites are easily recovered, however, in skin scrapings from persons with crusted scabies.

## III. SCABIES PREVENTION AND CONTROL PROGRAMS

It is recommended that HCF incorporate a scabies prevention program that involves all levels of the health care team. The program should include an assessment of the skin, hair and nail beds of all new admissions as soon as possible following arrival. Pruritus, rashes and skin lesions should be documented and brought to the attention of the nursing supervisor and the attending physician for further follow-up.

Essential elements of a successful scabies prevention program include:

1. Written policies and procedures for prevention and control of nosocomial scabies;
2. Health care workers who are trained to be suspicious of scabies in themselves or their patients if unexplained rash or pruritus occurs in themselves or their patients, and to report such occurrences to their supervisors;
3. A policy to screen newly admitted patients for scabies during the initial assessment (especially if transferred from another healthcare facility) and any suspect patient will immediately be placed on contact isolation until examined for scabies;
4. A policy that all new employees (especially employees who work at more than one facility) will be screened for scabies as part of pre-employment screening;

5. Access to and use as needed of the diagnostic skills of a consultant experienced in recognizing scabies to evaluate difficult or unusual cases or response to treatment;
6. Assurance of adequate support from hospital administration, medical staff, infection control, employee health and line staff for appropriate evaluation and treatment of employees, in-house patients and exposed discharged patients should an outbreak of nosocomial scabies occur.

## **IV. SCABIES OUTBREAK MANAGEMENT**

### **A. OVERVIEW**

The primary goal of an outbreak investigation is to identify risk factors contributing to the outbreak and to take corrective action to prevent further transmission of scabies cases. In general, an outbreak can be defined as an increase in the incidence of new cases above baseline within a defined period of time and within a defined geographical location (nursing unit, one floor or one wing, a department or, in some cases, the entire hospital).

A “baseline” for scabies is not a standard hospital measurement and the definition of a scabies outbreak may be multifaceted. A case definition should be developed to determine whether an outbreak has occurred or to estimate the magnitude of the outbreak. The following are examples of scabies hospital outbreak definitions:

- Two (2) or more confirmed (positive skin scraping) cases of scabies identified in patients, healthcare workers, volunteers and/or visitors during a two (2) week period of time, or
- One (1) confirmed (positive skin scraping) and at least two (2) clinically suspect cases identified in patients, healthcare workers, volunteers and/or visitors during a two (2) week period of time, or
- At least two (2) clinically suspect cases identified in patients, healthcare workers, volunteers and/or visitors during a two (2) week period of time.

Nosocomial transmission is highly probable if scabies is confirmed in two or more health care workers who have worked in the same area of the facility within the previous six weeks and who do not have an apparent source of exposure outside the facility.

Outbreak management should include planning for protective equipment and pharmacy supplies. Provisions should be made for obtaining additional personal protective equipment (PPE) such as disposable, long sleeve gowns and gloves. The pharmacy should also make arrangements for obtaining permethrin (Elimite) 5% cream. If ivermectin (Stromectol) is recommended as a scabies treatment option the pharmaceutical company should be notified and arrangements made for overnight shipping if necessary.

## B. SUMMARY OF ACTION STEPS

1. Immediately remove from work any health care worker with signs and symptoms of scabies and refer to employee health or other designated consultant or clinician experienced in the diagnosis of scabies.
2. Evaluate patients on affected units and immediately place patients with suspected scabies in contact isolation (Appendix F)
3. Report nosocomial scabies outbreaks and **single cases** of crusted scabies infestation to LAC ACDC.
4. Meet with key staff to coordinate control measures. Representatives from the following departments should be included: Administration, Employee Health, Environmental Services, Infection Control, Pharmacy, Medical Staff, and Nursing. One person, generally the Infection Preventionist (formerly Infection Control Professional), should coordinate control measures and should be given adequate resources to accomplish this objective in a timely and efficient manner.
5. Search for a possible source case. If two or more employees working in the same unit/area are diagnosed with scabies, it is likely that the source case was a patient with **crusted scabies** infestation.
6. Confirm the presence of scabies by microscopic identification of the mite or its products (skin scraping) in one or more symptomatic patients or employees.
7. Prepare a line listing of symptomatic patients and health care workers and a separate line listing of their contacts (Appendix G and H). Evaluate contacts for scabies.
8. Treat symptomatic patients and health care workers with an approved scabicide, provide prophylactic scabicide to all contacts of symptomatic cases, and perform environmental cleaning of affected units. Ideally, these steps (treatment, prophylaxis, and environmental cleaning) should all be accomplished within the same 24 hour period to prevent re-infestation of treated or prophylaxed individuals.
9. Provide training to all staff on the signs and symptoms of scabies. Stress that people can be infested and contagious for up to 6 weeks before any symptoms start.
10. Perform environmental cleaning of affected units (Appendix F).

11. Arrange for follow-up evaluation and prophylactic treatment of discharged patients who were contacts to scabies. (Appendix I and J).

## C. MANAGEMENT OF SYMPTOMATIC CASES

Often the first indications of a scabies outbreak are complaints of itching and rash in two or more health care workers or patients/residents. Properly performed skin scrapings will almost always be positive in persons with crusted scabies but are generally negative in cases of typical scabies, even when performed by experienced operators. None-the-less, it is recommended that efforts be made to confirm the diagnosis of scabies (perform skin scraping) in at least one symptomatic individual.

### 1. Symptomatic Health Care Workers

Health care workers (HCW) refer to all facility employees, contract employees, medical staff, house staff, students, religious workers and volunteers, etc.

- a. Immediately remove from work any HCW with signs or symptoms consistent with scabies and refer to employee health or other designated consultant experienced in the diagnosis of scabies. Confirm the presence of scabies by microscopic identification of the mite or its products in one or more symptomatic patient or employee (Appendix A).
- b. Prepare a line-listing of **symptomatic HCWs** that includes name, age, gender, symptoms, date of onset, result of scabies evaluation, any prior treatment for scabies, usual work and "float" assignments from six weeks before onset of symptoms until the current date, and symptoms in household or other close contacts (Appendix G).
- c. Treat all HCWs with confirmed or suspected scabies infestation with an approved scabicide according to consultant's recommendation or as outlined in Appendix B. **Review scabicide package insert before prescribing, dispensing, or applying scabicide. Provide clearly written instructions for proper application of dispensed scabicide** (Appendix D). Re-evaluate cases weekly to monitor response to treatment.
- d. Symptomatic HCWs can return to work as soon as treatment is completed but should use gowns and gloves for direct patient care to prevent reinfestation until all control measures for affected units/areas have been completed.
- e. Provide scabicide prophylaxis, along with written instructions for application, for all household contacts of symptomatic HCWs.

## 2. Symptomatic Patients

- a. Immediately place any patient in whom scabies infestation is suspected in contact isolation as outlined in the CDC “Guideline for Isolation Precautions in Hospitals” and Appendix F, “Isolation and Environmental Control for Patients with Scabies.” Maintain contact isolation until treatment is completed and/or case is determined by dermatology consultant or other experienced designee to be non-infectious.
- b. Attempt to confirm the diagnosis of scabies by microscopic identification of the mite, its eggs, or fecal pellets if the patient is the suspected source of the outbreak or if the clinical diagnosis of scabies is in question (Appendix A). If patient is a suspected case of **crusted scabies**, obtain dermatology consult for assistance with diagnosis and management.
- c. Treat with an approved scabicide according to consultant’s recommendation or as described in Appendix B.
- d. Perform environmental cleaning of case-patient’s room/ area as described in Appendix F.

## D. MANAGEMENT OF CONTACTS

**Contacts to typical (non-crusted) scabies** cases are defined as persons who had “hands-on” contact, handled infested clothing or bed linen, or slept in the same bed as the case-patient during the exposure period.

**Contacts to atypical (crusted) scabies** also include persons who had substantial contact with a crusted scabies case-patient’s environment, including HCWs who worked (regular or “float” assignment) on the same unit/area as the case-patient during the exposure period. If the case-patient was housed on more than one unit before control measures were initiated, each unit must be considered affected.

The outbreak **exposure period** is the period between the admission date of a scabies case and the date the condition is correctly diagnosed and control measures are implemented. If the case-patient is a long-term care resident, this period extends from six weeks prior to onset of symptoms. The identification of two or more symptomatic HCWs assigned to a particular unit suggests that prophylaxis is indicated for all unit/area contacts, whether or not they were direct contacts of a known scabies case.

Occasionally, a source case cannot be identified. In this situation, the exposure period should be considered to extend from 6 weeks before onset of symptoms (most often in HCWs) to the date of implementation of control measures.

## 1. HCW Contacts

- a. Identify and prepare a line listing of all HCWs who were direct contacts to a patient or fellow employee with scabies during the exposure period (Appendix G).
- b. Interview HCW contacts to determine presence of scabies symptoms and possible source of exposure; manage as a case if symptomatic.
- c. Provide prophylactic scabicide along with written instructions for application, as described in Appendix B, to all HCW with direct contact to a scabies case. HCW scabies contacts who refuse prophylactic treatment must be required to **wear gowns and gloves for contacts with patients or fellow employees for 6 weeks from the date of last potential exposure** (usually 6 weeks from implementation of control measures).

## 2. Patient Contacts

- a. Identify and prepare a line listing of all patients who were contacts to a patient with scabies or employee with scabies during the exposure period (Appendix H). This includes patients who resided on the same ward as a crusted scabies case during the exposure period and those who were already discharged.
- b. Examine **in-house patient contacts** to determine presence of signs and symptoms of scabies. If symptomatic, manage as a case.
- c. Apply prophylactic scabicide, as described in Appendix B, to in-house patients with direct contact to a scabies case.
- d. Patients who resided on the same ward as a crusted scabies patient with no direct contact to the case should be monitored with daily skin observations until six weeks past the date of last potential exposure and follow-up as appropriate.
- e. Notify **discharged patient contacts** of their potential exposure to scabies. Screen discharged patients for symptoms of scabies. Symptomatic patients should receive treatment and their family contacts should receive prophylaxis. Asymptomatic patients should be directed to follow up with their physician regarding possible scabies exposure and at a minimum, observe skin daily until six weeks past the date of last potential exposure. **It is the responsibility of the facility to ensure that all discharged exposed patients receive appropriate follow-up**

(Appendices I and J, “Sample letters to discharged patients and their physicians”).

- f. Notify facilities to which patient contacts have been transferred of their potential exposure.

#### **E. EXTENDING RECOMMENDATIONS FOR PROPHYLAXIS**

Facility-wide (mass) prophylaxis of all patients and at-risk employees (employees involved in direct patient care or exposed to patient care environment) should be considered if positive scrapings are found in patients or employees assigned to 2 or more areas of the facility where no direct link with an infested case-patient or HCW can be established.

#### **F. NOTIFICATION OF STAFF, VISITORS AND HOUSEHOLD MEMBERS**

Healthcare workers, volunteers, family members, sexual partners, and anyone else who have had contact with a scabies case should be notified immediately of the facility outbreak and assessed for symptoms. A scabies fact sheet and notification letter, which includes information about the scope of the outbreak and strategies that are being implemented to control the outbreak and prevent future cases, should be distributed to the above groups.

### **V. REPORTING**

Single cases of atypical (crusted) scabies and all outbreaks of scabies are required to be reported to Los Angeles County Department of Public Health during **normal business hours, Monday through Friday, 8:00 A.M. to 5:00 P.M.** Acute care facilities should report by phone to their ACDC Liaison Public Health Nurse or the Hospital Outreach Unit at (213) 240-7941. Long term care facilities and home health agencies should report to the Morbidity Unit at (888) 397-3993. Outbreaks are also reportable to Los Angeles County Department of Public Health, Health Facilities Division of Licensing and Certification.

The Scabies Prevention and Control Guidelines Acute and Sub-Acute Care Facilities and additional scabies information and resources can be found at the ACDC website [www.ph.lacounty.gov/acd/diseases/scabies.htm](http://www.ph.lacounty.gov/acd/diseases/scabies.htm).

# APPENDICES

## APPENDIX A

### DIAGNOSIS OF SCABIES BY SKIN SCRAPING

The diagnosis of scabies can be confirmed by microscopic identification of mites, eggs or scybala (fecal pellets) from specimens obtained by skin scrapings. A physician or nurse from the facility can be taught the procedure by a dermatologist, the consulting physician or by a nurse or technician who has had professional training to perform the procedure. A confirmed diagnosis of scabies should be made in at least one symptomatic case before recommending wide-spread scabicide prophylaxis in health care facility outbreaks. A physician, nurse or other healthcare professional who has been trained to perform the procedure should only do skin scrapings.


#### Equipment

1. Gloves and gowns
2. Magnifying glass
3. Light source such as goose neck lamp
4. Felt tip pen, green or blue washable ink
5. Alcohol swabs/wipes
6. #15 scalpel blade and handle,
7. Glass slides or culettes for scraping
8. Mineral oil or immersion oil
9. Slides and cover slips
10. Laboratory requisition forms
11. Sharps container
12. Clear nail polish
13. Microscope

#### Procedure

Observe patient's skin with a magnifying lens and look for lesions suggestive of scabies infestation. The shoulders, back, abdomen, hands, wrists, elbows, buttocks, axillae, knees, thighs and breasts are common sites for burrows.

1. Use hand magnifying lens to identify recent burrows or papules. A bright light and magnifying lens will assist in visualizing the tiny dark speck (the mite) at the end of the burrow.



2. Identify these high yield lesions by applying mineral oil (best used over dry scaly areas) or by applying the burrow ink test to possible burrows. The burrow ink test is done by using a wide felt tip pen (blue or green are best) over burrows and then wiping off with an alcohol swab. The alcohol will remove most surface ink, but will not remove the ink taken up by the burrow, thus leaving a dark irregular line.

3. Apply mineral oil or preferably microscope immersion oil to lesions or scalpel blade and glass slides.
4. Scrape non-excoriated, non-inflamed areas (burrows and papules) vigorously with a #15 scalpel blade or glass slide held at a 90-degree angle to the skin, while holding the skin taut, until the stratum corneum is removed. (Vigorous scraping appropriately results in a few red blood cells visible under the microscope, but there should not be frank bleeding.) Some practitioners prefer using a small curette. Change blades or curettes between scrapings on different persons. Blades can be placed on and removed from the handle with a forceps. Used blades must be placed in a sharps container.
5. Transfer skin scrapings from at least 6 different sites to a single slide or to separate slides. These scrapings can be pushed onto the slide edge and then moved to the center of the slide.
6. Place the cover slip over the slide.
7. Examine entire slide methodically under low power (2.5-4x) and then at 25-50x magnification. Microscopic examination of the skin scrapings should be performed at the facility; however, if the practitioner is not trained in reading the prepared slides, the cover slip should be secured to the slide at all edges with clear nail polish and transported by courier, or by mail (in a secure mailer) to a hospital or laboratory or to a physician's office with prior-arrangements.

## APPENDIX B

### TREATMENT AND PROPHYLAXIS OF SCABIES

#### A. Application of Scabicides: General Principles

1. Gowns and gloves are worn when applying scabicides to patients.
2. Bathe patients as usual and change bed linens. Allow skin to cool completely.
3. Apply scabicide to every square inch of skin, from the posterior ear folds down over the entire body. Include intergluteal cleft, umbilicus, skin folds, palms and soles, and webs between fingers and toes. If scabicide is washed off during handwashing, toileting, or perineal care, it must be reapplied.
4. In infants and young toddlers, the elderly, and the immunocompromised, the head (forehead, temples, and scalp) requires application of scabicide. Pay close attention to the area behind the ears. Do not get the scabicide near the eyes or mouth. Prior treatment failure may be an indication to include the head upon retreatment.
5. Fingernails and toenails should be clipped and scabicide applied under nails.
6. Follow directions and precautions outlined in the package insert accompanying scabicide.
7. A cleansing bath is taken when scabicide is to be removed.
8. Linens and clothing are changed after treatment. Contaminated clothing and linens may be 1) dry-cleaned or 2) washed in the hot cycle of the washing machine and dried in the hot cycle of the dryer for 10-20 minutes.
9. Provide detailed written instructions for scabicide use when dispensing scabicide for home application by employees and household members.

#### B. Scabicides

1. 5% permethrin cream (*Elimite, Acticin*) - currently considered drug of choice.
  - a. The usual adult dose is 30 grams. A 60 gram tube should treat two adults.
  - b. For adults and children, the cream should be massaged into the skin from below the chin to the soles of the feet. Scabies rarely infests the scalp of adults, although the hairline, neck, temple, and forehead may be infested in

infants and geriatric patients. Infants should be treated on the scalp, temple and forehead.

- c. The patient should be instructed to remove the medication by thoroughly bathing 8 to 14 hours after application. Contact with the eyes and mouth should be avoided. If contact with the eyes occurs, they should be immediately flushed with water.
  - d. Permethrin is regarded as safe for children two months of age and older. No instance of toxicity following accidental ingestion has been reported. The most commonly reported side effects are pruritus, edema and erythema, which may continue for up to two weeks after treatment. Patients should be told that the itching or stinging of scabies infestation may continue after treatment, and repeated application of the scabicide should be avoided. Demonstrable living mites after 14 days indicate that retreatment is necessary.
  - e. Although animal studies showed no adverse effects to reproductive function or damage to the fetus, no adequate studies have been done on pregnant women. Therefore, permethrin should be used during pregnancy only when clearly necessary. If treatment is necessary for lactating mothers, breast-feeding should be discontinued during the treatment period.
2. 10% crotamiton cream or lotion (*Eurax*)
- a. Massage thoroughly into skin once a day for two to five days. Remove by bathing 48 hours after last application.
  - b. Crotamiton is 60% effective when full five-day course is given.
  - c. Can be used on young children and elderly with dry, sensitive, but no denuded skin.
  - d. Avoid contact with eyes and mucous membranes.

3. Ivermectin (*Mectizan* or *Stromectol*)

Ivermectin is an antiparasitic agent shown to be safe and effective for treatment of onchocerciasis in humans in Africa. Experience with the agent in a single dose administered orally is limited, but encouraging, for treatment of crusted scabies or for infestations that do not respond to topical therapy. It is not yet approved by the Food and Drug Administration, but can be obtained from the manufacturer (Merck and Co, West Point, Pa.). A recent report described increased mortality from all causes in a small series of elderly patients who had received ivermectin.

### **C. Treatment Regimen for Typical Scabies Infestation**

1. A single adequate application of 5% permethrin cream is usually sufficient to eradicate typical scabies, whether a symptomatic case or asymptomatic carrier. Reevaluate response to treatment in 14 days.
2. In facilities with recurrent or endemic scabies or when application of scabicide for treatment of symptomatic scabies is not performed by a trained individual, a second application 3-7 days after the first is recommended by some authorities.
3. Asymptomatic contacts, including household and sexual contacts, of persons with clinical or confirmed scabies require one treatment with reevaluation in 14 days.

#### **D. Treatment Regimens for Crusted (Norwegian) or Severe Atypical Scabies**

1. Cases of crusted scabies and other variants of severe atypical scabies are best managed with the assistance of a dermatologist. Controlled studies to determine the most effective regimen for treatment of crusted scabies infestation have not been performed. The following regimens were selected from several that have appeared in the literature and have been successful in single or small series of cases. They are included as examples and are not necessarily endorsed by ACDC.
2. Patients with crusted or keratotic lesions should be soaked in a tub of lukewarm water for 10 minutes immediately prior to application of scabicide to hydrate the skin; use of keratolytic agents (e.g., salicylic acid) may soften scales and enhance penetration of scabicide. Allow skin to cool before applying scabicide.
3. Regimen A
  - a. 5% permethrin cream for 12 hours, followed by repeat application of 5% permethrin cream for 12 hours, wash off.
  - b. After seven days, repeat step (a), above.
  - c. Seven days following last treatment, obtain scrapings from at least 3 sites. If scrapings are positive or if symptoms unabated, treat again.
4. Regimen B
  - a. Apply 5% permethrin cream, as previously described, on day one.
  - b. Apply 10% crotamiton lotion, as previously described, on days 2-6.
  - c. Reapply 5% permethrin cream on day 7.

- d. Reassess on days 7 through 14, obtain scrapings from at least 3 sites in one month. If scrapings are positive or if symptoms unabated, begin regimen again.
5. Regimen C
    - a. Ivermectin 200 ug/kg in a single oral dose in combination with 5% permethrin cream on day one.
    - b. Two weeks after therapy (day 15), obtain scrapings from at least 3 sites. If scrapings are positive or if symptoms unabated, treat again.

## E. Treatment Failures

1. Treatment failures can result from:
  - a. Inadequate application of scabicide;
  - b. Infected, crusted, or keratotic lesions with insufficient penetration of scabicide;
  - c. Reinfestation from untreated contacts;
  - d. Resistance of mites to scabicide.
2. Pruritus and rash can continue for 1-4 weeks after treatment and should not be considered evidence of treatment failure until one month after last treatment. To ameliorate these signs and symptoms, some dermatologists use 1% hydrocortisone cream or triamcinolone cream (0.1%-0.025%) applied to the most intense rash sites **after** the first scabicide treatment. Oral antihistamines are also used to alleviate the hypersensitivity response.

### APPENDIX C

Patient Being Treated	Treatment Options	Dose	How To Treat	How Long is Treatment	Who Can Be Treated?
Typical Scabies	<u>Treatment A</u> 5% permethrin cream ( <i>Elimite, Acticin</i> )	Adult dose – 30 grams  60 gram tube can treat two adults	Massage cream into skin from under chin to soles of feet  Attention to hairline, neck, temple in geriatric patients	One treatment usually sufficient  May repeat if needed 7 days after 1 <sup>st</sup> treatment	Cases > 2 months, healthy adults  Used for prophylaxis of asymptomatic contacts
	<u>Treatment B</u> Ivermectin ( <i>Mectizan or Stromectol</i> ) oral antiparasitic  Used for patients who have failed treatment with or cannot tolerate topical treatment	200 mcg/kg	Given orally to treat suspect/confirmed cases of scabies	Single dose; 2 <sup>nd</sup> dose may be necessary to eliminate infection	Cases > 12 years
Atypical Scabies	<u>Treatment A</u> 5% permethrin cream ( <i>Elimite, Acticin</i> )	Adult dose – 30 grams	Massage cream into skin from under chin to soles of feet	Apply once, 2 <sup>nd</sup> application 12 hrs later  May repeat if needed 7 days after 1 <sup>st</sup> round of treatment	Cases > 2 months, healthy adults
	<u>Treatment B</u> 5% permethrin cream ( <i>Elimite, Acticin</i> )  10% crotamiton lotion ( <i>Eurax</i> )	Adult dose – 30 grams  Enough lotion to cover skin chin to feet	Apply permethrin once as above and again 12 hrs later on day 1 and day 7  Apply crotamiton as above on days 2-6	One week long treatment sufficient; reassess 7 days after treatment completed	Cases > 2 months, healthy adults
	<u>Treatment C</u> 5% permethrin cream ( <i>Elimite, Acticin</i> )  Ivermectin ( <i>Mectizan or Stromectol</i> ) oral antiparasitic	Adult dose – 30 grams  200 mcg/kg	Apply permethrin once as above and again 12 hrs later  Single oral dose	One treatment; reassess 14 days after treatment	Cases > 12 years

## **APPENDIX D**

### **DIRECTIONS FOR SCABIES TREATMENT OR PROPHYLAXIS WITH ELIMITE**

Clothing, towels, and bed linen that have been used within the last four days should be machine washed and dried using the hot cycle for 10-20 minutes. Articles that cannot be washed can be dry cleaned or tied in a plastic bag for a week. Floors and carpets should be vacuumed and the vacuum bag placed in a plastic bag and discarded.

#### **DIRECTIONS**

1. Take a bath, soaping the body completely, rinse well, and then dry thoroughly. Allow your body to cool.
2. Apply Elimite into the skin from the chin to the soles of the feet. Scabies rarely infests the scalp of adults, although the hairline, neck, temple, and forehead may be infested in infants and geriatric patients. Pay particular attention to skin folds and creases. Avoid contact with the eyes. Reapply if washed off following use of the toilet, handwashing, etc.
3. Put on clean clothing. Use freshly laundered bed linens and towels.
4. Leave cream on for at least 8 hours but no more than 14 hours, and then take a warm shower or bath, soaping the body completely, rinsing and drying well.
5. Put on clean clothing. Re-laundry towels and bed linens used during treatment.
6. Itching may continue for days or weeks.
7. A single treatment is generally adequate.

#### **CAUTION**

1. If pregnant or a nursing mother, consult your private physician.
2. Elimite is approved for use in children two months of age and older.

## APPENDIX E

### SCABIES FACT SHEET

#### **What is scabies?**

Scabies is an infestation of the skin caused by a mite. The female mite burrows into the top layer of the skin. This forms a slightly raised tunnel where the mite lays eggs and leaves waste.

#### **Who gets scabies?**

Anyone can get scabies.

#### **How is scabies spread?**

The mite is passed from person to person by skin contact or sharing bedding, clothing or other linens with a person who has scabies.

#### **What are the symptoms and when do they appear?**

The most common symptom is an itchy rash. Often the rash itches most at night. It can appear anywhere on the body but is usually on the hands, wrists, elbows, breasts, armpits, waistline, and groin.

Persons who have never had scabies before usually notice symptoms about 4 to 6 weeks after their contact with someone with scabies. Persons who have had scabies before may notice their symptoms sooner, often within a few days to 1 week.

Elderly persons, persons in institutions, and persons whose immune system is weak may not have itching. Any unusual skin problem should be checked by a doctor.

#### **How long is a person able to spread scabies?**

A person is probably able to spread scabies from the moment of contact until after all treatment is complete.

#### **How is scabies diagnosed?**

Scabies is diagnosed by a doctor or nurse looking at the rash and/or by taking a scraping from the skin.

#### **What is the treatment?**

A medicated cream will be prescribed by your doctor. It is put on the skin, left on for several hours, and then washed off. You must put on clean clothes and use freshly laundered bed and bath linens. An oral medication may also be prescribed.

#### **Can a person get scabies again?**

Yes. In fact, the symptoms (itching and rash) will appear more quickly.

#### **Should infested persons be excluded from school or work?**

Yes, until treatment has been finished, but generally this is less than one day.

#### **What are the health problems associated with scabies?**

Usually none. Occasionally, secondary skin infections may occur from scratching.

**What can be done to prevent its spread?**

Persons with symptoms should be checked and treated by their doctor as quickly as possible. Household members and other persons with skin-to-skin contact should be preventively treated. Clothing, bedding, and bath linens used within the 4 days before initiation of therapy should be washed in a washer using hot water and dried using the hot drier cycle. Clothing and other items that cannot be laundered should be stored in a closed plastic bag for one week.

## APPENDIX F

### ISOLATION AND ENVIRONMENTAL CONTROL FOR PATIENTS WITH SCABIES

#### A. Typical Scabies

1. Place patients with typical scabies on contact precautions during the treatment period; 24 hours after application of 5% permethrin cream or 24 hours after last application of scabicides requiring more than one application.
2. HCWs must wear gloves and a long-sleeved gown for hands-on contact. Wash hands after removal of gloves.
3. Place bed linens, towels and clothing used by an affected person during the 4 days prior to initiation of treatment in plastic bags inside the patient's room, handled by gloved and gowned laundry workers without sorting, and washed in hot water for at least 10 minutes. The hot cycle of the dryer should be used for at least 10-20 minutes. Nonwashable blankets and articles can be placed in a plastic bag for 7 days, dry cleaned or tumbled in a hot dryer for 20 minutes.
4. Change all bed linens, towels and clothes daily.
5. Disinfect multiple patient-use items, such as walking belts, blood pressure cuffs, stethoscopes, wheelchairs, etc., before using on other patients. Discard all creams, lotions or ointments used prior to effective treatment.
6. Vacuum mattresses, upholstered furniture and carpeting. There is no need for special treatment of furniture, mattresses or rugs or fumigation of areas. General cleaning and thorough vacuuming is recommended.
7. Routine disinfection procedures are adequate.
8. Symptomatic employees should be allowed back to work the morning following overnight treatment with 5% permethrin cream. Disposable gloves should be worn 2-3 days by symptomatic staff who most provide extensive hands-on care to their patients.

## **B. Crusted (Atypical )Scabies**

(Maintain contact isolation until treatment is completed and/or case is determined by dermatology consultant or other experienced designee to be non-infectious).

1. Assign patient to a private room. Restrict visitors until treatment regimen completed; alternatively, require visitors to gown and glove as required for contact isolation precautions. If resources permit, cohort employees to care for this patient only (no other direct care responsibilities) until effective treatment is completed.
2. HCWs must wear gloves and a long-sleeved gown with the wrist area covered and shoe covers to attend to patient needs, for housekeeping duties, and handling of laundry. Consider spraying pyrethrin insect repellent to wrist (edge of glove and ribbing of sleeve area), arms and front of gown. Remove gown before leaving the room. Wash hands.
3. Bed linens, towels and clothing used by the affected persons during the 4 days prior to initiation of treatment should be placed in plastic bags inside the patient's room, handled by gloved and gowned laundry workers without sorting, and laundered in hot water for at least 10 minutes. The hot cycle of the dryer should be used for at least 10-20 minutes. Non-washable blankets and articles can be placed in a plastic bag for 7 days, dry cleaned or tumbled in a hot dryer for 20 minutes.
4. Change all bed linens, towels and clothes daily.
5. Blood pressure cuffs, walking belts, stethoscopes, etc. should be designated for single patient use and left in the patient's room. Discard all creams, lotions or ointments used prior to effective treatment.
6. Upholstered furniture containing any cloth fabric should be removed from the room and, if necessary, replaced with plastic or vinyl furniture. Mattresses must be covered with plastic or vinyl.
7. The patient's room should be vacuumed daily with a vacuum cleaner designated for this room alone, followed by routine room cleaning and disinfection. The vacuum cleaner bag should be changed daily; removal and disposal of contaminated bags should be performed in accordance with infection control protocol.
8. The room should be terminally cleaned upon discharge or upon transfer of the patient from the room.

**APPENDIX G**

**SCABIES CASE/CONTACT LINE LIST FORM: EMPLOYEES\***

Submitted by (Name & Title): \_\_\_\_\_

Date: \_\_\_/\_\_\_/\_\_\_

Facility: \_\_\_\_\_

Outbreak#: \_\_\_\_\_

Employee Name	Job Title	Location/Description of Rash	Date of Symptom Onset	Dx'd/Eval. By (Name)	Usual Work Assign	Skin Scraping	Scabicide Dates of Rx	Follow-Up/Dates, Results	Family Members Sx	Family Members Rx'd

\*Includes employees, family and other non-patient contacts

**APPENDIX H**

**SCABIES CASE/CONTACT LINE LIST FORM: PATIENTS**

Submitted by (Name & Title): \_\_\_\_\_

Date: \_\_\_/\_\_\_/\_\_\_

Facility: \_\_\_\_\_

Outbreak #: \_\_\_\_\_

Patient Name	Age/ Sex	Medical Record Number	Adm. Date	Dx Date	Curr. Unit/ Bed No.	Dates Exposed - Exposure Period	Bed/Unit since Exposed	Scabies Signs/ Symp.	Dr.'s Name / Date Evaluated	Skin Scraping Results	Scabicide Rx/ Dates

## APPENDIX I

### Sample Letter to Discharged Patients

Facility Letterhead

Date

Dear Patient:

During your recent hospitalization you may have been exposed to scabies. Although it is unlikely that the exposure will result in you becoming infected with scabies, we want to alert you to the possible exposure. We are working in collaboration with Los Angeles County Public Health and want to ensure that you are informed of the possible exposure and the appropriate follow-up steps are taken, if needed.

Scabies is a contagious skin condition caused by a mite that requires skin-to-skin contact. Signs and symptoms include a rash and itching, especially at night. It can appear anywhere on the body but is usually on the hands, wrists, elbows, breasts, armpits, waistline, and groin. Scabies has a long incubation period (the time from possible exposure to the time symptoms develop), usually four to six weeks.

Persons who have scabies are infectious to others, even before symptoms develop. Scabies is diagnosed by a physician or nurse looking at the rash and/or taking a scraping from the skin. Usually, scabies is easily treated by a medicated cream or oral medication prescribed by your physician.

Public Health recommends that you check your skin daily and if a rash and/or itching occurs, notify your personal physician as soon as possible, or you may call me at \_\_\_\_\_ if you have any questions.

Thank you for your cooperation.

Sincerely,

Name, Title

## APPENDIX J

### Sample Letter to Physicians of Discharged Patients

Facility Letterhead

Date

Dear Physician:

Our medical center is currently experiencing an outbreak of scabies. Attached is a list of your patients who may have been exposed. The period of potential exposure was from \_\_\_\_\_ to \_\_\_\_\_.

Patients whose last date of exposure was more than six weeks ago should be questioned regarding symptoms consistent with scabies. Those patients found to be symptomatic should be evaluated and treated as necessary. Family members and other close contacts should receive prophylactic scabicide therapy at the same time the patient is treated. Permethrin 5% cream (*Elimite*) is now the recommended agent for treatment of scabies.

Patients whose last date of exposure was less than six weeks from this date should be treated with topical scabicide or monitored for symptoms until the six week period is over. Close contacts of symptomatic patients should also receive treatment or monitoring as well.

Enclosed is a list of your patients who are considered exposed and their last date of exposure. Please notify \_\_\_ at \_\_\_ if any of your patients develop scabies. Thank you for your assistance.

Sincerely,

Name, Title

## Appendix K

### Scabies Outbreak Management Checklist

<b>Outbreak Interventions</b>				
<b>Communication</b>	<b>N/A</b>	<b>Completed</b>	<b>Date</b>	<b>Signature</b>
Notification of facility administration				
Notification of infection prevention and control team				
Outbreak reported to the local health officer – LAC Acute Communicable Disease Control Program				
Outbreak reported to CDPH Licensing & Certification local office – LAC Health Facilities Division				
Notification of patients and relatives/visitors				
Health facility closed to new admissions				
Health facility reopened to new admissions				
<b>Investigation and Monitoring</b>	<b>N/A</b>	<b>Completed</b>	<b>Date</b>	<b>Signature</b>
Symptomatic health care workers removed from work and referred to Employee Health and/or evaluated by a clinician				
Patients evaluated for scabies and placed in contact isolation				
Patient line list completed				
Employee line list completed				
Confirm confirmed skin scraping in symptomatic patients/employees				
Daily skin assessments documented on all patients				
<b>Treatment</b>	<b>N/A</b>	<b>Completed</b>	<b>Date</b>	<b>Signature</b>
Symptomatic patients/employees treated with scabicide (refer to Appendix B, C & D)				
Prophylactic treatment offered to staff				
Prophylactic treatment offered to family/visitors				
<b>Environmental Cleaning</b>	<b>N/A</b>	<b>Completed</b>	<b>Date</b>	<b>Signature</b>
Enhanced environmental cleaning done throughout the outbreak period				
<b>Education</b>	<b>N/A</b>	<b>Completed</b>	<b>Date</b>	<b>Signature</b>
Training provided to all staff on the signs and symptoms of scabies				

## References

- Sargent SJ. Ectoparasites. In Mayhall CG (ed): Hospital Epidemiology and Infection Control, Third Edition. Baltimore: Williams & Wilkins 2004:755-757.
- Schultz MW, Gomez M, Hansen RC, et al. Comparative study of 5% permethrin cream and 1% lindane lotion for the treatment of scabies. Arch Dermatol 1990; 126:167-170.
- Green MS. Epidemiology of scabies. Epidemiologic Reviews 1989; 11:126-150.
- Lettau LA. Nosocomial transmission and infection control aspects of parasitic and ectoparasitic diseases Part III. Ectoparasites/summary and conclusions. Infect Control Hosp Epidemiol 1991; 12:179-185.
- Carslaw RW, Dobson RM, Hood AJK, Taylor RN. Mites in the environment of cases of Norwegian scabies. Brit J Derm 1975;92:333-7.
- Arlian LG, Estes SA, Vyszynski-Moher MS. Prevalence of *Sarcoptes scabiei* in the homes and nursing homes of scabietic patients. J Am Acad Dermatol 1988;11:242-44.
- California Department of Public Health Division of Communicable Disease Control in Conjunction with Licensing and Certification. Prevention and Control of Scabies in California Long-Term Care Facilities. March 2008.
- Missouri Bureau of Communicable Disease Control. Guidelines for Scabies Prevention and Control. Missouri Epidemiologist. November-December 1994
- Siegel JD, Rhinehart E, Jackson M, Chiarello L, and the Healthcare Infection Control Practices Advisory Committee, 2007 Guideline for Isolation Precautions: Preventing Transmission of Infectious Agents in Healthcare Settings, June 2007  
<http://www.cdc.gov/ncidod/dhqp/pdf/isolation2007.pdf>
- Meinking TL, Taplin D. Safety of permethrin vs lindane for the treatment of scabies. Arch Dermatol 1996;132:959-62.
- Purvis RS, Tyring SK. An outbreak of lindane-resistant scabies treated successfully with permethrin 5% cream. J Am Acad Dermatol 1991;25:1015-6.
- Drugs for Parasitic Infections. The Medical Letter 2007;5(Suppl):e1-e15.
- Meinking TL, Taplin D, Hermida JL, Pardo R, Kerdel FA. The treatment of scabies with ivermectin. N Engl J Med 1995;333:26-30.
- Brown S, Becher J, Brady W. Treatment of ectoparasitic infections: Review of the English-language literature, 1982-1992. Clin Infect Dis 1995;2-(S):S104-9.

Barkwell R, Shields S. Death associated with ivermectin treatment of scabies. *Lancet* 1997;34:1145-45.

Haag ML, Brozena SJ, Fenske NA. Attack of the scabies: What to do when an outbreak occurs. *Geriatrics* 1993;48:45-53.

Wilson, BB. Scabies, In: Mandell GL, Douglas RG, Bennett JF eds. *Principles and Practice of Infectious Diseases*, New York; Churchill Livingstone 2005:3304-3307.

## Acknowledgements

The Los Angeles County Department of Public Health would like to thank Rachael K. Lee RN, MSN/MPH, CNS and Carol Peterson, MD, MPH for developing the *Guidelines for the Prevention and Control of Scabies in Los Angeles County Health Care Facilities, March 2006* upon which the current guidelines are based.

## Revisions

The current revision and expansion of the *Scabies Prevention and Control Guidelines for Acute and Sub-Acute Care Facilities* was prepared by the Acute Communicable Disease Control Program, in conjunction with Community Health Services Division.

L'Tanya English, RN, MPH  
Acute Communicable Disease Control

Dawn Terashita Gastelum, MD, MPH  
Acute Communicable Disease Control

Patricia Marquez, MPH  
Acute Communicable Disease Control

Lorraine Sisneros, RN, MSN/MPH  
Acute Communicable Disease Control

Rachel Civen, MD, MPH  
Acute Communicable Disease Control

Juliet Bugante, RN, BSN  
Acute Communicable Disease Control

Kelly Ivie, RN, BSN  
Community Health Services Division

Paula Marin, RN, BSN  
Community Health Services Division

Laurene Mascola, MD, MPH  
Acute Communicable Disease Control

David Dassey, MD, MPH  
Acute Communicable Disease Control