


COVID-19 VACCINE SCHEDULES

People Ages 12+

(If you have a weak immune system, see *Immunocompromised People 12+*)



¹ Timing of 2nd dose: People under 64 years, and especially males ages 12-39, may consider getting the 2nd primary dose of vaccine 8 weeks after the 1st dose. A longer time between the 1st and 2nd doses may increase how much protection the vaccines offer and reduce the rare risk of heart problems (myocarditis and pericarditis). 3 weeks (Pfizer/Novavax) or 4 weeks (Moderna) is recommended for those age 65+ and anyone more likely to get very sick from COVID-19, or who needs the fullest possible protection sooner. Talk with your doctor if you are not sure about the best timing for your situation. See [Stay Up to Date with Your COVID-19 Vaccines Including Boosters](#).

² In limited situations, people ages 18 and older may choose to receive a Novavax vaccine as a booster instead of an mRNA (Pfizer or Moderna) vaccine: if they are unable or unwilling to receive an mRNA booster AND they have completed primary series vaccination AND they have never had a booster before. The Novavax booster is given at least 6 months after completing the last dose of the primary series.


³ J&J vaccine should only be used in [some situations](#).

If you have COVID-19, wait until you have recovered (if you had symptoms) and finished isolation to get any COVID-19 vaccine doses. If you recently had COVID-19, you may consider delaying a vaccine dose (primary series or booster) by 3 months from when your symptoms started or your positive test (if you didn't have any symptoms).



COVID-19 VACCINE SCHEDULES

Immunocompromised People Ages 12+



- 1 In limited situations, people ages 18 and older may choose to receive a Novavax vaccine as a booster instead of an mRNA (Pfizer or Moderna) vaccine: if they are unable or unwilling to receive an mRNA booster AND they have completed primary series vaccination AND they have never had a booster before. The Novavax booster is given at least 6 months after completing the last dose of the primary series.
 - 2 J&J vaccine should only be used in [some situations](#).
- **If you have COVID-19**, wait until you have recovered (if you had symptoms) and finished isolation to get a vaccine dose.
 - **You may also be eligible for a monoclonal antibody medicine called [Evusheld](#)** if you have a [moderately or severely weakened immune system](#) (e.g., a blood-related cancer, organ transplant, certain autoimmune condition, or certain treatment for cancer). It is given by injection every six months to help prevent you from getting COVID-19. It is for people ages 12+ who weigh at least 88 lbs. Evusheld can be given at least 2 weeks after any COVID-19 vaccine. Talk with your doctor about Evusheld and the best timing of vaccine doses based on your risks and treatment plan.