






COVID-19 VACCINE SCHEDULES

Children Ages 6 months – 11 years

(If your child has a weak immune system, see *Immunocompromised Children*)

Age	Vaccine	Primary Series	Updated Booster
6 months - 4 years	Pfizer		Booster not currently recommended
6 months - 4 years	Moderna		Booster not currently recommended
5-11 years	Pfizer		<p>Children age 5 get the Pfizer booster.</p> <p>Children ages 6+ get either the Moderna or Pfizer booster.</p>
5- 11 years	Moderna		



* The updated booster is recommended at least 2 months after the last COVID-19 vaccine (either the final primary series dose or last booster).

- **COVID-19 vaccine dosage is based on your child's age on the day of vaccination, not on their size or weight.** Children ages 11 and under get a lower dose of vaccine than teens and adults.
- **COVID-19 vaccines and other routine childhood vaccines can be given at the same visit.**
- **Timing of 2nd dose:** 8 weeks may be best for some people because a longer time between the 1st and 2nd doses may increase how much protection the vaccines offer and reduce the rare risk of heart problems (myocarditis and pericarditis). A 3-week (Pfizer) or 4-week (Moderna) interval is recommended for those people more likely to get very sick from COVID-19 or anyone who needs the fullest possible protection sooner. Talk with your doctor if you are not sure about the best timing for your situation.
- **If your child already had COVID-19, they should still get vaccinated.** Wait until they have recovered (if they had symptoms) and finished isolation to get any vaccine, including a COVID-19 vaccine. If they recently had COVID-19, you may consider delaying a COVID-19 vaccine dose (primary series or booster) by 3 months from when their symptoms started or when they first had a positive test (if they didn't have any symptoms).
- **For more information,** see the CDC webpage, [Stay Up to Date with Your COVID-19 Vaccines Including Boosters](https://www.cdc.gov/covid19/vaccines-including-boosters).

COVID-19 VACCINE SCHEDULES

Immunocompromised Children Ages 6 months – 11 years

Age	Vaccine	Primary Series	Updated Booster
6 months - 4 years	Pfizer		Booster not currently recommended
6 months - 4 years	Moderna		Booster not currently recommended
5-11 years	Pfizer		<p>Children age 5 get the Pfizer booster.</p> <p>Children ages 6+ get either Moderna or Pfizer booster.</p>
5-11 years	Moderna		

*The updated booster is recommended at least 2 months after the last COVID-19 vaccine (either the final primary series dose or last booster).

- **COVID vaccine dosage is based on age on the day of vaccination, not on size or weight.** Children ages 11 and under get a lower dose of vaccine than teens and adults. For more information, see the CDC webpage, [Stay Up to Date with Your COVID-19 Vaccines Including Boosters](#).
- **COVID-19 vaccines and other routine childhood vaccines can be given at the same visit.**
- **If your child already had COVID-19, they should still get vaccinated.** Wait until they have recovered (if they had symptoms) and finished isolation to get any vaccine, including a COVID-19 vaccine dose. If they recently had COVID-19, you may consider delaying a COVID-19 vaccine dose (primary series or booster) by 3 months from when their symptoms started or when they first had a positive test (if they didn't have any symptoms).
- **For more information,** see the CDC webpage, [COVID-19 Vaccines for People who are Moderately or Severely Immunocompromised](#).

