



COUNTY OF LOS ANGELES
Public Health

Sexually Transmitted Disease Program

STD Clinic Morbidity Report

Los Angeles County

2010



2615 South Grand Avenue, Room 500
Los Angeles, CA 90007
Tel: (213) 744-3070 Fax: (213) 749-9606

<http://publichealth.lacounty.gov/std>



JONATHAN E. FIELDING, M.D., M.P.H.
Director and Health Officer

JONATHAN E. FREEDMAN
Chief Deputy Director

PETER R. KERNDT, M.D., M.P.H.
Director, Sexually Transmitted Disease Program
2615 South Grand Avenue, Room 500
Los Angeles, California 90007
TEL (213) 744-3070 • FAX (213) 749-9606
<http://publichealth.lacounty.gov/std>

BOARD OF SUPERVISORS

Gloria Molina
First District

Mark Ridley-Thomas
Second District

Zev Yaroslavsky
Third District

Don Knabe
Fourth District

Michael D. Antonovich
Fifth District

September 22, 2011

Dear Colleague:

I am pleased to release the *Los Angeles County Sexually Transmitted Disease Clinic Morbidity Report, 2010*. This report summarizes chlamydia, gonorrhea, syphilis, HIV and herpes testing in the Los Angeles County Department of Public Health's Sexually Transmitted Disease (STD) clinics. These twelve clinics are Antelope Valley, Central, Curtis R. Tucker, Hollywood/Wilshire, Monrovia, North Hollywood, Pomona, Ruth Temple, Simms Mann Health and Wellness Center, South (now Martin Luther King, Jr. Center for Public Health), Torrance, and Whittier Health Centers.

In 2010, there were nearly 44,000 visits to the health centers for STD services and over 28,000 patients were tested for chlamydia, gonorrhea, syphilis and/or HIV. Chlamydia was the most commonly diagnosed STD (10.9%) followed by gonorrhea (4.7%), HIV (0.8%) and early syphilis (0.5%).

The report is organized with summaries of testing first, followed by sections on individual health centers. New to the 2010 report are summaries of clinical diagnoses, cervical cancer screening and the primary reason for clinic visit, as well as many additional figures to better illustrate disease prevalence, disease trends, and STD co-infections. Gender of sex partner is now included in all the major tables throughout the report.

We hope that this report serves as a resource to better understand STD morbidity in our clinics and serves as a guide in designing population-specific STD prevention and intervention programs for those seeking care at the Los Angeles County Department of Public Health STD clinics.

Sincerely,

A handwritten signature in black ink that reads "Peter R. Kerndt". The signature is fluid and cursive.

Peter R. Kerndt, M.D., M.P.H.
Director, Sexually Transmitted Disease Program
Los Angeles County Department of Public Health

PRK:zz/as

County of Los Angeles Public Health

Jonathan E. Fielding, MD, MPH
Director and Health Officer

Jonathan E. Freedman
Chief Deputy Director

Robert J. Kim-Farley, MD, MPH
Director, Disease Control Programs & Prevention

Deborah Davenport RN, MS
Director, CHS

Nandini Sodhi, MD, MPH
Area Health Officer SPA 1&2

Cristin Mondy, RN,
Acting Area Health Officer SPA 3&4

Jan King, MD, MPH
Area Health Officer SPA 5&6

Silvia Prieto, MD, MPH
Area Health Officer SPA 7&8

Jeffrey D Gunzenhauser, MD, MPH
Medical Director, CHS

Michael Palmer
Public Health Information System

Sexually Transmitted Disease Program

Peter R. Kerndt, MD, MPH
Director

Sarah Guerry, MD
Medical Director

Christine Wigen, MD, MPH
Assistant Medical Director

Ali Stirland, MBChB, MSc
STD Clinic Liaison

Getahun Aynalem, MD, MPH, Michael Chien, MPH
Epidemiology

Kai-Jen Cheng, MS, Waley Zeng, MD, MPH, Teofuldo Cruz
Information Systems & Data Management

Public Health Centers:

Antelope Valley Health Center
335B East Avenue K-6
Lancaster, CA 93535
(661) 723-4526

Elise Pomerance, MD, MPH, Area Medical Director
Angela Lozano, RN, Nurse Manager

Simms Mann Health & Wellness Center
2509 Pico Blvd
Santa Monica, CA 90405
(310) 998-3203

Maxine Liggins, MD, Area Medical Director
Lanet Williams, RN, Nurse Manager

Central Health Center
241 N. Figueroa Street, Los Angeles, CA 90012
(213) 240-8225

Sheree Poitier, MD, Area Medical Director
Rachael Lee, RN, Nurse Manager

Hollywood-Wilshire Health Center
5205 Melrose Avenue, Los Angeles, CA 90038
(323) 769-7932

Sheree Poitier, MD, Area Medical Director
Rachael Lee, RN, Nurse Manager

Monrovia Health Center
330 W. Maple Avenue, Monrovia, CA 91016
(626) 256-1600

Sheree Poitier, MD, Area Medical Director
Leticia Kouchak-Eftekhar, RN, Nurse Manager

North Hollywood Health Center
5300 Tujunga Avenue
North Hollywood, CA 91601
(818) 766-3982
Elise Pomerance, MD, MPH, Area Medical Director
Joanne Hobbs, RN, Nurse Manager

Pomona Health Center
750 S. Park Avenue, Pomona, CA 91766
(909) 868-0235
Sheree Poitier, MD, Area Medical Director
Leticia Kouchak-Eftekhar, RN, Nurse Manager

South Health Center
1522 E. 102nd Street, Los Angeles, CA 90002
(323) 563-4112
Maxine Liggins, MD, Area Medical Director
Lanet Williams, RN, Nurse Manager

Ruth Temple Health Center
3834 S. Western Avenue, Los Angeles CA 90062
(323) 730-3576
Maxine Liggins, MD, Area Medical Director
Lanet Williams, RN, Nurse Manager

Torrance Health Center
711 Del Amo Blvd, Torrance, CA 90501
(310) 222-6571
Meri Rathbun, MD, MPH, Area Medical Director
Barbara Smith, RN, Nurse Manager

Curtis Tucker Health Center
123 W. Manchester Blvd, Inglewood, CA 90301
(310) 419-5362
Meri Rathbun, MD, MPH, Area Medical Director
Barbara Smith, RN, Nurse Manager

Whittier Health Center
7643 S. Painter Avenue, Whittier CA 90602
(562) 464-5350
Meri Rathbun, MD, MPH, Area Medical Director
Norina Cadena, RN, Nurse Manager

The Sexually Transmitted Disease Clinic Morbidity Report is published on a yearly basis by the Sexually Transmitted Disease Program of the Los Angeles County Department of Public Health. This report is also available in PDF format, at <http://publichealth.lacounty.gov/std/>

If you would like to receive surveillance reports and other information from the STD Program, register for the STDInfo listserv. To subscribe send an email to: LISTSERV@LISTSERV.PH.LACOUNTY.GOV and type SUBSCRIBE STDINFO anywhere in the body of the email.

SUGGESTED CITATION

Sexually Transmitted Disease Program. Los Angeles County Department of Public Health. STD Clinic Morbidity Report, Los Angeles County, 2010: pp I-1--XIV-8.

CONTENTS

I. All Public Health Centers

Overall

| | | |
|------------|---|-------|
| Figure 1.1 | Map of Los Angeles County Public Health Centers | I-1 |
| Table 1.1 | Summary Visit Information by Health Center | I-2,3 |
| Table 1.2 | Summary Morbidity Information | I-4 |
| Figure 1.2 | STD Prevalence in Public Health Centers, 2010 | I-5 |
| Table 1.3 | Patient Visits and Tests | I-6 |

By Gender, Age and Race/Ethnicity

| | | |
|-------------|---|------|
| Figure 1.3 | Primary Reason(s) for New Visits among Males | I-7 |
| Figure 1.4 | Primary Reason(s) for New Visits among Females | I-8 |
| Table 1.4 | Patient Visits and Tests by Gender and Age | I-9 |
| Table 1.5 | Patient Visits and Tests by Gender and Race/Ethnicity | I-10 |
| Figure 1.5 | Chlamydia Cases Among Males, by Age Group and Race/Ethnicity | I-11 |
| Figure 1.6 | Chlamydia Cases Among Females, by Age Group and Race/Ethnicity | I-12 |
| Figure 1.7 | Gonorrhea Cases Among Males, by Age Group and Race/Ethnicity | I-13 |
| Figure 1.8 | Gonorrhea Cases Among Females, by Age Group and Race/Ethnicity | I-14 |
| Figure 1.9 | Early Syphilis Cases Among Males, by Age Group and Race/Ethnicity | I-15 |
| Figure 1.10 | HIV Cases Among Males, by Age Group and Race/Ethnicity | I-16 |
| Figure 1.11 | New HIV Cases Among Males, by Age Group and Race/Ethnicity | I-17 |
| Figure 1.12 | New HIV Cases Among Females, by Age Group and Race/Ethnicity | I-18 |

By Trend

| | | |
|--------------|---|------|
| Figure 1.13a | Sexually Transmitted Disease Prevalence, 2006-2010 | I-19 |
| Figure 1.13b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | I-20 |
| Figure 1.14 | Chlamydia Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-21 |
| Figure 1.15 | Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-22 |
| Figure 1.16 | Early Syphilis Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-23 |
| Figure 1.17 | HIV Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-24 |
| Figure 1.18 | New HIV Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-25 |

By Site of Infection

| | | |
|-------------|--|---------|
| Table 1.6 | Site of Chlamydia Infection by Gender | I-26,27 |
| Table 1.7 | Site of Gonorrhea Infection by Gender | I-28,29 |
| Figure 1.19 | Urogenital Chlamydia Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-30 |
| Figure 1.20 | Rectal Chlamydia Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-31 |
| Figure 1.21 | Urethral Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-32 |
| Figure 1.22 | Pharyngeal Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-33 |
| Figure 1.23 | Rectal Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010 | I-34 |
| Figure 1.24 | Distribution of Anatomic Site of Infection among All Patients with Chlamydia Infection | I-35 |
| Figure 1.25 | Distribution of Anatomic Site of Infection among All Patients with Gonorrhea Infection | I-36 |

| | | |
|-------------|---|------|
| Figure 1.26 | Distribution of Anatomic Site of Infection among MSM/W Patients with Chlamydia Infection | I-37 |
| Figure 1.27 | Distribution of Anatomic Site of Infection among MSM/W Patients with Gonorrhea Infection | I-38 |
| Figure 1.28 | Distribution of Anatomic Site of Infection among Female Patients with Chlamydia Infection | I-39 |
| Figure 1.29 | Distribution of Anatomic Site of Infection among Female Patients with Gonorrhea Infection | I-40 |

Concurrent New HIV and other STD Diagnosis

| | | |
|-------------|---|------|
| Table 1.8 | Patients Diagnosed with New HIV and Other STD on Same Visit | I-41 |
| Table 1.9 | Patients Diagnosed with New HIV and Other STD on Same Visit, by Gender and Age | I-42 |
| Table 1.10 | Patients Diagnosed with New HIV and Other STD on Same Visit, by Gender and Race/Ethnicity | I-43 |
| Figure 1.30 | Other STDs Diagnosed on Same Visit as HIV among Patients with New HIV | I-44 |
| Figure 1.31 | Distribution of Other STDs Diagnosed on Same Visit as HIV among Patients with New HIV | I-45 |
| Figure 1.32 | Distribution of Anatomic Site of Chlamydia Infection Diagnosed on Same Visit as HIV among Patients with New HIV | I-46 |
| Figure 1.33 | Distribution of Anatomic Site of Gonorrhea Infection Diagnosed on Same Visit as HIV among Patients with New HIV | I-47 |
| Figure 1.34 | Distribution of Other STDs Diagnosed on Same Visit as HIV among MSM/W Patients with New HIV | I-48 |
| Figure 1.35 | Distribution of Anatomic Site of Chlamydia Infection Diagnosed on Same Visit as HIV among MSM/W Patients with New HIV | I-49 |
| Figure 1.36 | Distribution of Anatomic Site of Gonorrhea Infection Diagnosed on Same Visit as HIV among MSM/W Patients with New HIV | I-50 |

Herpes Simplex Virus Testing

| | | |
|------------|--|------|
| Table 1.11 | Herpes Simplex Virus Testing of Symptomatic Patients Summary by Gender | I-51 |
|------------|--|------|

Paps Testing

| | | |
|------------|--|---------|
| Table 1.12 | Summary of Paps Performed by Health Center | I-52,53 |
|------------|--|---------|

Other Diagnoses

| | | |
|-------------|--|------|
| Figure 1.37 | Clinical Diagnoses among Males at New Visits | I-54 |
|-------------|--|------|

| | | |
|-------------|--|------|
| Figure 1.38 | Clinical Diagnoses among Females at New Visits | I-55 |
|-------------|--|------|

II. Antelope Valley Health Center

| | | |
|-----------|--------------------------|------|
| Table 2.1 | Patient Visits and Tests | II-1 |
|-----------|--------------------------|------|

| | | |
|-----------|--|------|
| Table 2.2 | Patient Visits and Tests by Gender and Age | II-2 |
|-----------|--|------|

| | | |
|-----------|---|------|
| Table 2.3 | Patient Visits and Tests by Gender and Race/Ethnicity | II-3 |
|-----------|---|------|

| | | |
|-------------|--|------|
| Figure 2.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | II-4 |
|-------------|--|------|

| | | |
|-------------|--|------|
| Figure 2.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | II-5 |
|-------------|--|------|

III. Simms Mann Health & Wellness Center

| | | |
|-----------|--------------------------|-------|
| Table 3.1 | Patient Visits and Tests | III-1 |
|-----------|--------------------------|-------|

| | | |
|-----------|--|-------|
| Table 3.2 | Patient Visits and Tests by Gender and Age | III-2 |
|-----------|--|-------|

| | | |
|-----------|---|-------|
| Table 3.3 | Patient Visits and Tests by Gender and Race/Ethnicity | III-3 |
|-----------|---|-------|

| | | |
|-------------|--|-------|
| Figure 3.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | III-4 |
|-------------|--|-------|

| | | |
|-------------|--|-------|
| Figure 3.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | III-5 |
|-------------|--|-------|

IV. Central Health Center

| | | |
|-----------|--------------------------|------|
| Table 4.1 | Patient Visits and Tests | IV-1 |
|-----------|--------------------------|------|

| | | |
|-----------|--|------|
| Table 4.2 | Patient Visits and Tests by Gender and Age | IV-2 |
|-----------|--|------|

| | | |
|-----------|---|------|
| Table 4.3 | Patient Visits and Tests by Gender and Race/Ethnicity | IV-3 |
|-----------|---|------|

Figure 4.1a Sexually Transmitted Disease Prevalence, 2006-2010 IV-4

Figure 4.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 IV-5

V. Hollywood-Wilshire Health Center

Table 5.1 Patient Visits and Tests V-1

Table 5.2 Patient Visits and Tests by Gender and Age V-2

Table 5.3 Patient Visits and Tests by Gender and Race/Ethnicity V-3

Figure 5.1a Sexually Transmitted Disease Prevalence, 2006-2010 V-4

Figure 5.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 V-5

VI. Monrovia Health Center

Table 6.1 Patient Visits and Tests VI-1

Table 6.2 Patient Visits and Tests by Gender and Age VI-2

Table 6.3 Patient Visits and Tests by Gender and Race/Ethnicity VI-3

Figure 6.1a Sexually Transmitted Disease Prevalence, 2006-2010 VI-4

Figure 6.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 VI-5

VII. North Hollywood Health Center

Table 7.1 Patient Visits and Tests VII-1

Table 7.2 Patient Visits and Tests by Gender and Age VII-2

Table 7.3 Patient Visits and Tests by Gender and Race/Ethnicity VII-3

Figure 7.1a Sexually Transmitted Disease Prevalence, 2006-2010 VII-4

Figure 7.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 VII-5

VIII. Pomona Health Center

Table 8.1 Patient Visits and Tests VIII-1

Table 8.2 Patient Visits and Tests by Gender and Age VIII-2

| | | |
|-------------|--|--------|
| Table 8.3 | Patient Visits and Tests by Gender and Race/Ethnicity | VIII-3 |
| Figure 8.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | VIII-4 |
| Figure 8.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | VIII-5 |

IX. South Health Center

| | | |
|-------------|--|------|
| Table 9.1 | Patient Visits and Tests | IX-1 |
| Table 9.2 | Patient Visits and Tests by Gender and Age | IX-2 |
| Table 9.3 | Patient Visits and Tests by Gender and Race/Ethnicity | IX-3 |
| Figure 9.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | IX-4 |
| Figure 9.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | IX-5 |

X. Ruth Temple Health Center

| | | |
|--------------|--|-----|
| Table 10.1 | Patient Visits and Tests | X-1 |
| Table 10.2 | Patient Visits and Tests by Gender and Age | X-2 |
| Table 10.3 | Patient Visits and Tests by Gender and Race/Ethnicity | X-3 |
| Figure 10.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | X-4 |
| Figure 10.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | X-5 |

XI. Torrance Health Center

| | | |
|--------------|--|------|
| Table 11.1 | Patient Visits and Tests | XI-1 |
| Table 11.2 | Patient Visits and Tests by Gender and Age | XI-2 |
| Table 11.3 | Patient Visits and Tests by Gender and Race/Ethnicity | XI-3 |
| Figure 11.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | XI-4 |
| Figure 11.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | XI-5 |

XII. Curtis Tucker Health Center

| | | |
|------------|--------------------------|-------|
| Table 12.1 | Patient Visits and Tests | XII-1 |
|------------|--------------------------|-------|

| | | |
|--------------|--|-------|
| Table 12.2 | Patient Visits and Tests by Gender and Age | XII-2 |
| Table 12.3 | Patient Visits and Tests by Gender and Race/Ethnicity | XII-3 |
| Figure 12.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | XII-4 |
| Figure 12.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | XII-5 |

XIII. Whittier Health Center

| | | |
|--------------|--|--------|
| Table 13.1 | Patient Visits and Tests | XIII-1 |
| Table 13.2 | Patient Visits and Tests by Gender and Age | XIII-2 |
| Table 13.3 | Patient Visits and Tests by Gender and Race/Ethnicity | XIII-3 |
| Figure 13.1a | Sexually Transmitted Disease Prevalence, 2006-2010 | XIII-4 |
| Figure 13.1b | Early Syphilis, HIV, and New HIV Prevalence, 2006-2010 | XIII-5 |

XIV. Appendices

| | | |
|-------------|---|-----------|
| Appendix A. | Laboratory Diagnostic Tests Used for the Diagnosis of Sexually Transmitted Diseases | XIV-1 |
| Appendix B. | Data Management | XIV-2,3 |
| Appendix C | STD Clinic Registration Schedule, June 2010 | XIV-4,5 |
| Appendix D | STD Clinic Intake Form (H2541) | XIV-6,7,8 |

Sexually Transmitted Disease Clinic Morbidity Report, 2010

I: All Public Health Centers

Figure 1. Map of Los Angeles County Public Health Centers (HC)



Table 1.1 Summary Visit Information by Health Center

| | All Clinics | | Antelope Valley | | Simms Mann | | Central | | Hollywood/ Wilshire | | North Hollywood | | Monrovia | |
|---------------------------------|-------------|-------|-----------------|-------|------------|-------|---------|-------|------------------------|-------|-----------------|-------|----------|-------|
| | n* | % | n | % | n | % | n | % | n | % | n | % | n | % |
| Total | 43,973 | 100.0 | 1,613 | 100.0 | 806 | 100.0 | 5,882 | 100.0 | 6,205 | 100.0 | 5,596 | 100.0 | 1,222 | 100.0 |
| Gender | | | | | | | | | | | | | | |
| Male | 28,441 | 64.7 | 740 | 45.9 | 624 | 77.4 | 4,175 | 71.0 | 4,684 | 75.5 | 3,976 | 71.1 | 813 | 66.5 |
| Female | 15,424 | 35.1 | 873 | 54.1 | 181 | 22.5 | 1,678 | 28.5 | 1,480 | 23.9 | 1,604 | 28.7 | 406 | 33.2 |
| Transgender† | 108 | 0.2 | 0 | 0.0 | 1 | 0.1 | 29 | 0.5 | 41 | 0.7 | 16 | 0.3 | 3 | 0.2 |
| Gender of Sex Partners†† | | | | | | | | | | | | | | |
| MSM/W | 4,963 | 11.3 | 83 | 5.1 | 99 | 12.3 | 991 | 16.8 | 1,240 | 20.0 | 854 | 15.3 | 172 | 14.1 |
| MSW | 22,210 | 50.5 | 625 | 38.7 | 510 | 63.3 | 2,956 | 50.3 | 3,223 | 51.9 | 2,888 | 51.6 | 592 | 48.4 |
| WSM | 13,416 | 30.5 | 770 | 47.7 | 174 | 21.6 | 1,485 | 25.2 | 1,272 | 20.5 | 1,400 | 25.0 | 359 | 29.4 |
| WSW/M | 1,104 | 2.5 | 78 | 4.8 | 4 | 0.5 | 82 | 1.4 | 110 | 1.8 | 79 | 1.4 | 9 | 0.7 |
| Unknown | 2,280 | 5.2 | 57 | 3.5 | 19 | 2.4 | 368 | 6.3 | 360 | 5.8 | 375 | 6.7 | 90 | 7.4 |
| Age | | | | | | | | | | | | | | |
| 0-14 | 82 | 0.2 | 6 | 0.4 | 29 | 3.6 | 6 | 0.1 | 3 | 0.0 | 6 | 0.1 | 3 | 0.2 |
| 15-19 | 3,636 | 8.3 | 302 | 18.7 | 163 | 20.2 | 273 | 4.6 | 278 | 4.5 | 200 | 3.6 | 71 | 5.8 |
| 20-24 | 9,737 | 22.1 | 594 | 36.8 | 205 | 25.4 | 1,016 | 17.3 | 1,189 | 19.2 | 1,022 | 18.3 | 277 | 22.7 |
| 25-29 | 8,691 | 19.8 | 282 | 17.5 | 153 | 19.0 | 1,124 | 19.1 | 1,412 | 22.8 | 1,248 | 22.3 | 248 | 20.3 |
| 30-34 | 6,272 | 14.3 | 149 | 9.2 | 165 | 20.5 | 913 | 15.5 | 1,079 | 17.4 | 997 | 17.8 | 135 | 11.0 |
| 35-44 | 8,123 | 18.5 | 165 | 10.2 | 68 | 8.4 | 1,279 | 21.7 | 1,284 | 20.7 | 1,224 | 21.9 | 252 | 20.6 |
| 45-54 | 5,148 | 11.7 | 83 | 5.1 | 21 | 2.6 | 813 | 13.8 | 682 | 11.0 | 555 | 9.9 | 140 | 11.5 |
| 55-64 | 1,798 | 4.1 | 30 | 1.9 | 2 | 0.2 | 341 | 5.8 | 233 | 3.8 | 270 | 4.8 | 56 | 4.6 |
| 65+ | 486 | 1.1 | 2 | 0.1 | 0 | 0.0 | 117 | 2.0 | 45 | 0.7 | 74 | 1.3 | 40 | 3.3 |
| Race/Ethnicity | | | | | | | | | | | | | | |
| White | 6,935 | 15.8 | 304 | 18.8 | 454 | 56.3 | 694 | 11.8 | 2,028 | 32.7 | 2,002 | 35.8 | 186 | 15.2 |
| African American | 17,464 | 39.7 | 690 | 42.8 | 116 | 14.4 | 1,068 | 18.2 | 1,224 | 19.7 | 807 | 14.4 | 117 | 9.6 |
| Hispanic | 15,942 | 36.3 | 438 | 27.2 | 103 | 12.8 | 3,690 | 62.7 | 2,080 | 33.5 | 2,180 | 39.0 | 588 | 48.1 |
| Asian | 1,890 | 4.3 | 28 | 1.7 | 84 | 10.4 | 120 | 2.0 | 528 | 8.5 | 317 | 5.7 | 253 | 20.7 |
| Native American | 109 | 0.2 | 3 | 0.2 | 2 | 0.2 | 11 | 0.2 | 16 | 0.3 | 25 | 0.4 | 3 | 0.2 |
| Other | 1,410 | 3.2 | 119 | 7.4 | 37 | 4.6 | 275 | 4.7 | 319 | 5.1 | 238 | 4.3 | 68 | 5.6 |
| Unknown | 223 | 0.5 | 31 | 1.9 | 10 | 1.2 | 24 | 0.4 | 10 | 0.2 | 27 | 0.5 | 7 | 0.6 |
| Type of Visit | | | | | | | | | | | | | | |
| Clinic session | 3,347 | 100.0 | 104 | 3.1 | 104 | 3.1 | 427 | 12.8 | 392 | 11.7 | 435 | 13.0 | 148 | 4.4 |
| New Visits per Session††† | 8.2 | -- | 10.5 | -- | 6.6 | -- | 6.9 | -- | 9.1 | -- | 7.3 | -- | 5.3 | -- |
| New Visit | 27,497 | 62.5 | 1,097 | 68.0 | 684 | 84.9 | 2,965 | 50.4 | 3,567 | 57.5 | 3,185 | 56.9 | 785 | 64.2 |
| Return Visit | 16,476 | 37.5 | 516 | 32.0 | 122 | 15.1 | 2,917 | 49.6 | 2,638 | 42.5 | 2,411 | 43.1 | 437 | 35.8 |

*Number of visits

†Transgendered in total visits for all clinics include 36 male-to-female (MTF) and 72 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women;

"WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

†††Average number of new visits per session i.e. total number of new visits 2010 / total number of sessions 2010. See data section (Appendix B) for definition of a new visit. A session is a half-day when the STD clinic is open.

Table 1.1 (continued) Summary Visit Information by Health Center

| | All Clinics | | Pomona | | South | | R. Temple | | Torrance | | C. Tucker | | Whittier | |
|---------------------------------|-------------|-------|--------|-------|-------|-------|-----------|-------|----------|-------|-----------|-------|----------|-------|
| | n* | % | n | % | n | % | n | % | n | % | n | % | n | % |
| Total | 43,973 | 100.0 | 1,900 | 100.0 | 6,713 | 100.0 | 5,471 | 100.0 | 793 | 100.0 | 5,608 | 100.0 | 2,164 | 100.0 |
| Gender | | | | | | | | | | | | | | |
| Male | 28,441 | 64.7 | 1,265 | 66.6 | 3,554 | 52.9 | 3,176 | 58.1 | 555 | 70.0 | 3,378 | 60.2 | 1,501 | 69.4 |
| Female | 15,424 | 35.1 | 626 | 32.9 | 3,154 | 47.0 | 2,292 | 41.9 | 238 | 30.0 | 2,229 | 39.7 | 663 | 30.6 |
| Transgender† | 108 | 0.2 | 9 | 0.5 | 5 | 0.1 | 3 | 0.1 | 0 | 0.0 | 1 | 0.0 | 0 | 0.0 |
| Gender of Sex Partners†† | | | | | | | | | | | | | | |
| MSM/W | 4,963 | 11.3 | 330 | 17.4 | 261 | 3.9 | 235 | 4.3 | 79 | 10.0 | 329 | 5.9 | 290 | 13.4 |
| MSW | 22,210 | 50.5 | 893 | 47.0 | 3,173 | 47.3 | 2,849 | 52.1 | 444 | 56.0 | 2,942 | 52.5 | 1,114 | 51.5 |
| WSM | 13,416 | 30.5 | 498 | 26.2 | 2,730 | 40.7 | 2,018 | 36.9 | 181 | 22.8 | 1,951 | 34.8 | 578 | 26.7 |
| WSW/M | 1,104 | 2.5 | 45 | 2.4 | 281 | 4.2 | 176 | 3.2 | 21 | 2.6 | 189 | 3.4 | 30 | 1.4 |
| Unknown | 2,280 | 5.2 | 134 | 7.1 | 268 | 4.0 | 193 | 3.5 | 68 | 8.6 | 197 | 3.5 | 152 | 7.0 |
| Age | | | | | | | | | | | | | | |
| 0-14 | 82 | 0.2 | 9 | 0.5 | 15 | 0.2 | 8 | 0.1 | 2 | 0.3 | 18 | 0.3 | 6 | 0.3 |
| 15-19 | 3,636 | 8.3 | 192 | 10.1 | 923 | 13.7 | 522 | 9.5 | 45 | 5.7 | 604 | 10.8 | 197 | 9.1 |
| 20-24 | 9,737 | 22.1 | 528 | 27.8 | 1,698 | 25.3 | 1,234 | 22.6 | 175 | 22.1 | 1,300 | 23.2 | 541 | 25.0 |
| 25-29 | 8,691 | 19.8 | 380 | 20.0 | 1,108 | 16.5 | 983 | 18.0 | 198 | 25.0 | 1,052 | 18.8 | 451 | 20.8 |
| 30-34 | 6,272 | 14.3 | 213 | 11.2 | 775 | 11.5 | 706 | 12.9 | 102 | 12.9 | 758 | 13.5 | 292 | 13.5 |
| 35-44 | 8,123 | 18.5 | 313 | 16.5 | 1,083 | 16.1 | 909 | 16.6 | 124 | 15.6 | 959 | 17.1 | 366 | 16.9 |
| 45-54 | 5,148 | 11.7 | 187 | 9.8 | 822 | 12.2 | 800 | 14.6 | 84 | 10.6 | 689 | 12.3 | 225 | 10.4 |
| 55-64 | 1,798 | 4.1 | 66 | 3.5 | 226 | 3.4 | 267 | 4.9 | 39 | 4.9 | 184 | 3.3 | 65 | 3.0 |
| 65+ | 486 | 1.1 | 12 | 0.6 | 63 | 0.9 | 42 | 0.8 | 24 | 3.0 | 44 | 0.8 | 21 | 1.0 |
| Race/Ethnicity | | | | | | | | | | | | | | |
| White | 6,935 | 15.8 | 273 | 14.4 | 148 | 2.2 | 135 | 2.5 | 160 | 20.2 | 301 | 5.4 | 250 | 11.6 |
| African American | 17,464 | 39.7 | 304 | 16.0 | 4,761 | 70.9 | 4,312 | 78.8 | 233 | 29.4 | 3,742 | 66.7 | 90 | 4.2 |
| Hispanic | 15,942 | 36.3 | 1,115 | 58.7 | 1,708 | 25.4 | 881 | 16.1 | 261 | 32.9 | 1,250 | 22.3 | 1,648 | 76.2 |
| Asian | 1,890 | 4.3 | 145 | 7.6 | 32 | 0.5 | 61 | 1.1 | 98 | 12.4 | 111 | 2.0 | 113 | 5.2 |
| Native American | 109 | 0.2 | 5 | 0.3 | 15 | 0.2 | 4 | 0.1 | 1 | 0.1 | 14 | 0.2 | 10 | 0.5 |
| Other | 1,410 | 3.2 | 46 | 2.4 | 42 | 0.6 | 72 | 1.3 | 27 | 3.4 | 123 | 2.2 | 44 | 2.0 |
| Unknown | 223 | 0.5 | 12 | 0.6 | 7 | 0.1 | 6 | 0.1 | 13 | 1.6 | 67 | 1.2 | 9 | 0.4 |
| Type of Visit | | | | | | | | | | | | | | |
| Clinic session | 3,347 | 100.0 | 160 | 4.8 | 509 | 15.2 | 449 | 13.4 | 74 | 2.2 | 392 | 11.7 | 153 | 4.6 |
| New Visits per Session††† | 8.2 | -- | 5.8 | -- | 10.1 | -- | 7.5 | -- | 7.6 | -- | 9.6 | -- | 9.6 | -- |
| New Visit | 27,497 | 62.5 | 921 | 48.5 | 5,128 | 76.4 | 3,373 | 61.7 | 566 | 71.4 | 3,758 | 67.0 | 1,468 | 67.8 |
| Return Visit | 16,476 | 37.5 | 979 | 51.5 | 1,585 | 23.6 | 2,098 | 38.3 | 227 | 28.6 | 1,850 | 33.0 | 696 | 32.2 |

*Number of visits

†Transgendered in total visits for all clinics include 36 male-to-female (MTF) and 72 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

†††Average number of new visits per session i.e. total number of new visits 2010 / total number of sessions 2010. See data section (Appendix B) for definition of a new visit. A session is a half-day when the STD clinic is open.

**Table 1.2 Summary Morbidity Information
January - December 2010
All Health Centers (HC)**

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|---------------|------------------|-----------------|---------------|------------------|------------------|----------------|--------------------------------------|-----|------------------|-----------------|---------------|-------------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** n % | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Positive % | Visits Tested | Positive n % | Positive % | Visits Tested | Positive* n % | Positive* % | | | Visits Tested | Positive n % | Positive % | New HIV*** n % | |
| All Sites | 43,973 | 100.0 | 28,275 | 100.0 | 27,173 | 2,954 | 10.9 | 27,266 | 1,292 | 4.7 | 25,137 | 1,433 | 5.7 | 120 | 0.5 | 23,285 | 177 | 0.8 | 131 | 0.6 |
| Antelope Valley HC | 1,613 | 3.7 | 1,195 | 4.2 | 1,149 | 173 | 15.1 | 1,156 | 50 | 4.3 | 1,071 | 33 | 3.1 | 0 | 0.0 | 1,045 | 4 | 0.4 | 3 | 0.3 |
| Simms Mann HC | 806 | 1.8 | 714 | 2.5 | 700 | 34 | 4.9 | 701 | 12 | 1.7 | 648 | 7 | 1.1 | 1 | 0.2 | 613 | 1 | 0.2 | 1 | 0.2 |
| Central HC | 5,882 | 13.4 | 3,135 | 11.1 | 3,003 | 231 | 7.7 | 3,008 | 105 | 3.5 | 2,919 | 288 | 9.9 | 19 | 0.7 | 2,728 | 28 | 1.0 | 20 | 0.7 |
| Hollywood/Wilshire HC | 6,205 | 14.1 | 3,747 | 13.3 | 3,529 | 286 | 8.1 | 3,536 | 144 | 4.1 | 3,112 | 215 | 6.9 | 22 | 0.7 | 2,758 | 26 | 0.9 | 20 | 0.7 |
| Monrovia HC | 1,222 | 2.8 | 913 | 3.2 | 861 | 98 | 11.4 | 862 | 28 | 3.2 | 838 | 69 | 8.2 | 6 | 0.7 | 757 | 5 | 0.7 | 4 | 0.5 |
| North Hollywood HC | 5,596 | 12.7 | 3,302 | 11.7 | 3,099 | 235 | 7.6 | 3,105 | 109 | 3.5 | 3,014 | 157 | 5.2 | 17 | 0.6 | 2,720 | 25 | 0.9 | 21 | 0.8 |
| Pomona HC | 1,900 | 4.3 | 1,061 | 3.8 | 987 | 167 | 16.9 | 990 | 40 | 4.0 | 966 | 48 | 5.0 | 6 | 0.6 | 884 | 4 | 0.5 | 4 | 0.5 |
| South HC | 6,713 | 15.3 | 5,139 | 18.2 | 5,091 | 721 | 14.2 | 5,095 | 323 | 6.3 | 4,513 | 153 | 3.4 | 15 | 0.3 | 4,306 | 17 | 0.4 | 12 | 0.3 |
| Temple HC | 5,471 | 12.4 | 3,569 | 12.6 | 3,507 | 443 | 12.6 | 3,523 | 229 | 6.5 | 3,158 | 210 | 6.6 | 14 | 0.4 | 2,890 | 28 | 1.0 | 16 | 0.6 |
| Torrance HC | 793 | 1.8 | 613 | 2.2 | 580 | 66 | 11.4 | 583 | 31 | 5.3 | 579 | 37 | 6.4 | 2 | 0.3 | 526 | 8 | 1.5 | 6 | 1.1 |
| C. Tucker HC | 5,608 | 12.8 | 3,554 | 12.6 | 3,399 | 346 | 10.2 | 3,431 | 180 | 5.2 | 3,091 | 121 | 3.9 | 10 | 0.3 | 2,879 | 19 | 0.7 | 17 | 0.6 |
| Whittier HC | 2,164 | 4.9 | 1,333 | 4.7 | 1,268 | 154 | 12.1 | 1,276 | 41 | 3.2 | 1,227 | 95 | 7.7 | 8 | 0.7 | 1,179 | 12 | 1.0 | 7 | 0.6 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on H2541 Intake Form

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested; similarly for RPR/TPPA, EIA/WB, etc.)

Figure 1.2 Sexually Transmitted Disease Prevalence in Public Health Centers, Los Angeles County, 2010




Table 1.3 Patient Visits and Tests, January - December 2010, All Public Health Centers

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|---------------------------------|--|-------|------------------------------------|-------|------------------|---------------|------|------------------|---------------|------|------------------|----------------|------|-------------------------------|-------------|------------------|---------------|-----|-----------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive* n | % | n | % | Visits Tested | Positive n | % | New HIV*** n | % |
| Total | 43,973 | 100.0 | 28,275 | 100.0 | 27,173 | 2,954 | 10.9 | 27,266 | 1,292 | 4.7 | 25,137 | 1,433 | 5.7 | 120 | 0.5 | 23,285 | 177 | 0.8 | 131 | 0.6 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 28,441 | 64.7 | 17,959 | 63.5 | 17,326 | 1,845 | 10.6 | 17,374 | 972 | 5.6 | 16,155 | 981 | 6.1 | 104 | 0.6 | 14,921 | 160 | 1.1 | 121 | 0.8 |
| Female | 15,424 | 35.1 | 10,271 | 36.3 | 9,804 | 1,108 | 11.3 | 9,848 | 315 | 3.2 | 8,939 | 443 | 5.0 | 16 | 0.2 | 8,328 | 17 | 0.2 | 10 | 0.1 |
| Transgender† | 108 | 0.2 | 45 | 0.2 | 43 | 1 | 2.3 | 44 | 5 | 11.4 | 43 | 9 | 20.9 | 0 | 0.0 | 36 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partners†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 4,963 | 11.3 | 3,000 | 10.6 | 2,821 | 331 | 11.7 | 2,836 | 392 | 13.8 | 2,689 | 444 | 16.5 | 70 | 2.6 | 2225 | 125 | 5.6 | 94 | 4.2 |
| MSW | 22,210 | 50.5 | 14,370 | 50.8 | 14,005 | 1,482 | 10.6 | 14,036 | 561 | 4.0 | 12,920 | 458 | 3.5 | 30 | 0.2 | 12244 | 29 | 0.2 | 22 | 0.2 |
| WSM | 13,416 | 30.5 | 9,055 | 32.0 | 8,700 | 1,010 | 11.6 | 8,738 | 267 | 3.1 | 7,879 | 364 | 4.6 | 16 | 0.2 | 7397 | 13 | 0.2 | 9 | 0.1 |
| WSW/M | 1,104 | 2.5 | 767 | 2.7 | 745 | 64 | 8.6 | 751 | 43 | 5.7 | 645 | 18 | 2.8 | 0 | 0.0 | 605 | 2 | 0.3 | 0 | 0.0 |
| Unknown | 2,280 | 5.2 | 1,083 | 3.8 | 902 | 67 | 7.4 | 905 | 29 | 3.2 | 1,004 | 149 | 14.8 | 4 | 0.4 | 814 | 8 | 1.0 | 6 | 0.7 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 82 | 0.2 | 47 | 0.2 | 46 | 15 | 32.6 | 46 | 3 | 6.5 | 43 | 0 | 0.0 | 0 | 0.0 | 42 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 3,636 | 8.3 | 2,557 | 9.0 | 2,492 | 623 | 25.0 | 2,499 | 216 | 8.6 | 2,258 | 46 | 2.0 | 5 | 0.2 | 2,165 | 10 | 0.5 | 8 | 0.4 |
| 20-24 | 9,737 | 22.1 | 6,611 | 23.4 | 6,460 | 1,086 | 16.8 | 6,483 | 395 | 6.1 | 5,832 | 166 | 2.8 | 24 | 0.4 | 5,573 | 39 | 0.7 | 32 | 0.6 |
| 25-29 | 8,691 | 19.8 | 5,678 | 20.1 | 5,530 | 569 | 10.3 | 5,550 | 261 | 4.7 | 5,068 | 181 | 3.6 | 23 | 0.5 | 4,813 | 34 | 0.7 | 25 | 0.5 |
| 30-34 | 6,272 | 14.3 | 3,968 | 14.0 | 3,835 | 285 | 7.4 | 3,857 | 161 | 4.2 | 3,523 | 156 | 4.4 | 14 | 0.4 | 3,271 | 34 | 1.0 | 25 | 0.8 |
| 35-44 | 8,123 | 18.5 | 5,058 | 17.9 | 4,841 | 267 | 5.5 | 4,849 | 153 | 3.2 | 4,488 | 343 | 7.6 | 24 | 0.5 | 4,044 | 38 | 0.9 | 28 | 0.7 |
| 45-54 | 5,148 | 11.7 | 3,096 | 10.9 | 2,907 | 83 | 2.9 | 2,918 | 80 | 2.7 | 2,768 | 302 | 10.9 | 19 | 0.7 | 2,451 | 13 | 0.5 | 9 | 0.4 |
| 55-64 | 1,798 | 4.1 | 1,019 | 3.6 | 880 | 23 | 2.6 | 882 | 19 | 2.2 | 927 | 177 | 19.1 | 6 | 0.6 | 761 | 7 | 0.9 | 3 | 0.4 |
| 65+ | 486 | 1.1 | 241 | 0.9 | 182 | 3 | 1.6 | 182 | 4 | 2.2 | 230 | 62 | 27.0 | 5 | 2.2 | 165 | 2 | 1.2 | 1 | 0.6 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 6,935 | 15.8 | 4,472 | 15.8 | 4,324 | 284 | 6.6 | 4,334 | 126 | 2.9 | 3,979 | 99 | 2.5 | 11 | 0.3 | 3748 | 18 | 0.5 | 15 | 0.4 |
| African American | 17,464 | 39.7 | 12,220 | 43.2 | 11,944 | 1,600 | 13.4 | 11,994 | 829 | 6.9 | 10,563 | 417 | 3.9 | 41 | 0.4 | 9832 | 74 | 0.8 | 52 | 0.5 |
| Hispanic | 15,942 | 36.3 | 9,283 | 32.8 | 8,706 | 874 | 10.0 | 8,734 | 271 | 3.1 | 8,504 | 801 | 9.4 | 58 | 0.7 | 7764 | 76 | 1.0 | 55 | 0.7 |
| Asian | 1,890 | 4.3 | 1,191 | 4.2 | 1,117 | 99 | 8.9 | 1,120 | 24 | 2.1 | 1,095 | 72 | 6.6 | 6 | 0.5 | 989 | 3 | 0.3 | 3 | 0.3 |
| Native American | 109 | 0.2 | 75 | 0.3 | 74 | 8 | 10.8 | 74 | 6 | 8.1 | 71 | 9 | 12.7 | 0 | 0.0 | 67 | 0 | 0.0 | 0 | 0.0 |
| Other | 1,410 | 3.2 | 904 | 3.2 | 881 | 74 | 8.4 | 883 | 29 | 3.3 | 804 | 30 | 3.7 | 4 | 0.5 | 772 | 5 | 0.6 | 5 | 0.6 |
| Unknown | 223 | 0.5 | 130 | 0.5 | 127 | 15 | 11.8 | 127 | 7 | 5.5 | 121 | 5 | 4.1 | 0 | 0.0 | 113 | 1 | 0.9 | 1 | 0.9 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 27,497 | 62.5 | 26,291 | 93.0 | 25,885 | 2,824 | 10.9 | 25,942 | 1,224 | 4.7 | 23,978 | 1,227 | 5.1 | 103 | 0.4 | 22,661 | 158 | 0.7 | 124 | 0.5 |
| Return Visit | 16,476 | 37.5 | 1,984 | 7.0 | 1,288 | 130 | 10.1 | 1,324 | 68 | 5.1 | 1,159 | 206 | 17.8 | 17 | 1.5 | 624 | 19 | 3.0 | 7 | 1.1 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator


***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic (H2541) Intake Form

†Transgendered in total visits include 36 male-to-female (MTF) and 72 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested; similarly for RPR/TPPA, EIA/WB, etc.)

Figure 1.3
Primary Reason(s) for New Visit* among Males
January - December 2010
All Health Centers




*Visit classification and clinical diagnoses as indicated by the Clinic Intake Form

**PHI=Public health investigator, PHN=Public health nurse

Note: Categories other than "check-up only" are non-exclusive

Figure 1.4
Primary Reason(s) for New Visit* among Females
January - December 2010
All Health Centers



*Visit classification and clinical diagnoses as indicated by the Clinic Intake Form

**PHI=Public health investigator, PHN=Public health nurse

Note: Categories other than "check-up only" are non-exclusive

Table 1.4 Patient Visits and Tests by Gender and Age, January - December 2010, All Public Health Centers

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|-----------------|--|-------|----------------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|-------|---------------|----------------------------|-------------------|-------------|-----|-----|-----|-----|
| | Total STD Visits n % | | No. Visits | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | with STD Test n % | Visits Tested | Positive n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 43,973 | 100.0 | 28,275 | 100.0 | 27,173 | 2,954 | 10.9 | 27,266 | 1,292 | 4.7 | 25,137 | 1,433 | 5.7 | 120 | 0.5 | 23,285 | 177 | 0.8 | 131 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 28 | 0.1 | 13 | 0.1 | 13 | 3 | 23.1 | 13 | 0 | 0.0 | 13 | 0 | 0.0 | 0 | 0.0 | 13 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 1,625 | 5.7 | 1,121 | 6.2 | 1,098 | 256 | 23.3 | 1,102 | 120 | 10.9 | 1,024 | 32 | 3.1 | 4 | 0.4 | 983 | 7 | 0.7 | 6 | 0.6 |
| 20-24 | 5,839 | 20.5 | 3,885 | 21.6 | 3,805 | 654 | 17.2 | 3,816 | 278 | 7.3 | 3,502 | 126 | 3.6 | 21 | 0.6 | 3,345 | 38 | 1.1 | 31 | 0.9 |
| 25-29 | 6,015 | 21.1 | 3,848 | 21.4 | 3,766 | 410 | 10.9 | 3,776 | 220 | 5.8 | 3,467 | 131 | 3.8 | 23 | 0.7 | 3,305 | 34 | 1.0 | 25 | 0.8 |
| 30-34 | 4,348 | 15.3 | 2,734 | 15.2 | 2,655 | 222 | 8.4 | 2,669 | 121 | 4.5 | 2,433 | 114 | 4.7 | 13 | 0.5 | 2,269 | 30 | 1.3 | 22 | 1.0 |
| 35-44 | 5,568 | 19.6 | 3,420 | 19.0 | 3,296 | 213 | 6.5 | 3,302 | 141 | 4.3 | 3,052 | 208 | 6.8 | 20 | 0.7 | 2,737 | 35 | 1.3 | 27 | 1.0 |
| 45-54 | 3,460 | 12.2 | 2,055 | 11.4 | 1,934 | 69 | 3.6 | 1,936 | 70 | 3.6 | 1,857 | 216 | 11.6 | 16 | 0.9 | 1,612 | 10 | 0.6 | 8 | 0.5 |
| 55-64 | 1,253 | 4.4 | 729 | 4.1 | 637 | 15 | 2.4 | 638 | 18 | 2.8 | 659 | 118 | 17.9 | 2 | 0.3 | 546 | 4 | 0.7 | 1 | 0.2 |
| 65+ | 319 | 1.1 | 159 | 0.9 | 127 | 3 | 2.4 | 127 | 4 | 3.1 | 152 | 38 | 25.0 | 5 | 3.3 | 115 | 2 | 1.7 | 1 | 0.9 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 28,455 | 100.0 | 17,964 | 100.0 | 17,331 | 1,845 | 10.6 | 17,379 | 972 | 5.6 | 16,159 | 983 | 6.1 | 104 | 0.6 | 14,925 | 160 | 1.1 | 121 | 0.8 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 54 | 0.3 | 34 | 0.3 | 33 | 12 | 36.4 | 33 | 3 | 9.1 | 30 | 0 | 0.0 | 0 | 0.0 | 29 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 2,007 | 13.0 | 1,434 | 14.0 | 1,392 | 367 | 26.4 | 1,395 | 96 | 6.9 | 1,232 | 14 | 1.1 | 1 | 0.1 | 1,180 | 3 | 0.3 | 2 | 0.2 |
| 20-24 | 3,883 | 25.1 | 2,720 | 26.5 | 2,649 | 432 | 16.3 | 2,661 | 117 | 4.4 | 2,324 | 37 | 1.6 | 3 | 0.1 | 2,222 | 1 | 0.0 | 1 | 0.0 |
| 25-29 | 2,658 | 17.2 | 1,823 | 17.7 | 1,757 | 159 | 9.0 | 1,767 | 41 | 2.3 | 1,594 | 49 | 3.1 | 0 | 0.0 | 1,502 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 1,905 | 12.3 | 1,222 | 11.9 | 1,170 | 63 | 5.4 | 1,178 | 38 | 3.2 | 1,081 | 42 | 3.9 | 1 | 0.1 | 994 | 4 | 0.4 | 3 | 0.3 |
| 35-44 | 2,538 | 16.4 | 1,628 | 15.8 | 1,536 | 53 | 3.5 | 1,537 | 10 | 0.7 | 1,427 | 133 | 9.3 | 4 | 0.3 | 1,301 | 3 | 0.2 | 1 | 0.1 |
| 45-54 | 1,684 | 10.9 | 1,039 | 10.1 | 971 | 14 | 1.4 | 980 | 10 | 1.0 | 909 | 85 | 9.4 | 3 | 0.3 | 837 | 3 | 0.4 | 1 | 0.1 |
| 55-64 | 545 | 3.5 | 290 | 2.8 | 243 | 8 | 3.3 | 244 | 1 | 0.4 | 268 | 59 | 22.0 | 4 | 1.5 | 215 | 3 | 1.4 | 2 | 0.9 |
| 65+ | 167 | 1.1 | 82 | 0.8 | 55 | 0 | 0.0 | 55 | 0 | 0.0 | 78 | 24 | 30.8 | 0 | 0.0 | 50 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 15,439 | 100.0 | 10,272 | 100.0 | 9,806 | 1,108 | 11.3 | 9,850 | 316 | 3.2 | 8,943 | 443 | 5.0 | 16 | 0.2 | 8,330 | 17 | 0.2 | 10 | 0.1 |
| Others † | 79 | ~ | 39 | ~ | 36 | 1 | 2.8 | 37 | 4 | 10.8 | 35 | 7 | 20.0 | 0 | 0.0 | 30 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 1.5 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, All Public Health Centers

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | HIV Testing | | | | | | |
|-----------------------|--|-------|----------------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|--------------------------------------|-------------------|--------|-----|-----|-----|-----|
| | Total STD Visits n % | | No. Visits | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** n % | HIV | | | | | |
| | | | with STD Test n % | Visits Tested | Positive n % | | New HIV*** n % | | | | | |
| Total | 43,973 | 100.0 | 28,275 | 100.0 | 27,173 | 2,954 | 10.9 | 27,266 | 1,292 | 4.7 | 25,137 | 1,433 | 5.7 | 120 | 0.5 | 23,285 | 177 | 0.8 | 131 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 5,138 | 18.0 | 3,307 | 18.4 | 3,209 | 210 | 6.5 | 3,215 | 108 | 3.4 | 2,957 | 83 | 2.8 | 9 | 0.3 | 2,761 | 18 | 0.7 | 15 | 0.5 |
| African American | 10,136 | 35.6 | 6,946 | 38.7 | 6,804 | 969 | 14.2 | 6,821 | 589 | 8.6 | 6,100 | 252 | 4.1 | 34 | 0.6 | 5,636 | 66 | 1.2 | 47 | 0.8 |
| Hispanic | 10,881 | 38.2 | 6,275 | 34.9 | 5,925 | 548 | 9.2 | 5,949 | 223 | 3.7 | 5,783 | 576 | 10.0 | 53 | 0.9 | 5,296 | 68 | 1.3 | 51 | 1.0 |
| Asian | 1,168 | 4.1 | 730 | 4.1 | 698 | 67 | 9.6 | 699 | 23 | 3.3 | 676 | 40 | 5.9 | 5 | 0.7 | 616 | 2 | 0.3 | 2 | 0.3 |
| Native American | 65 | 0.2 | 47 | 0.3 | 47 | 6 | 12.8 | 47 | 5 | 10.6 | 43 | 7 | 16.3 | 0 | 0.0 | 41 | 0 | 0.0 | 0 | 0.0 |
| Other | 900 | 3.2 | 567 | 3.2 | 559 | 36 | 6.4 | 559 | 19 | 3.4 | 514 | 20 | 3.9 | 3 | 0.6 | 496 | 5 | 1.0 | 5 | 1.0 |
| Unknown | 167 | 0.6 | 92 | 0.5 | 89 | 9 | 10.1 | 89 | 5 | 5.6 | 86 | 5 | 5.8 | 0 | 0.0 | 79 | 1 | 1.3 | 1 | 1.3 |
| Subtotal | 28,455 | 100.0 | 17,964 | 100.0 | 17,331 | 1,845 | 10.6 | 17,379 | 972 | 5.6 | 16,160 | 983 | 6.1 | 104 | 0.6 | 14,925 | 160 | 1.1 | 121 | 0.8 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 1,794 | 11.6 | 1,163 | 11.3 | 1,113 | 74 | 6.6 | 1,117 | 18 | 1.6 | 1,020 | 16 | 1.6 | 2 | 0.2 | 985 | 0 | 0.0 | 0 | 0.0 |
| African American | 7,322 | 47.4 | 5,269 | 51.3 | 5,135 | 631 | 12.3 | 5,168 | 239 | 4.6 | 4,458 | 165 | 3.7 | 7 | 0.2 | 4,194 | 8 | 0.2 | 5 | 0.1 |
| Hispanic | 5,005 | 32.4 | 2,984 | 29.0 | 2,759 | 325 | 11.8 | 2,762 | 45 | 1.6 | 2,698 | 218 | 8.1 | 5 | 0.2 | 2,447 | 8 | 0.3 | 4 | 0.2 |
| Asian | 718 | 4.6 | 459 | 4.5 | 417 | 32 | 7.7 | 419 | 1 | 0.2 | 417 | 32 | 7.7 | 1 | 0.2 | 371 | 1 | 0.3 | 1 | 0.3 |
| Native American | 44 | 0.3 | 28 | 0.3 | 27 | 2 | 7.4 | 27 | 1 | 3.7 | 28 | 2 | 7.1 | 0 | 0.0 | 26 | 0 | 0.0 | 0 | 0.0 |
| Other | 500 | 3.2 | 331 | 3.2 | 317 | 38 | 12.0 | 319 | 10 | 3.1 | 287 | 10 | 3.5 | 1 | 0.3 | 273 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 82 | 0.5 | 38 | 0.4 | 38 | 6 | 15.8 | 38 | 2 | 5.3 | 35 | 0 | 0.0 | 0 | 0.0 | 34 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 15,439 | 100.0 | 10,272 | 100.0 | 9,806 | 1,108 | 11.3 | 9,850 | 316 | 3.2 | 8,943 | 443 | 5.0 | 16 | 0.2 | 8,330 | 17 | 0.2 | 10 | 0.1 |
| Others † | 79 | ~ | 39 | ~ | 36 | 1 | 2.8 | 37 | 4 | 10.8 | 35 | 7 | 20.0 | 0 | 0.0 | 30 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinical Intake Form

†Includes transgender and gender unspecified

Figure 1.5. Chlamydia Cases Among Males in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




Figure 1.6. Chlamydia Cases Among Females in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




Figure 1.7. Gonorrhea Cases Among Males in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




Figure 1.8. Gonorrhea Cases Among Females in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




Figure 1.9. Early Syphilis Cases Among Males in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




Figure 1.10. HIV Cases Among Males in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010






Figure 1.11. New HIV Cases* Among Males in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




* Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

Figure 1.12. New HIV Cases* Among Females in Los Angeles County STD Clinics, by Age Group and Race/Ethnicity, 2010




* Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

**Figure 1.13a. Sexually Transmitted Disease Prevalence, 2006-2010
All Health Centers (HC)**




Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

**Figure 1.13b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010
All Health Centers (HC)**




**Figure 1.14. Chlamydia Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.15. Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.16. Early Syphilis Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic data collection on gender of sex partners started in latter half of that year.

**Figure 1.17. HIV Prevalence by Gender/Sexual Orientation, 2006-2010
All Health Centers (HC)**



Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.18 New HIV Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**



Note: MSM/W and MSW specific prevalence not available for 2006, as H2541 data collection on gender of sex partners started in latter half of that year.

**Table 1.6 Site of Chlamydia Infection by Gender
January - December 2010**

| | Chlamydia by Anatomic Site* | | | | | | | | | | | |
|--------------------------------|-----------------------------|------------|------|---------------|------------|------|---------------|------------|------|------------------------|-------------------------|------|
| | Overall Infections | | | | | | | | | Concomitant Infections | | |
| | Total | | | Urogenital | | | Rectal | | | Urogenital/Rectal | | |
| | Visits Tested** | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested† | Positive ¹ n | % |
| All Health Centers (HC) | | | | | | | | | | | | |
| Total | 27,173 | 2,954 | 10.9 | 27,133 | 2,735 | 10.1 | 2,090 | 283 | 13.5 | 2,051 | 64 | 3.1 |
| Male | 17,326 | 1,845 | 10.6 | 17,295 | 1,642 | 9.5 | 1,617 | 219 | 13.5 | 1,586 | 16 | 1.0 |
| Female | 9,804 | 1,108 | 11.3 | 9,796 | 1,093 | 11.2 | 446 | 63 | 14.1 | 439 | 48 | 10.9 |
| Transgender | 43 | 1 | 2.3 | 42 | 0 | 0.0 | 27 | 1 | 3.7 | 26 | 0 | 0.0 |
| Antelope Valley HC | | | | | | | | | | | | |
| Total | 1,149 | 173 | 15.1 | 1,145 | 163 | 14.2 | 66 | 12 | 18.2 | 62 | 2 | 3.2 |
| Male | 522 | 87 | 16.7 | 518 | 77 | 14.9 | 59 | 11 | 18.6 | 55 | 1 | 1.8 |
| Female | 627 | 86 | 13.7 | 627 | 86 | 13.7 | 7 | 1 | 14.3 | 7 | 1 | 14.3 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| Simms Mann HC | | | | | | | | | | | | |
| Total | 700 | 34 | 4.9 | 700 | 34 | 4.9 | 28 | 2 | 7.1 | 28 | 2 | 7.1 |
| Male | 548 | 25 | 4.6 | 548 | 25 | 4.6 | 25 | 1 | 4.0 | 25 | 1 | 4.0 |
| Female | 152 | 9 | 5.9 | 152 | 9 | 5.9 | 3 | 1 | 33.3 | 3 | 1 | 33.3 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| Central HC | | | | | | | | | | | | |
| Total | 3,003 | 231 | 7.7 | 2,999 | 197 | 6.6 | 461 | 48 | 10.4 | 457 | 14 | 3.1 |
| Male | 2,133 | 160 | 7.5 | 2,130 | 131 | 6.2 | 356 | 32 | 9.0 | 353 | 3 | 0.8 |
| Female | 861 | 70 | 8.1 | 860 | 66 | 7.7 | 98 | 15 | 15.3 | 97 | 11 | 11.3 |
| Transgender | 9 | 1 | 11.1 | 9 | 0 | 0.0 | 7 | 1 | 14.3 | 7 | 0 | 0.0 |
| Hollywood-Wilshire HC | | | | | | | | | | | | |
| Total | 3,529 | 286 | 8.1 | 3,517 | 256 | 7.3 | 319 | 34 | 10.7 | 307 | 4 | 1.3 |
| Male | 2,642 | 222 | 8.4 | 2,631 | 192 | 7.3 | 296 | 32 | 10.8 | 285 | 2 | 0.7 |
| Female | 870 | 64 | 7.4 | 870 | 64 | 7.4 | 15 | 2 | 13.3 | 15 | 2 | 13.3 |
| Transgender | 17 | 0 | 0.0 | 16 | 0 | 0.0 | 8 | 0 | 0.0 | 7 | 0 | 0.0 |
| Monrovia HC | | | | | | | | | | | | |
| Total | 861 | 98 | 11.4 | 859 | 78 | 9.1 | 86 | 20 | 23.3 | 85 | 0 | 0.0 |
| Male | 585 | 74 | 12.6 | 584 | 54 | 9.2 | 82 | 20 | 24.4 | 81 | 0 | 0.0 |
| Female | 274 | 24 | 8.8 | 273 | 24 | 8.8 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| Transgender | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| North Hollywood HC | | | | | | | | | | | | |
| Total | 3,099 | 235 | 7.6 | 3,092 | 180 | 5.8 | 348 | 59 | 17.0 | 341 | 4 | 1.2 |
| Male | 2,231 | 175 | 7.8 | 2,224 | 121 | 5.4 | 327 | 57 | 17.4 | 320 | 3 | 0.9 |
| Female | 860 | 60 | 7.0 | 860 | 59 | 6.9 | 16 | 2 | 12.5 | 16 | 1 | 6.3 |
| Transgender | 8 | 0 | 0.0 | 8 | 0 | 0.0 | 5 | 0 | 0.0 | 5 | 0 | 0.0 |

¹Positive at both urogenital and rectal sites on the same visit

*Anatomic site categories are not mutually exclusive (eg. Infections listed under the concomitant category "urogenital/rectal" are also included under the overall categories "urogenital" and "rectal")

***"Visits Tested" for chlamydia refers to number of patient visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested under "Total" chlamydia)

†Tested at both urogenital and rectal sites on the same visit

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

**Table 1.6 (continued) Site of Chlamydia Infection by Gender
January - December 2010**

| | Chlamydia by Anatomic Site* | | | | | | | | | | | |
|-------------------------|-----------------------------|------------|------|---------------|------------|------|---------------|------------|------|------------------------|-------------------------|------|
| | Overall Infections | | | | | | | | | Concomitant Infections | | |
| | Total | | | Urogenital | | | Rectal | | | Urogenital/Rectal | | |
| | Visits Tested** | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested† | Positive ¹ n | % |
| Pomona HC | | | | | | | | | | | | |
| Total | 987 | 167 | 16.9 | 985 | 152 | 15.4 | 112 | 22 | 19.6 | 110 | 7 | 6.4 |
| Male | 671 | 97 | 14.5 | 669 | 83 | 12.4 | 89 | 14 | 15.7 | 87 | 0 | 0.0 |
| Female | 316 | 70 | 22.2 | 316 | 69 | 21.8 | 23 | 8 | 34.8 | 23 | 7 | 30.4 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| South HC | | | | | | | | | | | | |
| Total | 5,091 | 721 | 14.2 | 5,091 | 710 | 13.9 | 168 | 24 | 14.3 | 168 | 13 | 7.7 |
| Male | 2,686 | 371 | 13.8 | 2,686 | 362 | 13.5 | 72 | 10 | 13.9 | 72 | 1 | 1.4 |
| Female | 2,401 | 350 | 14.6 | 2,401 | 348 | 14.5 | 94 | 14 | 14.9 | 94 | 12 | 12.8 |
| Transgender | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| Ruth Temple HC | | | | | | | | | | | | |
| Total | 3,507 | 443 | 12.6 | 3,506 | 432 | 12.3 | 90 | 15 | 16.7 | 89 | 4 | 4.5 |
| Male | 2,005 | 262 | 13.1 | 2,005 | 253 | 12.6 | 62 | 11 | 17.7 | 62 | 2 | 3.2 |
| Female | 1,500 | 181 | 12.1 | 1,499 | 179 | 11.9 | 26 | 4 | 15.4 | 25 | 2 | 8.0 |
| Transgender | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| Torrance HC | | | | | | | | | | | | |
| Total | 580 | 66 | 11.4 | 579 | 59 | 10.2 | 49 | 9 | 18.4 | 48 | 2 | 4.2 |
| Male | 410 | 48 | 11.7 | 409 | 41 | 10.0 | 37 | 9 | 24.3 | 36 | 2 | 5.6 |
| Female | 170 | 18 | 10.6 | 170 | 18 | 10.6 | 12 | 0 | 0.0 | 12 | 0 | 0.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| Curtis Tucker HC | | | | | | | | | | | | |
| Total | 3,399 | 346 | 10.2 | 3,394 | 333 | 9.8 | 202 | 22 | 10.9 | 197 | 9 | 4.6 |
| Male | 2,027 | 228 | 11.2 | 2,026 | 218 | 10.8 | 102 | 11 | 10.8 | 101 | 1 | 1.0 |
| Female | 1,371 | 118 | 8.6 | 1,367 | 115 | 8.4 | 99 | 11 | 11.1 | 95 | 8 | 8.4 |
| Transgender | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 |
| Whittier HC | | | | | | | | | | | | |
| Total | 1,268 | 154 | 12.1 | 1,266 | 141 | 11.1 | 161 | 16 | 9.9 | 159 | 3 | 1.9 |
| Male | 866 | 96 | 11.1 | 865 | 85 | 9.8 | 110 | 11 | 10.0 | 109 | 0 | 0.0 |
| Female | 402 | 58 | 14.4 | 401 | 56 | 14.0 | 51 | 5 | 9.8 | 50 | 3 | 6.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |

¹Positive at both urogenital and rectal sites on the same visit

*Anatomic site categories are not mutually exclusive (eg. Infections listed under the concomitant category "urogenital/rectal" are also included under the overall categories "urogenital" and "rectal")

**"Visits Tested" for chlamydia refers to number of patient visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested under "Total" chlamydia)

†Tested at both urogenital and rectal sites on the same visit

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

**Table 1.7 Site of Gonorrhea Infection by Gender
January - December 2010**

| | Gonorrhea by Anatomic Site* | | | | | | | | | | | | | | | | | | | | | | | |
|--------------------------------|-----------------------------|------------|------|---------------|------------|------|---------------|------------|------|---------------|------------|------|------------------------|-------------------------|------|-------------------|-------------------------|------|----------------|-------------------------|------|-------------------|-------------------------|------|
| | Overall Infections | | | | | | | | | | | | Concomitant Infections | | | | | | | | | | | |
| | Total | | | Urogenital | | | Pharyngeal | | | Rectal | | | Urogen./Pharyn. | | | Urogenital/Rectal | | | Pharyn./Rectal | | | Urog./Phar./Rect. | | |
| | Visits Tested** | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested† | Positive ¹ n | % | Visits Tested† | Positive ² n | % | Visits Tested† | Positive ³ n | % | Visits Tested† | Positive ⁴ n | % |
| All Health Centers (HC) | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 27,266 | 1,292 | 4.7 | 27,133 | 1,018 | 3.8 | 6,800 | 335 | 4.9 | 2,090 | 193 | 9.2 | 6,688 | 131 | 2.0 | 2,051 | 73 | 3.6 | 1,865 | 84 | 4.5 | 1,847 | 34 | 1.8 |
| Male | 17,374 | 972 | 5.6 | 17,295 | 734 | 4.2 | 4,505 | 257 | 5.7 | 1,617 | 178 | 11.0 | 4,442 | 87 | 2.0 | 1,586 | 61 | 3.8 | 1,533 | 76 | 5.0 | 1,518 | 27 | 1.8 |
| Female | 9,848 | 315 | 3.2 | 9,796 | 282 | 2.9 | 2,262 | 75 | 3.3 | 446 | 13 | 2.9 | 2,215 | 43 | 1.9 | 439 | 11 | 2.5 | 307 | 7 | 2.3 | 305 | 6 | 2.0 |
| Transgender | 44 | 5 | 11.4 | 42 | 2 | 4.8 | 33 | 3 | 9.1 | 27 | 2 | 7.4 | 31 | 1 | 3.2 | 26 | 1 | 3.8 | 25 | 1 | 4.0 | 24 | 1 | 4.2 |
| Antelope Valley HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 1,156 | 50 | 4.3 | 1,145 | 43 | 3.8 | 105 | 12 | 11.4 | 66 | 5 | 7.6 | 96 | 6 | 6.3 | 62 | 3 | 4.8 | 61 | 3 | 4.9 | 59 | 2 | 3.4 |
| Male | 525 | 33 | 6.3 | 518 | 27 | 5.2 | 79 | 10 | 12.7 | 59 | 5 | 8.5 | 74 | 5 | 6.8 | 55 | 3 | 5.5 | 56 | 3 | 5.4 | 54 | 2 | 3.7 |
| Female | 631 | 17 | 2.7 | 627 | 16 | 2.6 | 26 | 2 | 7.7 | 7 | 0 | 0.0 | 22 | 1 | 4.5 | 7 | 0 | 0.0 | 5 | 0 | 0.0 | 5 | 0 | 0.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| Simms Mann HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 701 | 12 | 1.7 | 700 | 9 | 1.3 | 65 | 4 | 6.2 | 28 | 2 | 7.1 | 64 | 2 | 3.1 | 28 | 0 | 0.0 | 26 | 1 | 3.8 | 26 | 0 | 0.0 |
| Male | 548 | 9 | 1.6 | 548 | 6 | 1.1 | 57 | 4 | 7.0 | 25 | 2 | 8.0 | 57 | 2 | 3.5 | 25 | 0 | 0.0 | 25 | 1 | 4.0 | 25 | 0 | 0.0 |
| Female | 153 | 3 | 2.0 | 152 | 3 | 2.0 | 8 | 0 | 0.0 | 3 | 0 | 0.0 | 7 | 0 | 0.0 | 3 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| Central HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 3,008 | 105 | 3.5 | 2,999 | 69 | 2.3 | 511 | 37 | 7.2 | 461 | 30 | 6.5 | 504 | 10 | 2.0 | 457 | 13 | 2.8 | 365 | 14 | 3.8 | 363 | 6 | 1.7 |
| Male | 2,137 | 93 | 4.4 | 2,130 | 59 | 2.8 | 472 | 36 | 7.6 | 356 | 28 | 7.9 | 466 | 10 | 2.1 | 353 | 12 | 3.4 | 350 | 14 | 4.0 | 348 | 6 | 1.7 |
| Female | 862 | 12 | 1.4 | 860 | 10 | 1.2 | 30 | 1 | 3.3 | 98 | 2 | 2.0 | 29 | 0 | 0.0 | 97 | 1 | 1.0 | 8 | 0 | 0.0 | 8 | 0 | 0.0 |
| Transgender | 9 | 0 | 0.0 | 9 | 0 | 0.0 | 9 | 0 | 0.0 | 7 | 0 | 0.0 | 9 | 0 | 0.0 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 7 | 0 | 0.0 |
| Hollywood-Wilshire HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 3,536 | 144 | 4.1 | 3,517 | 106 | 3.0 | 597 | 46 | 7.7 | 319 | 35 | 11.0 | 583 | 16 | 2.7 | 307 | 19 | 6.2 | 287 | 17 | 5.9 | 280 | 9 | 3.2 |
| Male | 2,647 | 137 | 5.2 | 2,631 | 99 | 3.8 | 504 | 43 | 8.5 | 296 | 34 | 11.5 | 493 | 13 | 2.6 | 285 | 18 | 6.3 | 270 | 16 | 5.9 | 264 | 8 | 3.0 |
| Female | 872 | 5 | 0.6 | 870 | 5 | 0.6 | 83 | 2 | 2.4 | 15 | 0 | 0.0 | 81 | 2 | 2.5 | 15 | 0 | 0.0 | 10 | 0 | 0.0 | 10 | 0 | 0.0 |
| Transgender | 17 | 2 | 11.8 | 16 | 2 | 12.5 | 10 | 1 | 10.0 | 8 | 1 | 12.5 | 9 | 1 | 11.1 | 7 | 1 | 14.3 | 7 | 1 | 14.3 | 6 | 1 | 16.7 |
| Monrovia HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 862 | 28 | 3.2 | 859 | 15 | 1.7 | 113 | 5 | 4.4 | 86 | 10 | 11.6 | 110 | 0 | 0.0 | 85 | 0 | 0.0 | 82 | 2 | 2.4 | 81 | 0 | 0.0 |
| Male | 587 | 24 | 4.1 | 584 | 11 | 1.9 | 106 | 5 | 4.7 | 82 | 10 | 12.2 | 103 | 0 | 0.0 | 81 | 0 | 0.0 | 79 | 2 | 2.5 | 78 | 0 | 0.0 |
| Female | 273 | 4 | 1.5 | 273 | 4 | 1.5 | 6 | 0 | 0.0 | 2 | 0 | 0.0 | 6 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| Transgender | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 1 | 0 | 0.0 | 2 | 0 | 0.0 | 1 | 0 | 0.0 | 2 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 |
| North Hollywood HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 3,105 | 109 | 3.5 | 3,092 | 62 | 2.0 | 602 | 36 | 6.0 | 348 | 44 | 12.6 | 595 | 7 | 1.2 | 341 | 14 | 4.1 | 324 | 16 | 4.9 | 323 | 4 | 1.2 |
| Male | 2,234 | 100 | 4.5 | 2,224 | 56 | 2.5 | 524 | 34 | 6.5 | 327 | 43 | 13.1 | 520 | 7 | 1.3 | 320 | 14 | 4.4 | 311 | 16 | 5.1 | 310 | 4 | 1.3 |
| Female | 862 | 6 | 0.7 | 860 | 6 | 0.7 | 70 | 0 | 0.0 | 16 | 0 | 0.0 | 68 | 0 | 0.0 | 16 | 0 | 0.0 | 8 | 0 | 0.0 | 8 | 0 | 0.0 |
| Transgender | 9 | 3 | 33.3 | 8 | 0 | 0.0 | 8 | 2 | 25.0 | 5 | 1 | 20.0 | 7 | 0 | 0.0 | 5 | 0 | 0.0 | 5 | 0 | 0.0 | 5 | 0 | 0.0 |

¹Positive at both urogenital and pharyngeal sites on the same visit

²Positive at both urogenital and rectal sites on the same visit

³Positive at both pharyngeal and rectal sites on the same visit

⁴Positive at urogenital, pharyngeal, and rectal sites on the same visit

*Anatomic site categories are not mutually exclusive (eg. Infections listed under the concomitant category "urogenital/rectal" are also included under the overall categories "urogenital" and "rectal")

**"Visits Tested" for gonorrhea refers to number of patient visits and not individual tests (ie. a patient who is tested for urethral, pharyngeal, and rectal gonorrhea on the same visit is only counted as one visit tested under "Total" gonorrhea)

†Tested at the specified anatomic sites on the same visit

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

**Table 1.7 (continued) Site of Gonorrhea Infection by Gender
January - December 2010**

| | Gonorrhea by Anatomic Site* | | | | | | | | | | | | | | | | | | | | | | | |
|-------------------------|-----------------------------|------------|-----|---------------|------------|-----|---------------|------------|-----|---------------|------------|------|------------------------|-------------------------|-------------------|----------------|-------------------------|-----|-------------------|-------------------------|------|----------------|-------------------------|-----|
| | Overall Infections | | | | | | | | | | | | Concomitant Infections | | | | | | | | | | | |
| | Total | | | Urogenital | | | Pharyngeal | | | Rectal | | | Urogen./Pharyn. | | Urogenital/Rectal | | Pharyn./Rectal | | Urog./Phar./Rect. | | | | | |
| | Visits Tested** | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested† | Positive ¹ n | % | Visits Tested† | Positive ² n | % | Visits Tested† | Positive ³ n | % | Visits Tested† | Positive ⁴ n | % |
| Pomona HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 990 | 40 | 4.0 | 985 | 20 | 2.0 | 550 | 22 | 4.0 | 112 | 10 | 8.9 | 546 | 8 | 1.5 | 110 | 1 | 0.9 | 105 | 4 | 3.8 | 104 | 1 | 1.0 |
| Male | 674 | 32 | 4.7 | 669 | 13 | 1.9 | 375 | 19 | 5.1 | 89 | 10 | 11.2 | 371 | 6 | 1.6 | 87 | 1 | 1.1 | 83 | 4 | 4.8 | 82 | 1 | 1.2 |
| Female | 316 | 8 | 2.5 | 316 | 7 | 2.2 | 175 | 3 | 1.7 | 23 | 0 | 0.0 | 175 | 2 | 1.1 | 23 | 0 | 0.0 | 22 | 0 | 0.0 | 22 | 0 | 0.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| South HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 5,095 | 323 | 6.3 | 5,091 | 297 | 5.8 | 711 | 42 | 5.9 | 168 | 12 | 7.1 | 707 | 20 | 2.8 | 168 | 6 | 3.6 | 158 | 5 | 3.2 | 158 | 3 | 1.9 |
| Male | 2,688 | 195 | 7.3 | 2,686 | 181 | 6.7 | 215 | 14 | 6.5 | 72 | 7 | 9.7 | 213 | 4 | 1.9 | 72 | 1 | 1.4 | 69 | 2 | 2.9 | 69 | 0 | 0.0 |
| Female | 2,403 | 128 | 5.3 | 2,401 | 116 | 4.8 | 494 | 28 | 5.7 | 94 | 5 | 5.3 | 492 | 16 | 3.3 | 94 | 5 | 5.3 | 87 | 3 | 3.4 | 87 | 3 | 3.4 |
| Transgender | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| Ruth Temple HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 3,523 | 229 | 6.5 | 3,506 | 203 | 5.8 | 763 | 41 | 5.4 | 90 | 16 | 17.8 | 747 | 20 | 2.7 | 89 | 6 | 6.7 | 82 | 8 | 9.8 | 82 | 3 | 3.7 |
| Male | 2,011 | 155 | 7.7 | 2,005 | 135 | 6.7 | 427 | 25 | 5.9 | 62 | 14 | 22.6 | 421 | 10 | 2.4 | 62 | 4 | 6.5 | 58 | 7 | 12.1 | 58 | 2 | 3.4 |
| Female | 1,510 | 74 | 4.9 | 1,499 | 68 | 4.5 | 334 | 16 | 4.8 | 26 | 2 | 7.7 | 324 | 10 | 3.1 | 25 | 2 | 8.0 | 22 | 1 | 4.5 | 22 | 1 | 4.5 |
| Transgender | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 |
| Torrance HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 583 | 31 | 5.3 | 579 | 21 | 3.6 | 326 | 13 | 4.0 | 49 | 5 | 10.2 | 322 | 5 | 1.6 | 48 | 2 | 4.2 | 46 | 1 | 2.2 | 45 | 0 | 0.0 |
| Male | 412 | 29 | 7.0 | 409 | 20 | 4.9 | 225 | 11 | 4.9 | 37 | 5 | 13.5 | 222 | 4 | 1.8 | 36 | 2 | 5.6 | 35 | 1 | 2.9 | 34 | 0 | 0.0 |
| Female | 171 | 2 | 1.2 | 170 | 1 | 0.6 | 101 | 2 | 2.0 | 12 | 0 | 0.0 | 100 | 1 | 1.0 | 12 | 0 | 0.0 | 11 | 0 | 0.0 | 11 | 0 | 0.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |
| Curtis Tucker HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 3,431 | 180 | 5.2 | 3,394 | 156 | 4.6 | 1,707 | 52 | 3.0 | 202 | 16 | 7.9 | 1,672 | 32 | 1.9 | 197 | 7 | 3.6 | 183 | 10 | 5.5 | 180 | 5 | 2.8 |
| Male | 2,039 | 134 | 6.6 | 2,026 | 119 | 5.9 | 1,013 | 33 | 3.3 | 102 | 12 | 11.8 | 1,000 | 22 | 2.2 | 101 | 4 | 4.0 | 94 | 7 | 7.4 | 93 | 3 | 3.2 |
| Female | 1,391 | 46 | 3.3 | 1,367 | 37 | 2.7 | 693 | 19 | 2.7 | 99 | 4 | 4.0 | 671 | 10 | 1.5 | 95 | 3 | 3.2 | 88 | 3 | 3.4 | 86 | 2 | 2.3 |
| Transgender | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 |
| Whittier HC | | | | | | | | | | | | | | | | | | | | | | | | |
| Total | 1,276 | 41 | 3.2 | 1,266 | 17 | 1.3 | 750 | 25 | 3.3 | 161 | 8 | 5.0 | 742 | 5 | 0.7 | 159 | 2 | 1.3 | 146 | 3 | 2.1 | 146 | 1 | 0.7 |
| Male | 872 | 31 | 3.6 | 865 | 8 | 0.9 | 508 | 23 | 4.5 | 110 | 8 | 7.3 | 502 | 4 | 0.8 | 109 | 2 | 1.8 | 103 | 3 | 2.9 | 103 | 1 | 1.0 |
| Female | 404 | 10 | 2.5 | 401 | 9 | 2.2 | 242 | 2 | 0.8 | 51 | 0 | 0.0 | 240 | 1 | 0.4 | 50 | 0 | 0.0 | 43 | 0 | 0.0 | 43 | 0 | 0.0 |
| Transgender | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- |

¹Positive at both urogenital and pharyngeal sites on the same visit

²Positive at both urogenital and rectal sites on the same visit

³Positive at both pharyngeal and rectal sites on the same visit

⁴Positive at urogenital, pharyngeal, and rectal sites on the same visit


*Anatomic site categories are not mutually exclusive (eg. Infections listed under the concomitant category "urogenital/rectal" are also included under the overall categories "urogenital" and "rectal")

**"Visits Tested" for gonorrhea refers to number of patient visits and not individual tests (ie. a patient who is tested for urethral, pharyngeal, and rectal gonorrhea on the same visit is only counted as one visit tested under "Total" gonorrhea)

†Tested at the specified anatomic sites on the same visit


Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

**Figure 1.19 Urogenital Chlamydia Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.20 Rectal Chlamydia Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.21 Urogenital Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.22 Pharyngeal Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

**Figure 1.23 Rectal Gonorrhea Prevalence by Gender/Gender of Sex Partners, 2006-2010
All Health Centers (HC)**




Note: MSM/W and MSW specific prevalence not available for 2006, as Clinic Intake Form data collection on gender of sex partners started in latter half of that year.

Figure 1.24. Distribution of Anatomic Site of Infection among All Patients with Chlamydia Infection
January - December 2010
All Health Centers




Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Figure 1.25. Distribution of Anatomic Site of Infection among All Patients with Gonorrhea Infection
 January - December 2010
 All Health Centers



Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested


Figure 1.26. Distribution of Anatomic Site of Infection among MSM/W* with Chlamydia Infection
 January - December 2010
 All Health Centers



*MSM/W=men who have sex with men, or men who have sex with men and women

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested


Figure 1.27. Distribution of Anatomic Site of Infection among MSM/W* with Gonorrhea Infection
 January - December 2010
 All Health Centers



*MSM/W=men who have sex with men, or men who have sex with men and women


Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Figure 1.28. Distribution of Anatomic Site of Infection among Female Patients with Chlamydia Infection
 January - December 2010
 All Health Centers



Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Figure 1.29. Distribution of Anatomic Site of Infection among Female Patients with Gonorrhea Infection
 January - December 2010
 All Health Centers



Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Table 1.8 Patient Visits and Tests
 Patients Diagnosed with New HIV and Other STD on Same Visit
 January - December 2010, All Health Centers

| | | Chlamydia, Gonorrhea, Syphilis and HIV Testing | | | | | | | | | | | | | | |
|---------------------------------|--|--|-------|------------------------------------|-------|------------------|-----------------|-----------------|--------------------------------|------------------------------------|--------------------------------|-----------|----|----------------|----|------|
| | | | | | | | | | | Concomitant Infection with New HIV | | | | | | |
| | | Total STD Visits n % | | No. Visits with STD Test n % | | HIV | | | | Chlamydia | | Gonorrhea | | Early Syphilis | | |
| | | | | | | Visits Tested | Positive n % | New HIV* n % | Positive ¹ n %** | Positive ² n %** | Positive ³ n %** | | | | | |
| Total*** | | 43,973 | 100.0 | 28,275 | 100.0 | 23,285 | 177 | 0.8 | 131 | 0.6 | 17 | 13.0 | 27 | 20.6 | 12 | 9.2 |
| Gender | | | | | | | | | | | | | | | | |
| Male | | 28,441 | 64.7 | 17,959 | 63.5 | 14,921 | 160 | 1.1 | 121 | 0.8 | 17 | 14.0 | 27 | 22.3 | 12 | 9.9 |
| Female | | 15,424 | 35.1 | 10,271 | 36.3 | 8,328 | 17 | 0.2 | 10 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | | 108 | 0.2 | 45 | 0.2 | 36 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| Gender of Sex Partners†† | | | | | | | | | | | | | | | | |
| MSM/W | | 4,963 | 11.3 | 3,000 | 10.6 | 2,225 | 125 | 5.6 | 94 | 4.2 | 16 | 17.0 | 24 | 25.5 | 9 | 9.6 |
| MSW | | 22,210 | 50.5 | 14,370 | 50.8 | 12,244 | 29 | 0.2 | 22 | 0.2 | 1 | 4.5 | 3 | 13.6 | 3 | 13.6 |
| WSM | | 13,416 | 30.5 | 9,055 | 32.0 | 7,397 | 13 | 0.2 | 9 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| WSW/M | | 1,104 | 2.5 | 767 | 2.7 | 605 | 2 | 0.3 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| Unknown | | 2,280 | 5.2 | 1,083 | 3.8 | 814 | 8 | 1.0 | 6 | 0.7 | 1 | 16.7 | 1 | 16.7 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | |
| 0-14 | | 82 | 0.2 | 47 | 0.2 | 42 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| 15-19 | | 3,636 | 8.3 | 2,557 | 9.0 | 2,165 | 10 | 0.5 | 8 | 0.4 | 1 | 12.5 | 1 | 12.5 | 0 | 0.0 |
| 20-24 | | 9,737 | 22.1 | 6,611 | 23.4 | 5,573 | 39 | 0.7 | 32 | 0.6 | 7 | 21.9 | 13 | 40.6 | 5 | 15.6 |
| 25-29 | | 8,691 | 19.8 | 5,678 | 20.1 | 4,813 | 34 | 0.7 | 25 | 0.5 | 2 | 8.0 | 9 | 36.0 | 4 | 16.0 |
| 30-34 | | 6,272 | 14.3 | 3,968 | 14.0 | 3,271 | 34 | 1.0 | 25 | 0.8 | 3 | 12.0 | 2 | 8.0 | 3 | 12.0 |
| 35-44 | | 8,123 | 18.5 | 5,058 | 17.9 | 4,044 | 38 | 0.9 | 28 | 0.7 | 3 | 10.7 | 2 | 7.1 | 0 | 0.0 |
| 45-54 | | 5,148 | 11.7 | 3,096 | 10.9 | 2,451 | 13 | 0.5 | 9 | 0.4 | 1 | 11.1 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | | 1,798 | 4.1 | 1,019 | 3.6 | 761 | 7 | 0.9 | 3 | 0.4 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 65+ | | 486 | 1.1 | 241 | 0.9 | 165 | 2 | 1.2 | 1 | 0.6 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | |
| White | | 6,935 | 15.8 | 4,472 | 15.8 | 3,748 | 18 | 0.5 | 15 | 0.4 | 2 | 13.3 | 4 | 26.7 | 0 | 0.0 |
| African American | | 17,464 | 39.7 | 12,220 | 43.2 | 9,832 | 74 | 0.8 | 52 | 0.5 | 8 | 15.4 | 11 | 21.2 | 10 | 19.2 |
| Hispanic | | 15,942 | 36.3 | 9,283 | 32.8 | 7,764 | 76 | 1.0 | 55 | 0.7 | 7 | 12.7 | 10 | 18.2 | 2 | 3.6 |
| Asian | | 1,890 | 4.3 | 1,191 | 4.2 | 989 | 3 | 0.3 | 3 | 0.3 | 0 | 0.0 | 2 | 66.7 | 0 | 0.0 |
| Native American | | 109 | 0.2 | 75 | 0.3 | 67 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| Other | | 1,410 | 3.2 | 904 | 3.2 | 772 | 5 | 0.6 | 5 | 0.6 | 0 | 0.0 | 1 | 20.0 | 0 | 0.0 |
| Unknown | | 223 | 0.5 | 130 | 0.5 | 113 | 1 | 0.9 | 1 | 0.9 | 0 | 0.0 | 1 | 100.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | |
| New Visit | | 27,497 | 62.5 | 26,291 | 93.0 | 22,661 | 158 | 0.7 | 124 | 0.5 | 17 | 13.7 | 26 | 21.0 | 12 | 9.7 |
| Return Visit | | 16,476 | 37.5 | 1,984 | 7.0 | 624 | 19 | 3.0 | 7 | 1.1 | 0 | 0.0 | 1 | 14.3 | 0 | 0.0 |

¹Positive for both chlamydia and new HIV; ²Positive for both gonorrhea and new HIV; ³Positive for both confirmed early syphilis and new HIV

*Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

**Percentage of new HIV cases

***Among new HIV cases, 10 were diagnosed with HIV and more than one other STD on the same visit: 5 with chlamydia and gonorrhea infections; 2 with chlamydia and early syphilis infections; 2 with gonorrhea and early syphilis infections; and 1 with chlamydia, gonorrhea, and early syphilis infections

†Transgendered in total visits include 36 male-to-female (MTF) and 72 unspecified

††Based on self-reported gender of sexual partners; "MSM/W" = men who have sex with men, or men who have sex with men and women; "MSW" = men who have sex with women; "WSM" = women who have sex with men; "WSW/M" = women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for HIV refers to number of individual visits and not individual tests (ie. a patient who is tested with an HIV EIA and Western Blot on the same visit is only counted as one visit tested)

Table 1.9 Patient Visits and Tests by Gender and Age
 Patients Diagnosed with New HIV and Other STD on Same Visit
 January - December 2010, All Health Centers

| Chlamydia, Gonorrhea, Syphilis and HIV Testing | | | | | | | | | | | | | | | |
|--|-------------------------|-------|------------------------------------|-------|------------------|----------|-----|----------|-----|------------------------------------|------|-----------|------|----------------|------|
| | Total STD Visits n % | | No. Visits with STD Test n % | | HIV | | | | | Concomitant Infection with New HIV | | | | | |
| | | | | | Visits Tested | Positive | | New HIV* | | Chlamydia | | Gonorrhea | | Early Syphilis | |
| | | | | | | n | % | n | % | n | % | n | %** | n | %** |
| Total*** | 43,973 | 100.0 | 28,275 | 100.0 | 23,285 | 177 | 0.8 | 131 | 0.6 | 17 | 13.0 | 27 | 20.6 | 12 | 9.2 |
| Male | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | |
| 0-14 | 28 | 0.1 | 13 | 0.1 | 13 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| 15-19 | 1625 | 5.7 | 1122 | 6.2 | 984 | 7 | 0.7 | 6 | 0.6 | 1 | 16.7 | 1 | 16.7 | 0 | 0.0 |
| 20-24 | 5837 | 20.5 | 3881 | 21.6 | 3,342 | 38 | 1.1 | 31 | 0.9 | 7 | 22.6 | 13 | 41.9 | 5 | 16.1 |
| 25-29 | 6015 | 21.1 | 3851 | 21.4 | 3,308 | 34 | 1.0 | 25 | 0.8 | 2 | 8.0 | 9 | 36.0 | 4 | 16.0 |
| 30-34 | 4349 | 15.3 | 2735 | 15.2 | 2,270 | 30 | 1.3 | 22 | 1.0 | 3 | 13.6 | 2 | 9.1 | 3 | 13.6 |
| 35-44 | 5557 | 19.5 | 3415 | 19.0 | 2,732 | 35 | 1.3 | 27 | 1.0 | 3 | 11.1 | 2 | 7.4 | 0 | 0.0 |
| 45-54 | 3458 | 12.2 | 2054 | 11.4 | 1,611 | 10 | 0.6 | 8 | 0.5 | 1 | 12.5 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 1253 | 4.4 | 729 | 4.1 | 546 | 4 | 0.7 | 1 | 0.2 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 319 | 1.1 | 159 | 0.9 | 115 | 2 | 1.7 | 1 | 0.9 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | -- | 0 | -- | 0 | -- | 0 | -- | 0 | -- |
| Subtotal | 28,441 | 100.0 | 17,959 | 100.0 | 14,921 | 160 | 1.1 | 121 | 0.8 | 17 | 14.0 | 27 | 22.3 | 12 | 9.9 |
| Female | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | |
| 0-14 | 54 | 0.4 | 34 | 0.3 | 29 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| 15-19 | 2007 | 13.0 | 1435 | 14.0 | 1,181 | 3 | 0.3 | 2 | 0.2 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 3883 | 25.2 | 2720 | 26.5 | 2,222 | 1 | 0.0 | 1 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 2657 | 17.2 | 1822 | 17.7 | 1,501 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| 30-34 | 1902 | 12.3 | 1225 | 11.9 | 995 | 4 | 0.4 | 3 | 0.3 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 2532 | 16.4 | 1627 | 15.8 | 1,300 | 3 | 0.2 | 1 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 1677 | 10.9 | 1036 | 10.1 | 835 | 3 | 0.4 | 1 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 545 | 3.5 | 290 | 2.8 | 215 | 3 | 1.4 | 2 | 0.9 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 167 | 1.1 | 82 | 0.8 | 50 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | -- | 0 | -- | 0 | -- | 0 | -- | 0 | -- |
| Subtotal | 15,424 | 100.0 | 10,271 | 100.0 | 8,328 | 17 | 0.2 | 10 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | 108 | 100.0 | 45 | 100.0 | 36 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- |

¹Positive for both chlamydia and new HIV; ²Positive for both gonorrhea and new HIV; ³Positive for both confirmed early syphilis and new HIV

*Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

**Percentage of new HIV cases

***Among male new HIV cases, 10 were diagnosed with HIV and more than one other STD on the same visit: 5 with chlamydia and gonorrhea infections; 2 with chlamydia and early syphilis infections; 2 with gonorrhea and early syphilis infections; and 1 with chlamydia, gonorrhea, and early syphilis infections

†Transgendered in total visits include 36 male-to-female (MTF) and 72 unspecified

Note: "Visits Tested" for HIV refers to number of individual visits and not individual tests (ie. a patient who is tested with an HIV EIA and Western Blot on the same visit is only counted as one visit tested)

Table 1.10 Patient Visits and Tests by Gender and Race/Ethnicity
 Patients Diagnosed with New HIV and Other STD on Same Visit
 January - December 2010, All Health Centers

| Chlamydia, Gonorrhea, Syphilis and HIV Testing | | | | | | | | | | | | | | | | |
|--|-------------------------|-------|------------------------------------|-------|--------|-----|--|-----|-----|----|------------------------------------|--------------------------------|--------------------------------|----|----------------|--|
| | Total STD Visits n % | | No. Visits with STD Test n % | | | | HIV Visits Tested Positive n % New HIV* n % | | | | Concomitant Infection with New HIV | | | | | |
| | | | | | | | | | | | Chlamydia | | Gonorrhea | | Early Syphilis | |
| | | | | | | | | | | | Positive ¹ n %** | Positive ² n %** | Positive ³ n %** | | | |
| Total*** | 43,973 | 100.0 | 28,275 | 100.0 | 23,285 | 177 | 0.8 | 131 | 0.6 | 17 | 13.0 | 27 | 20.6 | 12 | 9.2 | |
| Male | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | |
| White | 5138 | 18.1 | 3307 | 18.4 | 2,761 | 18 | 0.7 | 15 | 0.5 | 2 | 13.3 | 4 | 26.7 | 0 | 0.0 | |
| African American | 10133 | 35.6 | 6944 | 38.7 | 5,635 | 66 | 1.2 | 47 | 0.8 | 8 | 17.0 | 11 | 23.4 | 10 | 21.3 | |
| Hispanic | 10869 | 38.2 | 6272 | 34.9 | 5,293 | 68 | 1.3 | 51 | 1.0 | 7 | 13.7 | 10 | 19.6 | 2 | 3.9 | |
| Asian | 1167 | 4.1 | 730 | 4.1 | 616 | 2 | 0.3 | 2 | 0.3 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | |
| Native American | 65 | 0.2 | 47 | 0.3 | 41 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- | |
| Other | 902 | 3.2 | 567 | 3.2 | 496 | 5 | 1.0 | 5 | 1.0 | 0 | 0.0 | 1 | 20.0 | 0 | 0.0 | |
| Unknown | 167 | 0.6 | 92 | 0.5 | 79 | 1 | 1.3 | 1 | 1.3 | 0 | 0.0 | 1 | 100.0 | 0 | 0.0 | |
| Subtotal | 28,441 | 100.0 | 17,959 | 100.0 | 14,921 | 160 | 1.1 | 121 | 0.8 | 17 | 14.0 | 27 | 22.3 | 12 | 9.9 | |
| Female | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | |
| White | 1793 | 11.6 | 1163 | 11.3 | 985 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- | |
| African American | 7321 | 47.5 | 5269 | 51.3 | 4,195 | 8 | 0.2 | 5 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | |
| Hispanic | 4994 | 32.4 | 2979 | 29.0 | 2,442 | 8 | 0.3 | 4 | 0.2 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | |
| Asian | 716 | 4.6 | 458 | 4.5 | 370 | 1 | 0.3 | 1 | 0.3 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | |
| Native American | 44 | 0.3 | 28 | 0.3 | 26 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- | |
| Other | 500 | 3.2 | 336 | 3.3 | 276 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- | |
| Unknown | 56 | 0.4 | 38 | 0.4 | 34 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- | |
| Subtotal | 15,424 | 100.0 | 10,271 | 100.0 | 8,328 | 17 | 0.2 | 10 | 0.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | |
| Transgender[†] | 108 | 100.0 | 45 | 100.0 | 36 | 0 | 0.0 | 0 | 0.0 | 0 | -- | 0 | -- | 0 | -- | |

¹Positive for both chlamydia and new HIV; ²Positive for both gonorrhea and new HIV; ³Positive for both confirmed early syphilis and new HIV

*Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form


**Percentage of new HIV cases

***Among male new HIV cases, 10 were diagnosed with HIV and more than one other STD on the same visit: 5 with chlamydia and gonorrhea infections; 2 with chlamydia and early syphilis infections; 2 with gonorrhea and early syphilis infections; and 1 with chlamydia, gonorrhea, and early syphilis infections

[†]Transgendered in total visits include 36 male-to-female (MTF) and 72 unspecified

Note: "Visits Tested" for HIV refers to number of individual visits and not individual tests (ie. a patient who is tested with an HIV EIA and Western Blot on the same visit is only counted as one visit tested)

Figure 1.30. Other STDs Diagnosed on Same Visit as HIV among Patients with New HIV
 January - December 2010
 All Health Centers




*MSM/W = men who have sex with men, or men who have sex with men and women; MSW = men who have sex with women

**Chlamydia, Gonorrhea, or Early Syphilis


Note1: Number of new HIV cases: Overall=131; MSM/W=94; MSW=22; Females=0

Figure 1.31. Distribution of Other STDs Diagnosed on Same Visit as HIV among Patients with New HIV
 January - December 2010
 All Health Centers




Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Figure 1.32. Distribution of Anatomic Site of Chlamydia Infection Diagnosed on Same Visit as HIV among Patients with New HIV
January - December 2010
All Health Centers




Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Figure 1.33. Distribution of Anatomic Site of Gonorrhea Infection Diagnosed on Same Visit as HIV among Patients with New HIV
 January - December 2010
 All Health Centers



Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested


Figure 1.34. Distribution of Other STDs Diagnosed on Same Visit as HIV among MSM/W* Patients with New HIV
 January - December 2010
 All Health Centers



*MSM/W = men who have sex with men, or men who have sex with men and women

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested


Figure 1.35. Distribution of Anatomic Site of Chlamydia Infection Diagnosed on Same Visit as HIV among MSM/W* Patients with New HIV
January - December 2010
All Health Centers



*MSM/W = men who have sex with men, or men who have sex with men and women

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

Figure 1.36. Distribution of Anatomic Site of Gonorrhea Infection Diagnosed on Same Visit as HIV among MSM/W* Patients with New HIV
 January - December 2010
 All Health Centers



*MSM/W = men who have sex with men, or men who have sex with men and women

Note: Anatomic site of distribution may not be representative of actual prevalence but an artifact of sites tested

**Table 1.11 Herpes Simplex Virus Testing of Symptomatic Patients*
Summary by Gender
January - December 2010**

| | Herpes Simplex Virus | | | | | | | | | | | | | |
|------------------------------|----------------------|-----|------------------|-------|------------------|-------|-----------|------------------|----|------------------|-------|------------------|-------|-------|
| | Male | | | | | | Female | | | | | | | |
| | Total Tests | | | HSV 1 | | HSV 2 | | Total Tests | | | HSV 1 | | HSV 2 | |
| No. Tests | Positive Tests n | % | Positive Tests n | % | Positive Tests n | % | No. Tests | Positive Tests n | % | Positive Tests n | % | Positive Tests n | % | |
| All Centers | 432 | 146 | 33.8% | 27 | 6.3% | 119 | 27.5% | 209 | 69 | 33.0% | 22 | 10.5% | 47 | 22.5% |
| Antelope Valley HC | 9 | 1 | 11.1% | 1 | 11.1% | 0 | 0.0% | 9 | 3 | 33.3% | 1 | 11.1% | 2 | 22.2% |
| Simms Mann HC | 15 | 3 | 20.0% | 1 | 6.7% | 2 | 13.3% | 5 | 1 | 20.0% | 1 | 20.0% | 0 | 0.0% |
| Central HC | 48 | 21 | 43.8% | 7 | 14.6% | 14 | 29.2% | 17 | 5 | 29.4% | 1 | 5.9% | 4 | 23.5% |
| Hollywood-Wilshire HC | 84 | 23 | 27.4% | 3 | 3.6% | 20 | 23.8% | 26 | 11 | 42.3% | 3 | 11.5% | 8 | 30.8% |
| Monrovia HC | 20 | 9 | 45.0% | 3 | 15.0% | 6 | 30.0% | 9 | 2 | 22.2% | 1 | 11.1% | 1 | 11.1% |
| North Hollywood HC | 69 | 18 | 26.1% | 3 | 4.3% | 15 | 21.7% | 27 | 11 | 40.7% | 4 | 14.8% | 7 | 25.9% |
| Pomona HC | 12 | 3 | 25.0% | 0 | 0.0% | 3 | 25.0% | 5 | 1 | 20.0% | 1 | 20.0% | 0 | 0.0% |
| South HC | 48 | 18 | 37.5% | 0 | 0.0% | 18 | 37.5% | 40 | 15 | 37.5% | 7 | 17.5% | 8 | 20.0% |
| Ruth Temple HC | 43 | 21 | 48.8% | 3 | 7.0% | 18 | 41.9% | 18 | 7 | 38.9% | 1 | 5.6% | 6 | 33.3% |
| Torrance HC | 11 | 5 | 45.5% | 2 | 18.2% | 3 | 27.3% | 8 | 2 | 25.0% | 1 | 12.5% | 1 | 12.5% |
| Curtis Tucker HC | 51 | 18 | 35.3% | 3 | 5.9% | 15 | 29.4% | 36 | 9 | 25.0% | 0 | 0.0% | 9 | 25.0% |
| Whittier HC | 22 | 6 | 27.3% | 1 | 4.5% | 5 | 22.7% | 9 | 2 | 22.2% | 1 | 11.1% | 1 | 11.1% |

*Direct testing performed by PCR of symptomatic site

**Table 1.12 Summary of Paps Performed by Health Center
All Health Centers (HC)**

| | All Clinics | | Antelope Valley | | Simms Mann | | Central | | Hollywood/ Wilshire | | North Hollywood | | Monrovia | |
|--|-------------|-------|-----------------|-------|------------|-------|---------|-------|------------------------|-------|-----------------|-------|----------|-------|
| | n | % | n | % | n | % | n | % | n | % | n | % | n | % |
| Total Number of Paps | 1,298 | 100.0 | 89 | 100.0 | 50 | 100.0 | 208 | 100.0 | 36 | 100.0 | 168 | 100.0 | 65 | 100.0 |
| Abnormal Paps | 171 | 13.2 | 15 | 16.9 | 7 | 14.0 | 22 | 10.6 | 5 | 13.9 | 20 | 11.9 | 13 | 20.0 |
| Abnormal Results: | | | | | | | | | | | | | | |
| Women ≤ 20 years* ASC-US (Atypical Squamous Cells of Unetermined Significance) | 18 | 10.5 | 0 | 0.0 | 0 | 0.0 | 2 | 9.1 | 0 | 0.0 | 1 | 5.0 | 2 | 15.4 |
| Women ≤ 20 years* LSIL (Low-grade Squamous Intraepithelial Lesion) | 11 | 6.4 | 4 | 26.7 | 0 | 0.0 | 2 | 9.1 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| ≥ 21 years ASC-US w/ neg. HPV | 33 | 19.3 | 2 | 13.3 | 2 | 28.6 | 4 | 18.2 | 1 | 20.0 | 3 | 15.0 | 2 | 15.4 |
| ≥ 21 years ASC-US w/ no HPV | 5 | 2.9 | 0 | 0.0 | 0 | 0.0 | 1 | 4.5 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| ≥ 21 years ASC-US w/ pos. HPV | 46 | 26.9 | 3 | 20.0 | 3 | 42.9 | 5 | 22.7 | 1 | 20.0 | 6 | 30.0 | 4 | 30.8 |
| Women ≥ 21 years LSIL (Low-grade Squamous Intraepithelial Lesion) | 46 | 26.9 | 4 | 26.7 | 2 | 28.6 | 7 | 31.8 | 2 | 40.0 | 7 | 35.0 | 4 | 30.8 |
| High-grade Squamous Intraepithelial Lesion | 10 | 5.8 | 1 | 6.7 | 0 | 0.0 | 1 | 4.5 | 0 | 0.0 | 2 | 10.0 | 1 | 7.7 |
| Atypical Squamous Cells - cannot exclude HSIL | 1 | 0.6 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 5.0 | 0 | 0.0 |
| Atypical Glandular Cells - not otherwise specified | 1 | 0.6 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 1 | 20.0 | 0 | 0.0 | 0 | 0.0 |
| Atypical Glandular Cells - favor neoplasia | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Adenocarcinoma in situ | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Adenocarcinoma | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Squamous Cell Carcinoma | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |


*From May 2010, the Pap protocol changed to performing Paps every 2 years starting age 21 (previously Paps were performed yearly from 3 years after first sexual intercourse)

**Table 1.12 Summary of Paps Performed by Health Center
All Health Centers (HC)**

| | Pomona | | South | | R. Temple | | Torrance | | C. Tucker | | Whittier | |
|--|--------|-------|-------|-------|-----------|-------|----------|-------|-----------|-------|----------|-------|
| | n | % | n | % | n | % | n | % | n | % | n | % |
| Total Number of Paps | 53 | 100.0 | 245 | 100.0 | 183 | 100.0 | 29 | 100.0 | 130 | 100.0 | 42 | 100.0 |
| Abnormal Paps | 5 | 9.4 | 30 | 12.2 | 21 | 11.5 | 3 | 10.3 | 25 | 19.2 | 5 | 11.9 |
| Abnormal Results: | | | | | | | | | | | | |
| Women ≤ 20 years* ASC-US (Atypical Squamous Cells of Uneterminated Significance) | 0 | 0.0 | 5 | 16.7 | 3 | 14.3 | 1 | 33.3 | 3 | 12.0 | 1 | 20.0 |
| Women ≤ 20 years* LSIL (Low-grade Squamous Intraepithelial Lesion) | 1 | 20.0 | 2 | 6.7 | 2 | 9.5 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| ≥ 21 years ASC-US w/ neg. HPV | 1 | 20.0 | 9 | 30.0 | 4 | 19.0 | 1 | 33.3 | 4 | 16.0 | 0 | 0.0 |
| ≥ 21 years ASC-US w/ no HPV | 0 | 0.0 | 0 | 0.0 | 1 | 4.8 | 0 | 0.0 | 3 | 12.0 | 0 | 0.0 |
| ≥ 21 years ASC-US w/ pos. HPV | 1 | 20.0 | 7 | 23.3 | 8 | 38.1 | 1 | 33.3 | 6 | 24.0 | 1 | 20.0 |
| Women ≥ 21 years LSIL (Low-grade Squamous Intraepithelial Lesion) | 2 | 40.0 | 6 | 20.0 | 3 | 14.3 | 0 | 0.0 | 5 | 20.0 | 3 | 60.0 |
| High-grade Squamous Intraepithelial Lesion | 0 | 0.0 | 1 | 3.3 | 0 | 0.0 | 0 | 0.0 | 4 | 16.0 | 0 | 0.0 |
| Atypical Squamous Cells - cannot exclude HSIL | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Atypical Glandular Cells - not otherwise specified | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Atypical Glandular Cells - favor neoplasia | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Adenocarcinoma in situ | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Adenocarcinoma | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |
| Squamous Cell Carcinoma | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 | 0.0 |


*From May 2010, the Pap protocol changed to performing Paps every 2 years starting age 21 (previously Paps were performed yearly from 3 years after first sexual intercourse)

Figure 1.37. Clinical Diagnoses among Males at New Visits*
January - December 2010
All Health Centers



*Visit classification and clinical diagnoses as indicated by the Clinic Intake Form

Figure 1.38
 Clinical Diagnoses among Females at New Visits*
 January - December 2010
 All Health Centers



*Visit classification and clinical diagnoses as indicated by the Clinic Intake Form

**PID = Pelvic Inflammatory Disease

***Vaginitis (NOS) = Vaginitis (not otherwise specified)

Sexually Transmitted Disease Clinic Morbidity Report, 2010

II: Antelope Valley Health Center

Table 2.1 Patient Visits and Tests, January - December 2010, Antelope Valley Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|-------------------------|------------------|-----------------|-------------------|-------------|-------|---|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | Early Syphilis** n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 1,613 | 100.0 | 1,195 | 100.0 | 1,149 | 173 | 15.1 | 1,156 | 50 | 4.3 | 1,071 | 33 | 3.1 | 0 | 0.0 | 1,045 | 4 | 0.4 | 3 | 0.3 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 740 | 45.9 | 541 | 45.3 | 522 | 87 | 16.7 | 525 | 33 | 6.3 | 500 | 19 | 3.8 | 0 | 0.0 | 487 | 3 | 0.6 | 2 | 0.4 |
| Female | 873 | 54.1 | 654 | 54.7 | 627 | 86 | 13.7 | 631 | 17 | 2.7 | 571 | 14 | 2.5 | 0 | 0.0 | 558 | 1 | 0.2 | 1 | 0.2 |
| Transgender† | 0 | 0.0 | 0 | 0.0 | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | -- | 0 | 0 | -- | 0 | -- |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 83 | 5.1 | 64 | 5.4 | 60 | 15 | 25.0 | 61 | 10 | 16.4 | 58 | 8 | 13.8 | 0 | 0.0 | 53 | 3 | 5.7 | 2 | 3.8 |
| MSW | 625 | 38.7 | 459 | 38.4 | 448 | 71 | 15.8 | 450 | 23 | 5.1 | 426 | 8 | 1.9 | 0 | 0.0 | 421 | 0 | 0.0 | 0 | 0.0 |
| WSM | 770 | 47.7 | 581 | 48.6 | 557 | 78 | 14.0 | 561 | 15 | 2.7 | 506 | 8 | 1.6 | 0 | 0.0 | 498 | 1 | 0.2 | 1 | 0.2 |
| WSW/M | 78 | 4.8 | 58 | 4.9 | 57 | 7 | 12.3 | 57 | 2 | 3.5 | 51 | 3 | 5.9 | 0 | 0.0 | 49 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 57 | 3.5 | 33 | 2.8 | 27 | 2 | 7.4 | 27 | 0 | 0.0 | 30 | 6 | 20.0 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 6 | 0.4 | 2 | 0.2 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 302 | 18.7 | 238 | 19.9 | 232 | 50 | 21.6 | 233 | 13 | 5.6 | 218 | 2 | 0.9 | 0 | 0.0 | 216 | 1 | 0.5 | 1 | 0.5 |
| 20-24 | 594 | 36.8 | 446 | 37.3 | 425 | 80 | 18.8 | 429 | 23 | 5.4 | 393 | 6 | 1.5 | 0 | 0.0 | 385 | 2 | 0.5 | 1 | 0.3 |
| 25-29 | 282 | 17.5 | 210 | 17.6 | 208 | 29 | 13.9 | 209 | 11 | 5.3 | 194 | 3 | 1.5 | 0 | 0.0 | 192 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 149 | 9.2 | 104 | 8.7 | 102 | 7 | 6.9 | 102 | 1 | 1.0 | 86 | 2 | 2.3 | 0 | 0.0 | 84 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 165 | 10.2 | 119 | 10.0 | 114 | 3 | 2.6 | 114 | 1 | 0.9 | 111 | 7 | 6.3 | 0 | 0.0 | 106 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 83 | 5.1 | 53 | 4.4 | 47 | 3 | 6.4 | 48 | 1 | 2.1 | 47 | 6 | 12.8 | 0 | 0.0 | 44 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 30 | 1.9 | 22 | 1.8 | 18 | 0 | 0.0 | 18 | 0 | 0.0 | 19 | 7 | 36.8 | 0 | 0.0 | 15 | 1 | 6.7 | 1 | 6.7 |
| 65+ | 2 | 0.1 | 1 | 0.1 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 304 | 18.8 | 214 | 17.9 | 206 | 27 | 13.1 | 206 | 2 | 1.0 | 194 | 6 | 3.1 | 0 | 0.0 | 191 | 0 | 0.0 | 0 | 0.0 |
| African American | 690 | 42.8 | 539 | 45.1 | 519 | 86 | 16.6 | 524 | 39 | 7.4 | 461 | 16 | 3.5 | 0 | 0.0 | 449 | 3 | 0.7 | 2 | 0.4 |
| Hispanic | 438 | 27.2 | 307 | 25.7 | 292 | 40 | 13.7 | 294 | 5 | 1.7 | 285 | 8 | 2.8 | 0 | 0.0 | 278 | 1 | 0.4 | 1 | 0.4 |
| Asian | 28 | 1.7 | 22 | 1.8 | 22 | 3 | 13.6 | 22 | 0 | 0.0 | 22 | 1 | 4.5 | 0 | 0.0 | 21 | 0 | 0.0 | 0 | 0.0 |
| Native American | 3 | 0.2 | 3 | 0.3 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Other | 119 | 7.4 | 86 | 7.2 | 85 | 14 | 16.5 | 85 | 3 | 3.5 | 83 | 0 | 0.0 | 0 | 0.0 | 82 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 31 | 1.9 | 24 | 2.0 | 22 | 3 | 13.6 | 22 | 1 | 4.5 | 23 | 2 | 8.7 | 0 | 0.0 | 21 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit*** | | | | | | | | | | | | | | | | | | | | |
| New Visit | 1,097 | 68.0 | 1,089 | 91.1 | 1,071 | 167 | 15.6 | 1,075 | 47 | 4.4 | 1,034 | 18 | 1.7 | 0 | 0.0 | 1018 | 3 | 0.3 | 3 | 0.3 |
| Return Visit | 516 | 32.0 | 106 | 8.9 | 78 | 6 | 7.7 | 81 | 3 | 3.7 | 37 | 15 | 40.5 | 0 | 0.0 | 27 | 1 | 3.7 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include both male-to-female (MTF) and female-to-male (FTM)

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 2.2 Patient Visits and Tests by Gender and Age, January - December 2010, Antelope Valley Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-------|---|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 1,613 | 100.0 | 1,195 | 100.0 | 1,149 | 173 | 15.1 | 1,156 | 50 | 4.3 | 1,071 | 33 | 3.1 | 0 | 0.0 | 1,045 | 4 | 0.4 | 3 | 0.3 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 2 | 0.3 | 90 | 16.6 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 112 | 15.2 | 217 | 40.1 | 88 | 20 | 22.7 | 88 | 9 | 10.2 | 86 | 2 | 2.3 | 0 | 0.0 | 82 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 286 | 38.8 | 103 | 19.0 | 209 | 44 | 21.1 | 211 | 14 | 6.6 | 198 | 5 | 2.5 | 0 | 0.0 | 194 | 2 | 1.0 | 1 | 0.5 |
| 25-29 | 139 | 18.9 | 40 | 7.4 | 102 | 18 | 17.6 | 103 | 8 | 7.8 | 92 | 0 | 0.0 | 0 | 0.0 | 95 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 62 | 8.4 | 45 | 8.3 | 39 | 1 | 2.6 | 39 | 0 | 0.0 | 38 | 2 | 5.3 | 0 | 0.0 | 37 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 66 | 9.0 | 29 | 5.4 | 44 | 2 | 4.5 | 44 | 1 | 2.3 | 43 | 3 | 7.0 | 0 | 0.0 | 41 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 49 | 6.5 | 16 | 3.0 | 26 | 2 | 7.7 | 26 | 1 | 3.8 | 28 | 3 | 10.7 | 0 | 0.0 | 26 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 20 | 2.7 | 1 | 0.2 | 13 | 0 | 0.0 | 13 | 0 | 0.0 | 14 | 4 | 28.6 | 0 | 0.0 | 11 | 1 | 9.1 | 1 | 9.1 |
| 65+ | 2 | 0.3 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 738 | 100.0 | 541 | 100.0 | 522 | 87 | 16.7 | 525 | 33 | 6.3 | 500 | 19 | 3.8 | 0 | 0.0 | 487 | 3 | 0.6 | 2 | 0.4 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 4 | 0.5 | 2 | 0.3 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 190 | 21.8 | 148 | 22.6 | 144 | 30 | 20.8 | 145 | 4 | 2.8 | 132 | 0 | 0.0 | 0 | 0.0 | 134 | 1 | 0.7 | 1 | 0.7 |
| 20-24 | 306 | 35.1 | 229 | 35.0 | 216 | 36 | 16.7 | 218 | 9 | 4.1 | 195 | 1 | 0.5 | 0 | 0.0 | 191 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 143 | 16.4 | 107 | 16.4 | 106 | 11 | 10.4 | 106 | 3 | 2.8 | 102 | 3 | 2.9 | 0 | 0.0 | 97 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 87 | 10.0 | 64 | 9.8 | 63 | 6 | 9.5 | 63 | 1 | 1.6 | 48 | 0 | 0.0 | 0 | 0.0 | 47 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 99 | 11.3 | 74 | 11.3 | 70 | 1 | 1.4 | 70 | 0 | 0.0 | 68 | 4 | 5.9 | 0 | 0.0 | 65 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 34 | 3.9 | 24 | 3.7 | 21 | 1 | 4.8 | 22 | 0 | 0.0 | 19 | 3 | 15.8 | 0 | 0.0 | 18 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 10 | 1.1 | 6 | 0.9 | 5 | 0 | 0.0 | 5 | 0 | 0.0 | 5 | 3 | 60.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 873 | 100.0 | 654 | 100.0 | 627 | 86 | 13.7 | 631 | 17 | 2.7 | 571 | 14 | 2.5 | 0 | 0.0 | 558 | 1 | 0.2 | 1 | 0.2 |
| Others † | 2 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 2.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Antelope Valley Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-------|---|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 1,613 | 100.0 | 1,195 | 100.0 | 1,149 | 173 | 15.1 | 1,156 | 50 | 4.3 | 1,071 | 33 | 3.1 | | | 1,045 | 4 | 0.4 | 3 | 0.3 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 144 | 19.5 | 101 | 18.7 | 99 | 17 | 17.2 | 99 | 2 | 2.0 | 93 | 4 | 4.3 | 0 | 0.0 | 91 | 0 | 0.0 | 0 | 0.0 |
| African American | 280 | 37.9 | 219 | 40.5 | 210 | 39 | 18.6 | 212 | 25 | 11.8 | 198 | 9 | 4.5 | 0 | 0.0 | 194 | 3 | 1.5 | 2 | 1.0 |
| Hispanic | 222 | 30.1 | 151 | 27.9 | 145 | 20 | 13.8 | 146 | 3 | 2.1 | 139 | 3 | 2.2 | 0 | 0.0 | 136 | 0 | 0.0 | 0 | 0.0 |
| Asian | 15 | 2.0 | 13 | 2.4 | 13 | 1 | 7.7 | 13 | 0 | 0.0 | 13 | 1 | 7.7 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| Native American | 1 | 0.1 | 1 | 0.2 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 62 | 8.4 | 45 | 8.3 | 45 | 9 | 20.0 | 45 | 3 | 6.7 | 45 | 0 | 0.0 | 0 | 0.0 | 44 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 14 | 1.9 | 11 | 2.0 | 9 | 1 | 11.1 | 9 | 0 | 0.0 | 11 | 2 | 18.2 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 738 | 100.0 | 541 | 100.0 | 522 | 87 | 16.7 | 525 | 33 | 6.3 | 500 | 19 | 3.8 | 0 | 0.0 | 487 | 3 | 0.6 | 2 | 0.4 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 160 | 18.3 | 113 | 17.3 | 107 | 10 | 9.3 | 107 | 0 | 0.0 | 101 | 2 | 2.0 | 0 | 0.0 | 100 | 0 | 0.0 | 0 | 0.0 |
| African American | 410 | 47.0 | 320 | 48.9 | 309 | 47 | 15.2 | 312 | 14 | 4.5 | 263 | 7 | 2.7 | 0 | 0.0 | 255 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 216 | 24.7 | 156 | 23.9 | 147 | 20 | 13.6 | 148 | 2 | 1.4 | 146 | 5 | 3.4 | 0 | 0.0 | 142 | 1 | 0.7 | 1 | 0.7 |
| Asian | 13 | 1.5 | 9 | 1.4 | 9 | 2 | 22.2 | 9 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Native American | 2 | 0.2 | 2 | 0.3 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Other | 55 | 6.3 | 41 | 6.3 | 40 | 5 | 12.5 | 40 | 0 | 0.0 | 38 | 0 | 0.0 | 0 | 0.0 | 38 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 17 | 2.0 | 13 | 2.0 | 13 | 2 | 15.4 | 13 | 1 | 7.7 | 12 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 873 | 100.0 | 654 | 100.0 | 627 | 86 | 13.7 | 631 | 17 | 2.7 | 571 | 14 | 2.5 | 0 | 0.0 | 558 | 1 | 0.2 | 1 | 0.2 |
| Others † | 2 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 2.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Antelope Valley Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 2.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Antelope Valley Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

III: Simms Mann Health and Wellness Center

Table 3.1 Patient Visits and Tests, January - December 2010, Simms Mann Health & Wellness Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|---|------------------|-------------------------------|-------------------|-------------|---|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n | % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 806 | 100.0 | 714 | 100.0 | 700 | 34 | 4.9 | 701 | 12 | 1.7 | 648 | 7 | 1.1 | 1 | 0.2 | 613 | 1 | 0.2 | 1 | 0.2 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 624 | 77.4 | 556 | 77.9 | 548 | 25 | 4.6 | 548 | 9 | 1.6 | 502 | 5 | 1.0 | 0 | 0.0 | 476 | 1 | 0.2 | 1 | 0.2 |
| Female | 181 | 22.5 | 158 | 22.1 | 152 | 9 | 5.9 | 153 | 3 | 2.0 | 146 | 2 | 1.4 | 1 | 0.7 | 137 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | 1 | 0.1 | 0 | 0.0 | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | -- | 0 | 0 | -- | 0 | -- |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 99 | 12.3 | 80 | 11.2 | 76 | 3 | 3.9 | 76 | 8 | 10.5 | 73 | 5 | 6.8 | 0 | 0.0 | 65 | 1 | 1.5 | 1 | 1.5 |
| MSW | 510 | 63.3 | 463 | 64.8 | 459 | 21 | 4.6 | 459 | 1 | 0.2 | 418 | 0 | 0.0 | 0 | 0.0 | 401 | 0 | 0.0 | 0 | 0.0 |
| WSM | 174 | 21.6 | 152 | 21.3 | 147 | 9 | 6.1 | 148 | 3 | 2.0 | 140 | 1 | 0.7 | 1 | 0.7 | 132 | 0 | 0.0 | 0 | 0.0 |
| WSW/M | 4 | 0.5 | 4 | 0.6 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 4 | 1 | 25.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 19 | 2.4 | 15 | 2.1 | 15 | 1 | 6.7 | 15 | 0 | 0.0 | 13 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | -- | 0 | 0 | -- | 0 | -- |
| 15-19 | 29 | 3.6 | 28 | 3.9 | 28 | 1 | 3.6 | 28 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 | 22 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 163 | 20.2 | 151 | 21.1 | 150 | 14 | 9.3 | 151 | 3 | 2.0 | 139 | 0 | 0.0 | 0 | 0.0 | 134 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 205 | 25.4 | 177 | 24.8 | 174 | 10 | 5.7 | 174 | 4 | 2.3 | 157 | 0 | 0.0 | 0 | 0.0 | 152 | 1 | 0.7 | 1 | 0.7 |
| 30-34 | 153 | 19.0 | 132 | 18.5 | 131 | 4 | 3.1 | 131 | 1 | 0.8 | 123 | 0 | 0.0 | 0 | 0.0 | 120 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 165 | 20.5 | 148 | 20.7 | 142 | 3 | 2.1 | 142 | 2 | 1.4 | 131 | 5 | 3.8 | 1 | 0.8 | 117 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 68 | 8.4 | 61 | 8.5 | 59 | 2 | 3.4 | 59 | 2 | 3.4 | 58 | 1 | 1.7 | 0 | 0.0 | 54 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 21 | 2.6 | 15 | 2.1 | 15 | 0 | 0.0 | 15 | 0 | 0.0 | 15 | 1 | 6.7 | 0 | 0.0 | 13 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 2 | 0.2 | 2 | 0.3 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 454 | 56.3 | 408 | 57.1 | 404 | 17 | 4.2 | 404 | 8 | 2.0 | 370 | 2 | 0.5 | 0 | 0.0 | 352 | 0 | 0.0 | 0 | 0.0 |
| African American | 116 | 14.4 | 100 | 14.0 | 96 | 5 | 5.2 | 97 | 2 | 2.1 | 85 | 1 | 1.2 | 0 | 0.0 | 78 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 103 | 12.8 | 92 | 12.9 | 87 | 5 | 5.7 | 87 | 2 | 2.3 | 87 | 3 | 3.4 | 0 | 0.0 | 79 | 0 | 0.0 | 0 | 0.0 |
| Asian | 84 | 10.4 | 75 | 10.5 | 75 | 4 | 5.3 | 75 | 0 | 0.0 | 70 | 1 | 1.4 | 1 | 1.4 | 68 | 0 | 0.0 | 0 | 0.0 |
| Native American | 2 | 0.2 | 2 | 0.3 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Other | 37 | 4.6 | 31 | 4.3 | 30 | 1 | 3.3 | 30 | 0 | 0.0 | 29 | 0 | 0.0 | 0 | 0.0 | 29 | 1 | 3.4 | 1 | 3.4 |
| Unknown | 10 | 1.2 | 6 | 0.8 | 6 | 1 | 16.7 | 6 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 684 | 84.9 | 671 | 94.0 | 668 | 34 | 5.1 | 668 | 12 | 1.8 | 623 | 4 | 0.6 | 1 | 0.2 | 598 | 1 | 0.2 | 1 | 0.2 |
| Return Visit | 122 | 15.1 | 43 | 6.0 | 32 | 0 | 0.0 | 33 | 0 | 0.0 | 25 | 3 | 12.0 | 0 | 0.0 | 15 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 1 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 3.2 Patient Visits and Tests by Gender and Age, January - December 2010, Simms Mann Health & Wellness Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-----|-----|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 806 | 100.0 | 714 | 100.0 | 700 | 34 | 4.9 | 701 | 12 | 1.7 | 648 | 7 | 1.1 | 1 | 0.2 | 613 | 1 | 0.2 | 1 | 0.2 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 17 | 2.7 | 16 | 2.9 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 110 | 17.6 | 103 | 18.6 | 16 | 0 | 0.0 | 16 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 166 | 26.6 | 143 | 25.8 | 103 | 9 | 8.7 | 103 | 2 | 1.9 | 96 | 0 | 0.0 | 0 | 0.0 | 91 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 123 | 19.7 | 109 | 19.6 | 141 | 9 | 6.4 | 141 | 2 | 1.4 | 125 | 0 | 0.0 | 0 | 0.0 | 122 | 1 | 0.8 | 1 | 0.8 |
| 30-34 | 130 | 20.8 | 118 | 21.3 | 109 | 3 | 2.8 | 109 | 1 | 0.9 | 102 | 0 | 0.0 | 0 | 0.0 | 100 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 59 | 9.5 | 52 | 9.4 | 114 | 2 | 1.8 | 114 | 2 | 1.8 | 103 | 3 | 2.9 | 0 | 0.0 | 93 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 19 | 3.0 | 14 | 2.5 | 50 | 2 | 4.0 | 50 | 2 | 4.0 | 49 | 1 | 2.0 | 0 | 0.0 | 45 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 0 | 0.0 | 0 | 0.0 | 14 | 0 | 0.0 | 14 | 0 | 0.0 | 14 | 1 | 7.1 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 624 | 100.0 | 555 | 100.0 | 547 | 25 | 4.6 | 547 | 9 | 1.6 | 501 | 5 | 1.0 | 0 | 0.0 | 475 | 1 | 0.2 | 1 | 0.2 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 12 | 6.6 | 12 | 7.6 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 53 | 29.3 | 48 | 30.4 | 12 | 1 | 8.3 | 12 | 0 | 0.0 | 11 | 0 | 0.0 | 0 | 0.0 | 10 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 38 | 21.0 | 33 | 20.9 | 47 | 5 | 10.6 | 48 | 1 | 2.1 | 43 | 0 | 0.0 | 0 | 0.0 | 43 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 30 | 16.6 | 23 | 14.6 | 32 | 1 | 3.1 | 32 | 2 | 6.3 | 31 | 0 | 0.0 | 0 | 0.0 | 29 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 35 | 19.3 | 30 | 19.0 | 22 | 1 | 4.5 | 22 | 0 | 0.0 | 21 | 0 | 0.0 | 0 | 0.0 | 20 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 9 | 5.0 | 9 | 5.7 | 28 | 1 | 3.6 | 28 | 0 | 0.0 | 28 | 2 | 7.1 | 1 | 3.6 | 24 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 2 | 1.1 | 1 | 0.6 | 9 | 0 | 0.0 | 9 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 2 | 1.1 | 2 | 1.3 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 181 | 100.0 | 158 | 100.0 | 152 | 9 | 5.9 | 153 | 3 | 2.0 | 146 | 2 | 1.4 | 1 | 0.7 | 137 | 0 | 0.0 | 0 | 0.0 |
| Others † | 1 | ~ | 1 | ~ | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 3.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Simms Mann Health & Wellness Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|-----------------------|--|-------|------------------------------------|-------|---------------|------------|------------|---------------|------------|------------|---------------|------------|------------|----------------------------|-----|---------------|------------|------------|-------------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | | HIV | | | |
| | | | | | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | n | % | Visits Tested | Positive n | Positive % | New HIV*** n % | |
| Total | 806 | 100.0 | 714 | 100.0 | 700 | 34 | 4.9 | 701 | 12 | 1.7 | 648 | 7 | 1.1 | 1 | 0.2 | 613 | 1 | 0.2 | 1 | 0.2 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 368 | 59.0 | 333 | 60.0 | 330 | 14 | 4.2 | 330 | 7 | 2.1 | 301 | 2 | 0.7 | 0 | 0.0 | 286 | 0 | 0.0 | 0 | 0.0 |
| African American | 86 | 13.8 | 71 | 12.8 | 69 | 4 | 5.8 | 69 | 1 | 1.4 | 61 | 0 | 0.0 | 0 | 0.0 | 55 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 79 | 12.7 | 72 | 13.0 | 69 | 4 | 5.8 | 69 | 1 | 1.4 | 67 | 3 | 4.5 | 0 | 0.0 | 63 | 0 | 0.0 | 0 | 0.0 |
| Asian | 54 | 8.7 | 50 | 9.0 | 50 | 2 | 4.0 | 50 | 0 | 0.0 | 46 | 0 | 0.0 | 0 | 0.0 | 45 | 0 | 0.0 | 0 | 0.0 |
| Native American | 2 | 0.3 | 2 | 0.4 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Other | 27 | 4.3 | 23 | 4.1 | 23 | 0 | 0.0 | 23 | 0 | 0.0 | 21 | 0 | 0.0 | 0 | 0.0 | 21 | 1 | 4.8 | 1 | 4.8 |
| Unknown | 8 | 1.3 | 4 | 0.7 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 624 | 100.0 | 555 | 100.0 | 547 | 25 | 4.6 | 547 | 9 | 1.6 | 501 | 5 | 1.0 | 0 | 0.0 | 475 | 1 | 0.2 | 1 | 0.2 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 86 | 47.5 | 75 | 47.5 | 74 | 3 | 4.1 | 74 | 1 | 1.4 | 69 | 0 | 0.0 | 0 | 0.0 | 66 | 0 | 0.0 | 0 | 0.0 |
| African American | 29 | 16.0 | 28 | 17.7 | 26 | 1 | 3.8 | 27 | 1 | 3.7 | 23 | 1 | 4.3 | 0 | 0.0 | 22 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 24 | 13.3 | 20 | 12.7 | 18 | 1 | 5.6 | 18 | 1 | 5.6 | 20 | 0 | 0.0 | 0 | 0.0 | 16 | 0 | 0.0 | 0 | 0.0 |
| Asian | 30 | 16.6 | 25 | 15.8 | 25 | 2 | 8.0 | 25 | 0 | 0.0 | 24 | 1 | 4.2 | 1 | 4.2 | 23 | 0 | 0.0 | 0 | 0.0 |
| Native American | 10 | 5.5 | 8 | 5.1 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Other | 2 | 1.1 | 2 | 1.3 | 7 | 1 | 14.3 | 7 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 181 | 100.0 | 158 | 100.0 | 152 | 9 | 5.9 | 153 | 3 | 2.0 | 146 | 2 | 1.4 | 1 | 0.7 | 137 | 0 | 0.0 | 0 | 0.0 |
| Others † | 1 | ~ | 1 | ~ | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 3.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Simms Mann Health and Wellness Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 3.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Simms Mann Health and Wellness Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

IV: Central Health Center

Table 4.1 Patient Visits and Tests, January - December 2010, Central Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-------------------|-------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 5,882 | 100.0 | 3,135 | 100.0 | 3,003 | 231 | 7.7 | 3,008 | 105 | 3.5 | 2,919 | 288 | 9.9 | 19 | 0.7 | 2,728 | 28 | 1.0 | 20 | 0.7 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 4,175 | 71.0 | 2,229 | 71.1 | 2,133 | 160 | 7.5 | 2,137 | 93 | 4.4 | 2,083 | 218 | 10.5 | 14 | 0.7 | 1,934 | 24 | 1.2 | 18 | 0.9 |
| Female | 1,678 | 28.5 | 897 | 28.6 | 861 | 70 | 8.1 | 862 | 12 | 1.4 | 827 | 66 | 8.0 | 5 | 0.6 | 786 | 4 | 0.5 | 2 | 0.3 |
| Transgender† | 29 | 0.5 | 9 | 0.3 | 9 | 1 | 11.1 | 9 | 0 | 0.0 | 9 | 4 | 44.4 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 991 | 16.8 | 522 | 16.7 | 498 | 51 | 10.2 | 501 | 62 | 12.4 | 480 | 85 | 17.7 | 9 | 1.9 | 413 | 19 | 4.6 | 13 | 3.1 |
| MSW | 2,956 | 50.3 | 1,618 | 51.6 | 1,571 | 106 | 6.7 | 1,572 | 29 | 1.8 | 1,520 | 114 | 7.5 | 4 | 0.3 | 1,460 | 5 | 0.3 | 5 | 0.3 |
| WSM | 1,485 | 25.2 | 803 | 25.6 | 779 | 67 | 8.6 | 780 | 9 | 1.2 | 743 | 51 | 6.9 | 5 | 0.7 | 715 | 3 | 0.4 | 2 | 0.3 |
| WSW/M | 82 | 1.4 | 46 | 1.5 | 45 | 3 | 6.7 | 45 | 3 | 6.7 | 41 | 2 | 4.9 | 0 | 0.0 | 440 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 368 | 6.3 | 146 | 4.7 | 110 | 4 | 3.6 | 110 | 2 | 1.8 | 135 | 36 | 26.7 | 1 | 0.7 | 100 | 1 | 1.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 6 | 0.1 | 2 | 0.1 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 273 | 4.6 | 157 | 5.0 | 153 | 30 | 19.6 | 153 | 4 | 2.6 | 148 | 9 | 6.1 | 0 | 0.0 | 145 | 1 | 0.7 | 1 | 0.7 |
| 20-24 | 1,016 | 17.3 | 570 | 18.2 | 559 | 67 | 12.0 | 560 | 15 | 2.7 | 528 | 21 | 4.0 | 2 | 0.4 | 521 | 5 | 1.0 | 5 | 1.0 |
| 25-29 | 1,124 | 19.1 | 623 | 19.9 | 616 | 58 | 9.4 | 616 | 31 | 5.0 | 583 | 29 | 5.0 | 2 | 0.3 | 568 | 7 | 1.2 | 4 | 0.7 |
| 30-34 | 913 | 15.5 | 475 | 15.2 | 455 | 25 | 5.5 | 457 | 21 | 4.6 | 441 | 38 | 8.6 | 4 | 0.9 | 416 | 2 | 0.5 | 2 | 0.5 |
| 35-44 | 1,279 | 21.7 | 682 | 21.8 | 665 | 35 | 5.3 | 666 | 21 | 3.2 | 633 | 75 | 11.8 | 3 | 0.5 | 583 | 9 | 1.5 | 6 | 1.0 |
| 45-54 | 813 | 13.8 | 420 | 13.4 | 396 | 11 | 2.8 | 396 | 11 | 2.8 | 390 | 67 | 17.2 | 5 | 1.3 | 344 | 1 | 0.3 | 1 | 0.3 |
| 55-64 | 341 | 5.8 | 159 | 5.1 | 126 | 3 | 2.4 | 127 | 1 | 0.8 | 148 | 38 | 25.7 | 2 | 1.4 | 118 | 3 | 2.5 | 1 | 0.8 |
| 65+ | 117 | 2.0 | 47 | 1.5 | 31 | 1 | 3.2 | 31 | 1 | 3.2 | 46 | 11 | 23.9 | 1 | 2.2 | 31 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 694 | 11.8 | 402 | 12.8 | 389 | 24 | 6.2 | 391 | 12 | 3.1 | 370 | 9 | 2.4 | 1 | 0.3 | 357 | 3 | 0.8 | 3 | 0.8 |
| African American | 1,068 | 18.2 | 637 | 20.3 | 610 | 61 | 10.0 | 610 | 44 | 7.2 | 565 | 46 | 8.1 | 3 | 0.5 | 512 | 4 | 0.8 | 3 | 0.6 |
| Hispanic | 3,690 | 62.7 | 1,867 | 59.6 | 1,778 | 132 | 7.4 | 1,781 | 43 | 2.4 | 1,769 | 222 | 12.5 | 15 | 0.8 | 1,653 | 21 | 1.3 | 14 | 0.8 |
| Asian | 120 | 2.0 | 64 | 2.0 | 62 | 3 | 4.8 | 62 | 0 | 0.0 | 62 | 5 | 8.1 | 0 | 0.0 | 58 | 0 | 0.0 | 0 | 0.0 |
| Native American | 11 | 0.2 | 6 | 0.2 | 6 | 1 | 16.7 | 6 | 0 | 0.0 | 5 | 1 | 20.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Other | 275 | 4.7 | 155 | 4.9 | 154 | 10 | 6.5 | 154 | 6 | 3.9 | 145 | 5 | 3.4 | 0 | 0.0 | 141 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 24 | 0.4 | 4 | 0.1 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 2,965 | 50.4 | 2,916 | 93.0 | 2,905 | 223 | 7.7 | 2,906 | 102 | 3.5 | 2,789 | 270 | 9.7 | 15 | 0.5 | 2,694 | 27 | 1.0 | 20 | 0.7 |
| Return Visit | 2,917 | 49.6 | 219 | 7.0 | 98 | 8 | 8.2 | 102 | 3 | 2.9 | 130 | 18 | 13.8 | 4 | 3.1 | 34 | 1 | 2.9 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 12 male-to-female (MTF) and 17 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 4.2 Patient Visits and Tests by Gender and Age, January - December 2010, Central Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|------------------|-----------------|-------------------------------|-------------|-------|-----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 5,882 | 100.0 | 3,135 | 100.0 | 3,003 | 231 | 7.7 | 3,008 | 105 | 3.5 | 2,919 | 288 | 9.9 | 19 | 0.7 | 2,728 | 28 | 1.0 | 20 | 0.7 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 4 | 0.1 | 1 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 148 | 3.5 | 84 | 3.8 | 81 | 12 | 14.8 | 81 | 3 | 3.7 | 81 | 7 | 8.6 | 0 | 0.0 | 78 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 675 | 16.1 | 377 | 16.9 | 371 | 44 | 11.9 | 372 | 14 | 3.8 | 353 | 17 | 4.8 | 1 | 0.3 | 349 | 5 | 1.4 | 5 | 1.4 |
| 25-29 | 837 | 20.0 | 466 | 20.9 | 462 | 44 | 9.5 | 462 | 30 | 6.5 | 436 | 24 | 5.5 | 2 | 0.5 | 426 | 7 | 1.6 | 4 | 0.9 |
| 30-34 | 666 | 15.9 | 351 | 15.7 | 336 | 22 | 6.5 | 338 | 17 | 5.0 | 329 | 28 | 8.5 | 3 | 0.9 | 310 | 2 | 0.6 | 2 | 0.6 |
| 35-44 | 929 | 22.2 | 494 | 22.1 | 481 | 26 | 5.4 | 482 | 17 | 3.5 | 463 | 55 | 11.9 | 3 | 0.6 | 422 | 8 | 1.9 | 6 | 1.4 |
| 45-54 | 592 | 14.2 | 306 | 13.7 | 286 | 10 | 3.5 | 286 | 10 | 3.5 | 279 | 53 | 19.0 | 4 | 1.4 | 238 | 1 | 0.4 | 1 | 0.4 |
| 55-65 | 254 | 6.1 | 121 | 5.4 | 98 | 1 | 1.0 | 98 | 1 | 1.0 | 113 | 29 | 25.7 | 0 | 0.0 | 93 | 1 | 1.1 | 0 | 0.0 |
| 65+ | 78 | 1.9 | 32 | 1.4 | 20 | 1 | 5.0 | 20 | 1 | 5.0 | 31 | 7 | 22.6 | 1 | 3.2 | 20 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 4,183 | 100.0 | 2,232 | 100.0 | 2,136 | 160 | 7.5 | 2,140 | 93 | 4.3 | 2,086 | 220 | 10.5 | 14 | 0.7 | 1,937 | 24 | 1.2 | 18 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 2 | 0.1 | 1 | 0.1 | 1 | 1 | 100.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 122 | 7.3 | 72 | 8.0 | 71 | 18 | 25.4 | 71 | 1 | 1.4 | 66 | 2 | 3.0 | 0 | 0.0 | 66 | 1 | 1.5 | 1 | 1.5 |
| 20-24 | 337 | 20.1 | 192 | 21.4 | 187 | 23 | 12.3 | 187 | 1 | 0.5 | 174 | 3 | 1.7 | 1 | 0.6 | 171 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 285 | 17.0 | 156 | 17.4 | 153 | 14 | 9.2 | 153 | 1 | 0.7 | 146 | 5 | 3.4 | 0 | 0.0 | 141 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 243 | 14.5 | 122 | 13.6 | 117 | 3 | 2.6 | 117 | 4 | 3.4 | 112 | 10 | 8.9 | 1 | 0.9 | 106 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 343 | 20.4 | 186 | 20.8 | 182 | 8 | 4.4 | 182 | 4 | 2.2 | 168 | 19 | 11.3 | 0 | 0.0 | 159 | 1 | 0.6 | 0 | 0.0 |
| 45-54 | 221 | 13.2 | 114 | 12.7 | 110 | 1 | 0.9 | 110 | 1 | 0.9 | 111 | 14 | 12.6 | 1 | 0.9 | 106 | 0 | 0.0 | 0 | 0.0 |
| 55-65 | 87 | 5.2 | 38 | 4.2 | 28 | 2 | 7.1 | 29 | 0 | 0.0 | 35 | 9 | 25.7 | 2 | 5.7 | 25 | 2 | 8.0 | 1 | 4.0 |
| 65+ | 39 | 2.3 | 15 | 1.7 | 11 | 0 | 0.0 | 11 | 0 | 0.0 | 15 | 4 | 26.7 | 0 | 0.0 | 11 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,679 | 100.0 | 896 | 100.0 | 860 | 70 | 8.1 | 861 | 12 | 1.4 | 828 | 66 | 8.0 | 5 | 0.6 | 786 | 4 | 0.5 | 2 | 0.3 |
| Others † | 20 | ~ | 7 | ~ | 7 | 1.0 | 14.3 | 7 | 0.0 | 0.0 | 5 | 2.0 | 40.0 | 0 | 0.0 | 5 | 0.0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 4.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Central Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|-----------------------|--|-------|--------------------------|-------|---------------|------------|------------|---------------|------------|------------|---------------|------------|------------|----------------------------|-----|---------------|------------|------------|--------------|-------|
| | Total STD Visits | | No. Visits with STD Test | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | n | % | Visits Tested | Positive n | Positive % | New HIV*** n | *** % |
| n | % | n | % | | | | | | | | | | | | | | | | | |
| Total | 5,882 | 100.0 | 3,135 | 100.0 | 3,003 | 231 | 7.7 | 3,008 | 105 | 3.5 | 2,919 | 288 | 9.9 | 19 | 0.7 | 2,728 | 28 | 1.0 | 20 | 0.7 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 515 | 12.3 | 296 | 13.3 | 286 | 16 | 5.6 | 287 | 10 | 3.5 | 275 | 8 | 2.9 | 1 | 0.4 | 264 | 3 | 1.1 | 3 | 1.1 |
| African American | 775 | 18.5 | 458 | 20.5 | 433 | 46 | 10.6 | 433 | 41 | 9.5 | 411 | 29 | 7.1 | 2 | 0.5 | 365 | 3 | 0.8 | 2 | 0.5 |
| Hispanic | 2,602 | 62.2 | 1,325 | 59.4 | 1,267 | 89 | 7.0 | 1,270 | 39 | 3.1 | 1,258 | 173 | 13.8 | 11 | 0.9 | 1,172 | 18 | 1.5 | 13 | 1.1 |
| Asian | 72 | 1.7 | 38 | 1.7 | 36 | 3 | 8.3 | 36 | 0 | 0.0 | 37 | 4 | 10.8 | 0 | 0.0 | 34 | 0 | 0.0 | 0 | 0.0 |
| Native American | 7 | 0.2 | 4 | 0.2 | 4 | 1 | 25.0 | 4 | 0 | 0.0 | 3 | 1 | 33.3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Other | 192 | 4.6 | 107 | 4.8 | 106 | 5 | 4.7 | 106 | 3 | 2.8 | 99 | 5 | 5.1 | 0 | 0.0 | 96 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 20 | 0.5 | 4 | 0.2 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 4,183 | 100.0 | 2,232 | 100.0 | 2,136 | 160 | 7.5 | 2,140 | 93 | 4.3 | 2,086 | 220 | 10.5 | 14 | 0.7 | 1,937 | 24 | 1.2 | 18 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 179 | 10.7 | 106 | 11.8 | 103 | 8 | 7.8 | 104 | 2 | 1.9 | 95 | 1 | 1.1 | 0 | 0.0 | 93 | 0 | 0.0 | 0 | 0.0 |
| African American | 293 | 17.5 | 179 | 20.0 | 177 | 15 | 8.5 | 177 | 3 | 1.7 | 154 | 17 | 11.0 | 1 | 0.6 | 147 | 1 | 0.7 | 1 | 0.7 |
| Hispanic | 1,070 | 63.7 | 537 | 59.9 | 506 | 42 | 8.3 | 506 | 4 | 0.8 | 506 | 47 | 9.3 | 4 | 0.8 | 476 | 3 | 0.6 | 3 | 0.6 |
| Asian | 48 | 2.9 | 26 | 2.9 | 26 | 0 | 0.0 | 26 | 0 | 0.0 | 25 | 1 | 4.0 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 |
| Native American | 4 | 0.2 | 2 | 0.2 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 81 | 4.8 | 46 | 5.1 | 46 | 5 | 10.9 | 46 | 3 | 6.5 | 46 | 0 | 0.0 | 0 | 0.0 | 45 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 4 | 0.2 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,679 | 100.0 | 896 | 100.0 | 860 | 70 | 8.1 | 861 | 12 | 1.4 | 828 | 66 | 8.0 | 5 | 0.6 | 786 | 4 | 0.5 | 2 | 0.3 |
| Others † | 20 | ~ | 7 | ~ | 7 | 1 | 14.3 | 7 | 0 | 0.0 | 5 | 2 | 40.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 4.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Central Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 4.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Central Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

V: Hollywood-Wilshire Health Center

Table 5.1 Patient Visits and Tests, January - December 2010, Hollywood-Wilshire Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | | HIV Testing | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-----|-------|-------------|-----|-------------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | New HIV*** n % | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | | | | | | |
| Total | 6,205 | 100.0 | 3,747 | 100.0 | 3,529 | 286 | 8.1 | 3,536 | 144 | 4.1 | 3,112 | 215 | 6.9 | 22 | 0.7 | 2,758 | 26 | 0.9 | 20 | 0.7 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 4,684 | 75.5 | 2,796 | 74.6 | 2,642 | 222 | 8.4 | 2,647 | 137 | 5.2 | 2,334 | 178 | 7.6 | 22 | 0.9 | 2,055 | 24 | 1.2 | 18 | 0.9 |
| Female | 1,480 | 23.9 | 934 | 24.9 | 870 | 64 | 7.4 | 872 | 5 | 0.6 | 761 | 34 | 4.5 | 0 | 0.0 | 688 | 2 | 0.3 | 2 | 0.3 |
| Transgender† | 41 | 0.7 | 17 | 0.5 | 17 | 0 | 0.0 | 17 | 2 | 11.8 | 17 | 3 | 17.6 | 0 | 0.0 | 15 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 1,240 | 20.0 | 744 | 19.9 | 688 | 60 | 8.7 | 692 | 93 | 13.4 | 618 | 97 | 15.7 | 16 | 2.6 | 463 | 20 | 4.3 | 15 | 3.2 |
| MSW | 3,223 | 51.9 | 1,968 | 52.5 | 1,887 | 157 | 8.3 | 1,887 | 40 | 2.1 | 1,646 | 67 | 4.1 | 6 | 0.4 | 1538 | 3 | 0.2 | 3 | 0.2 |
| WSM | 1,272 | 20.5 | 806 | 21.5 | 757 | 56 | 7.4 | 758 | 4 | 0.5 | 661 | 21 | 3.2 | 0 | 0.0 | 604 | 2 | 0.3 | 2 | 0.3 |
| WSW/M | 110 | 1.8 | 77 | 2.1 | 73 | 6 | 8.2 | 74 | 1 | 1.4 | 58 | 3 | 5.2 | 0 | 0.0 | 51 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 360 | 5.8 | 152 | 4.1 | 124 | 7 | 5.6 | 125 | 6 | 4.8 | 129 | 27 | 20.9 | 0 | 0.0 | 102 | 1 | 1.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 3 | 0.0 | 1 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 278 | 4.5 | 167 | 4.5 | 158 | 31 | 19.6 | 158 | 8 | 5.1 | 136 | 1 | 0.7 | 0 | 0.0 | 129 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 1,189 | 19.2 | 759 | 20.3 | 739 | 86 | 11.6 | 741 | 28 | 3.8 | 613 | 28 | 4.6 | 2 | 0.3 | 577 | 3 | 0.5 | 2 | 0.3 |
| 25-29 | 1,412 | 22.8 | 883 | 23.6 | 855 | 59 | 6.9 | 857 | 33 | 3.9 | 758 | 29 | 3.8 | 4 | 0.5 | 716 | 4 | 0.6 | 4 | 0.6 |
| 30-34 | 1,079 | 17.4 | 668 | 17.8 | 644 | 43 | 6.7 | 647 | 28 | 4.3 | 563 | 29 | 5.2 | 2 | 0.4 | 504 | 7 | 1.4 | 6 | 1.2 |
| 35-44 | 1,284 | 20.7 | 756 | 20.2 | 702 | 49 | 7.0 | 702 | 27 | 3.8 | 617 | 44 | 7.1 | 9 | 1.5 | 527 | 9 | 1.7 | 6 | 1.1 |
| 45-54 | 682 | 11.0 | 374 | 10.0 | 326 | 16 | 4.9 | 326 | 18 | 5.5 | 306 | 52 | 17.0 | 3 | 1.0 | 223 | 3 | 1.3 | 2 | 0.9 |
| 55-64 | 233 | 3.8 | 121 | 3.2 | 92 | 2 | 2.2 | 92 | 2 | 2.2 | 101 | 23 | 22.8 | 1 | 1.0 | 73 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 45 | 0.7 | 18 | 0.5 | 12 | 0 | 0.0 | 12 | 0 | 0.0 | 17 | 9 | 52.9 | 1 | 5.9 | 8 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 2,028 | 32.7 | 1,302 | 34.7 | 1,264 | 81 | 6.4 | 1,264 | 38 | 3.0 | 1,075 | 31 | 2.9 | 4 | 0.4 | 996 | 6 | 0.6 | 4 | 0.4 |
| African American | 1,224 | 19.7 | 771 | 20.6 | 748 | 90 | 12.0 | 748 | 48 | 6.4 | 603 | 29 | 4.8 | 3 | 0.5 | 531 | 5 | 0.9 | 4 | 0.8 |
| Hispanic | 2,080 | 33.5 | 1,144 | 30.5 | 1,018 | 81 | 8.0 | 1,023 | 45 | 4.4 | 965 | 136 | 14.1 | 13 | 1.3 | 806 | 13 | 1.6 | 10 | 1.2 |
| Asian | 528 | 8.5 | 317 | 8.5 | 293 | 23 | 7.8 | 294 | 8 | 2.7 | 284 | 11 | 3.9 | 2 | 0.7 | 249 | 1 | 0.4 | 1 | 0.4 |
| Native American | 16 | 0.3 | 11 | 0.3 | 11 | 2 | 18.2 | 11 | 1 | 9.1 | 11 | 0 | 0.0 | 0 | 0.0 | 11 | 0 | 0.0 | 0 | 0.0 |
| Other | 319 | 5.1 | 194 | 5.2 | 188 | 9 | 4.8 | 189 | 4 | 2.1 | 166 | 7 | 4.2 | 0 | 0.0 | 158 | 1 | 0.6 | 1 | 0.6 |
| Unknown | 10 | 0.2 | 8 | 0.2 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 8 | 1 | 12.5 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 3,567 | 57.5 | 3,494 | 93.2 | 3,368 | 275 | 8.2 | 3,370 | 128 | 3.8 | 2,968 | 174 | 5.9 | 18 | 0.6 | 2,680 | 22 | 0.8 | 19 | 0.7 |
| Return Visit | 2,638 | 42.5 | 253 | 6.8 | 161 | 11 | 6.8 | 166 | 16 | 9.6 | 144 | 41 | 28.5 | 4 | 2.8 | 78 | 4 | 5.1 | 1 | 1.3 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 14 male-to-female (MTF) and 27 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 5.2 Patient Visits and Tests by Gender and Age, January - December 2010, Hollywood-Wilshire Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------|-------|----------------------|---------------|--------------|---------------|--------------|---------------|--------------|-----|-------|----------------------------|--------------|-------------------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | n | % | with STD Test n % | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n | % | Visits Tested | Positive n % | New HIV*** n % | | | | |
| Total | 6,205 | 100.0 | 3,747 | 100.0 | 3,529 | 286 | 8.1 | 3,536 | 144 | 4.1 | 3,112 | 215 | 6.9 | 22 | 0.7 | 2,758 | 26 | 0.9 | 20 | 0.7 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 3 | 0.1 | 1 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 153 | 3.3 | 91 | 3.3 | 87 | 13 | 14.9 | 87 | 5 | 5.7 | 78 | 1 | 1.3 | 0 | 0.0 | 73 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 826 | 17.6 | 520 | 18.6 | 506 | 59 | 11.7 | 507 | 27 | 5.3 | 437 | 22 | 5.0 | 2 | 0.5 | 415 | 3 | 0.7 | 2 | 0.5 |
| 25-29 | 1,076 | 23.0 | 661 | 23.6 | 641 | 52 | 8.1 | 642 | 32 | 5.0 | 577 | 27 | 4.7 | 4 | 0.7 | 544 | 4 | 0.7 | 4 | 0.7 |
| 30-34 | 869 | 18.5 | 529 | 18.9 | 513 | 41 | 8.0 | 516 | 26 | 5.0 | 438 | 25 | 5.7 | 2 | 0.5 | 390 | 6 | 1.5 | 5 | 1.3 |
| 35-44 | 1,043 | 22.3 | 614 | 22.0 | 577 | 40 | 6.9 | 577 | 27 | 4.7 | 494 | 34 | 6.9 | 9 | 1.8 | 419 | 8 | 1.9 | 5 | 1.2 |
| 45-54 | 510 | 10.9 | 276 | 9.9 | 240 | 15 | 6.3 | 240 | 18 | 7.5 | 225 | 44 | 19.6 | 3 | 1.3 | 157 | 3 | 1.9 | 2 | 1.3 |
| 55-64 | 173 | 3.7 | 89 | 3.2 | 65 | 2 | 3.1 | 65 | 2 | 3.1 | 70 | 20 | 28.6 | 1 | 1.4 | 47 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 33 | 0.7 | 14 | 0.5 | 11 | 0 | 0.0 | 11 | 0 | 0.0 | 13 | 5 | 38.5 | 1 | 7.7 | 8 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 4,686 | 100.0 | 2,795 | 100.0 | 2,641 | 222 | 8.4 | 2,646 | 137 | 5.2 | 2,333 | 178 | 7.6 | 22 | 0.9 | 2,054 | 24 | 1.2 | 18 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 124 | 8.3 | 75 | 8.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 355 | 23.8 | 234 | 25.1 | 70 | 18 | 25.7 | 70 | 3 | 4.3 | 57 | 0 | 0.0 | 0 | 0.0 | 55 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 332 | 22.3 | 220 | 23.6 | 228 | 27 | 11.8 | 229 | 1 | 0.4 | 171 | 4 | 2.3 | 0 | 0.0 | 157 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 199 | 13.4 | 131 | 14.0 | 212 | 7 | 3.3 | 213 | 1 | 0.5 | 179 | 2 | 1.1 | 0 | 0.0 | 171 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 240 | 16.1 | 142 | 15.2 | 124 | 2 | 1.6 | 124 | 1 | 0.8 | 118 | 4 | 3.4 | 0 | 0.0 | 107 | 1 | 0.9 | 1 | 0.9 |
| 35-44 | 168 | 11.3 | 96 | 10.3 | 125 | 9 | 7.2 | 125 | 0 | 0.0 | 123 | 10 | 8.1 | 0 | 0.0 | 108 | 1 | 0.9 | 1 | 0.9 |
| 45-54 | 60 | 4.0 | 32 | 3.4 | 84 | 1 | 1.2 | 84 | 0 | 0.0 | 79 | 7 | 8.9 | 0 | 0.0 | 64 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 12 | 0.8 | 4 | 0.4 | 27 | 0 | 0.0 | 27 | 0 | 0.0 | 31 | 3 | 9.7 | 0 | 0.0 | 26 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 4 | 4 | 100.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,490 | 100.0 | 934 | 100.0 | 871 | 64 | 7.3 | 873 | 6 | 0.7 | 762 | 34 | 4.5 | 0 | 0.0 | 688 | 2 | 0.3 | 2 | 0.3 |
| Others † | 29 | ~ | 18 | ~ | 17 | 0.0 | 0.0 | 17 | 1.0 | 5.9 | 17 | 3.0 | 17.6 | 0 | 0.0 | 16 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 5.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Hollywood-Wilshire Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|------------------|-----------------|-------------------------------|-----|-------------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 6,205 | 100.0 | 3,747 | 100.0 | 3,529 | 286 | 8.1 | 3,536 | 144 | 4.1 | 3,112 | 215 | 6.9 | 22 | 0.7 | 2,758 | 26 | 0.9 | 20 | 0.7 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 1,630 | 34.8 | 1,021 | 36.5 | 991 | 68 | 6.9 | 991 | 36 | 3.6 | 847 | 30 | 3.5 | 4 | 0.5 | 775 | 6 | 0.8 | 4 | 0.5 |
| African American | 927 | 19.8 | 578 | 20.7 | 566 | 70 | 12.4 | 566 | 45 | 8.0 | 453 | 25 | 5.5 | 3 | 0.7 | 400 | 5 | 1.3 | 4 | 1.0 |
| Hispanic | 1,555 | 33.2 | 852 | 30.5 | 755 | 55 | 7.3 | 760 | 43 | 5.7 | 724 | 111 | 15.3 | 13 | 1.8 | 597 | 12 | 2.0 | 9 | 1.5 |
| Asian | 338 | 7.2 | 199 | 7.1 | 187 | 21 | 11.2 | 187 | 8 | 4.3 | 179 | 6 | 3.4 | 2 | 1.1 | 156 | 0 | 0.0 | 0 | 0.0 |
| Native American | 12 | 0.3 | 8 | 0.3 | 8 | 2 | 25.0 | 8 | 1 | 12.5 | 8 | 0 | 0.0 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| Other | 214 | 4.6 | 129 | 4.6 | 127 | 6 | 4.7 | 127 | 4 | 3.1 | 114 | 5 | 4.4 | 0 | 0.0 | 111 | 1 | 0.9 | 1 | 0.9 |
| Unknown | 10 | 0.2 | 8 | 0.3 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 8 | 1 | 12.5 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 4,686 | 100.0 | 2,795 | 100.0 | 2,641 | 222 | 8.4 | 2,646 | 137 | 5.2 | 2,333 | 178 | 7.6 | 22 | 0.9 | 2,054 | 24 | 1.2 | 18 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 397 | 26.6 | 280 | 30.0 | 272 | 13 | 4.8 | 272 | 2 | 0.7 | 227 | 1 | 0.4 | 0 | 0.0 | 220 | 0 | 0.0 | 0 | 0.0 |
| African American | 294 | 19.7 | 191 | 20.4 | 180 | 20 | 11.1 | 180 | 3 | 1.7 | 148 | 4 | 2.7 | 0 | 0.0 | 130 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 506 | 34.0 | 281 | 30.1 | 252 | 26 | 10.3 | 252 | 1 | 0.4 | 230 | 22 | 9.6 | 0 | 0.0 | 199 | 1 | 0.5 | 1 | 0.5 |
| Asian | 188 | 12.6 | 117 | 12.5 | 105 | 2 | 1.9 | 106 | 0 | 0.0 | 104 | 5 | 4.8 | 0 | 0.0 | 92 | 1 | 1.1 | 1 | 1.1 |
| Native American | 4 | 0.3 | 3 | 0.3 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Other | 101 | 6.8 | 62 | 6.6 | 59 | 3 | 5.1 | 60 | 0 | 0.0 | 50 | 2 | 4.0 | 0 | 0.0 | 44 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,490 | 100.0 | 934 | 100.0 | 871 | 64 | 7.3 | 873 | 6 | 0.7 | 762 | 34 | 4.5 | 0 | 0.0 | 688 | 2 | 0.3 | 2 | 0.3 |
| Others † | 29 | ~ | 18 | ~ | 17 | 0 | 0.0 | 17 | 1 | 5.9 | 17 | 3 | 17.6 | 0 | 0.0 | 16 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 5.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Hollywood-Wilshire Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 5.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Hollywood-Wilshire Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Sexually Transmitted Disease Clinic Morbidity Report, 2010

VI: Monrovia Health Center

Table 6.1 Patient Visits and Tests, January - December 2010, Monrovia Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-----|-------------|---|-----|-------------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | New HIV*** n % | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | | | | | | |
| Total | 1,222 | 100.0 | 913 | 100.0 | 861 | 98 | 11.4 | 862 | 28 | 3.2 | 838 | 69 | 8.2 | 6 | 0.7 | 757 | 5 | 0.7 | 4 | 0.5 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 813 | 66.5 | 609 | 66.7 | 585 | 74 | 12.6 | 587 | 24 | 4.1 | 569 | 42 | 7.4 | 6 | 1.1 | 520 | 5 | 1.0 | 4 | 0.8 |
| Female | 406 | 33.2 | 302 | 33.1 | 274 | 24 | 8.8 | 273 | 4 | 1.5 | 267 | 27 | 10.1 | 0 | 0.0 | 235 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | 3 | 0.2 | 2 | 0.2 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 172 | 14.1 | 129 | 14.1 | 121 | 25 | 20.7 | 123 | 16 | 13.0 | 120 | 15 | 12.5 | 6 | 5.0 | 103 | 5 | 4.9 | 4 | 3.9 |
| MSW | 592 | 48.4 | 445 | 48.7 | 432 | 48 | 11.1 | 432 | 7 | 1.6 | 417 | 24 | 5.8 | 0 | 0.0 | 391 | 0 | 0.0 | 0 | 0.0 |
| WSM | 359 | 29.4 | 275 | 30.1 | 253 | 23 | 9.1 | 252 | 4 | 1.6 | 245 | 25 | 10.2 | 0 | 0.0 | 221 | 0 | 0.0 | 0 | 0.0 |
| WSW/M | 9 | 0.7 | 8 | 0.9 | 8 | 0 | 0.0 | 8 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 90 | 7.4 | 56 | 6.1 | 47 | 2 | 4.3 | 47 | 1 | 2.1 | 50 | 5 | 10.0 | 0 | 0.0 | 36 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 3 | 0.2 | 1 | 0.1 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 71 | 5.8 | 63 | 6.9 | 59 | 14 | 23.7 | 59 | 2 | 3.4 | 57 | 2 | 3.5 | 0 | 0.0 | 52 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 277 | 22.7 | 213 | 23.3 | 212 | 38 | 17.9 | 211 | 8 | 3.8 | 194 | 12 | 6.2 | 3 | 1.5 | 185 | 1 | 0.5 | 1 | 0.5 |
| 25-29 | 248 | 20.3 | 189 | 20.7 | 177 | 18 | 10.2 | 179 | 6 | 3.4 | 171 | 9 | 5.3 | 1 | 0.6 | 162 | 1 | 0.6 | 1 | 0.6 |
| 30-34 | 135 | 11.0 | 103 | 11.3 | 97 | 8 | 8.2 | 97 | 7 | 7.2 | 91 | 4 | 4.4 | 0 | 0.0 | 85 | 2 | 2.4 | 2 | 2.4 |
| 35-44 | 252 | 20.6 | 188 | 20.6 | 180 | 16 | 8.9 | 180 | 4 | 2.2 | 173 | 21 | 12.1 | 1 | 0.6 | 154 | 1 | 0.6 | 0 | 0.0 |
| 45-54 | 140 | 11.5 | 91 | 10.0 | 83 | 3 | 3.6 | 83 | 1 | 1.2 | 87 | 11 | 12.6 | 1 | 1.1 | 75 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 56 | 4.6 | 43 | 4.7 | 37 | 1 | 2.7 | 37 | 0 | 0.0 | 43 | 5 | 11.6 | 0 | 0.0 | 32 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 40 | 3.3 | 22 | 2.4 | 15 | 0 | 0.0 | 15 | 0 | 0.0 | 21 | 5 | 23.8 | 0 | 0.0 | 11 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 186 | 15.2 | 144 | 15.8 | 139 | 13 | 9.4 | 139 | 5 | 3.6 | 136 | 4 | 2.9 | 1 | 0.7 | 125 | 0 | 0.0 | 0 | 0.0 |
| African American | 117 | 9.6 | 96 | 10.5 | 95 | 15 | 15.8 | 95 | 4 | 4.2 | 88 | 2 | 2.3 | 0 | 0.0 | 84 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 588 | 48.1 | 439 | 48.1 | 416 | 49 | 11.8 | 416 | 16 | 3.8 | 398 | 27 | 6.8 | 4 | 1.0 | 364 | 3 | 0.8 | 2 | 0.5 |
| Asian | 253 | 20.7 | 177 | 19.4 | 155 | 16 | 10.3 | 156 | 2 | 1.3 | 162 | 32 | 19.8 | 0 | 0.0 | 132 | 1 | 0.8 | 1 | 0.8 |
| Native American | 3 | 0.2 | 2 | 0.2 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Other | 68 | 5.6 | 52 | 5.7 | 51 | 4 | 7.8 | 51 | 1 | 2.0 | 49 | 3 | 6.1 | 1 | 2.0 | 48 | 1 | 2.1 | 1 | 2.1 |
| Unknown | 7 | 0.6 | 3 | 0.3 | 3 | 1 | 33.3 | 3 | 0 | 0.0 | 3 | 1 | 33.3 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 785 | 64.2 | 759 | 83.1 | 754 | 83 | 11.0 | 754 | 19 | 2.5 | 716 | 46 | 6.4 | 4 | 0.6 | 682 | 2 | 0.3 | 2 | 0.3 |
| Return Visit | 437 | 35.8 | 154 | 16.9 | 107 | 15 | 14.0 | 108 | 9 | 8.3 | 122 | 23 | 18.9 | 2 | 1.6 | 75 | 3 | 4.0 | 2 | 2.7 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 3 male-to-female (MTF)

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 6.2 Patient Visits and Tests by Gender and Age, January - December 2010, Monrovia Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------------------|-----------------|------|-------------------------------|------------------|-----------------|-----|-----|------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV^ | | | New HIV*** | |
| | | | | | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | n % | | | |
| Total | 1,222 | 100.0 | 913 | 100.0 | 861 | 98 | 11.4 | 862 | 28 | 3.2 | 838 | 69 | 8.2 | 6 | 0.7 | 757 | 5 | 0.7 | 4 | 0.5 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 1 | 0.1 | 28 | 4.6 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 34 | 4.2 | 140 | 23.0 | 28 | 7 | 25.0 | 28 | 1 | 3.6 | 25 | 0 | 0.0 | 0 | 0.0 | 25 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 186 | 22.9 | 134 | 22.0 | 140 | 29 | 20.7 | 140 | 8 | 5.7 | 131 | 12 | 9.2 | 3 | 2.3 | 125 | 1 | 0.8 | 1 | 0.8 |
| 25-29 | 173 | 21.3 | 70 | 11.5 | 127 | 13 | 10.2 | 129 | 4 | 3.1 | 125 | 4 | 3.2 | 1 | 0.8 | 119 | 1 | 0.8 | 1 | 0.8 |
| 30-34 | 89 | 10.9 | 132 | 21.7 | 69 | 7 | 10.1 | 69 | 6 | 8.7 | 63 | 2 | 3.2 | 0 | 0.0 | 61 | 2 | 3.3 | 2 | 3.3 |
| 35-44 | 174 | 21.4 | 60 | 9.9 | 127 | 14 | 11.0 | 127 | 4 | 3.1 | 123 | 10 | 8.1 | 1 | 0.8 | 108 | 1 | 0.9 | 0 | 0.0 |
| 45-54 | 90 | 11.1 | 33 | 5.4 | 57 | 3 | 5.3 | 57 | 1 | 1.8 | 57 | 7 | 12.3 | 1 | 1.8 | 52 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 42 | 5.2 | 12 | 2.0 | 29 | 1 | 3.4 | 29 | 0 | 0.0 | 33 | 4 | 12.1 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 24 | 3.0 | 0 | 0.0 | 8 | 0 | 0.0 | 8 | 0 | 0.0 | 12 | 3 | 25.0 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 813 | 100.0 | 609 | 100.0 | 585 | 74 | 12.6 | 587 | 24 | 4.1 | 569 | 42 | 7.4 | 6 | 1.1 | 520 | 5 | 1.0 | 4 | 0.8 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 2 | 0.5 | 1 | 0.3 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 37 | 9.0 | 35 | 11.5 | 31 | 7 | 22.6 | 31 | 1 | 3.2 | 32 | 2 | 6.3 | 0 | 0.0 | 27 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 91 | 22.2 | 73 | 24.0 | 72 | 9 | 12.5 | 71 | 0 | 0.0 | 63 | 0 | 0.0 | 0 | 0.0 | 60 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 75 | 18.3 | 55 | 18.1 | 50 | 5 | 10.0 | 50 | 2 | 4.0 | 46 | 5 | 10.9 | 0 | 0.0 | 43 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 46 | 11.2 | 33 | 10.9 | 28 | 1 | 3.6 | 28 | 1 | 3.6 | 28 | 2 | 7.1 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 78 | 19.1 | 56 | 18.4 | 53 | 2 | 3.8 | 53 | 0 | 0.0 | 50 | 11 | 22.0 | 0 | 0.0 | 46 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 50 | 12.2 | 31 | 10.2 | 26 | 0 | 0.0 | 26 | 0 | 0.0 | 30 | 4 | 13.3 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 14 | 3.4 | 10 | 3.3 | 8 | 0 | 0.0 | 8 | 0 | 0.0 | 10 | 1 | 10.0 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 16 | 3.9 | 10 | 3.3 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 9 | 2 | 22.2 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 409 | 100.0 | 304 | 100.0 | 276 | 24 | 8.7 | 275 | 4 | 1.5 | 269 | 27 | 10.0 | 0 | 0.0 | 237 | 0 | 0.0 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 6.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Monrovia Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|------------------|-----------------|-------------------------------|-------------|-----|---|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 1,222 | 100.0 | 913 | 100.0 | 861 | 98 | 11.4 | 862 | 28 | 3.2 | 838 | 69 | 8.2 | 6 | 0.7 | 757 | 5 | 0.7 | 4 | 0.5 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 141 | 17.3 | 106 | 17.4 | 103 | 11 | 10.7 | 103 | 5 | 4.9 | 99 | 2 | 2.0 | 1 | 1.0 | 91 | 0 | 0.0 | 0 | 0.0 |
| African American | 82 | 10.1 | 69 | 11.3 | 69 | 13 | 18.8 | 69 | 4 | 5.8 | 64 | 1 | 1.6 | 0 | 0.0 | 61 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 401 | 49.3 | 297 | 48.8 | 284 | 37 | 13.0 | 285 | 12 | 4.2 | 279 | 22 | 7.9 | 4 | 1.4 | 256 | 3 | 1.2 | 2 | 0.8 |
| Asian | 145 | 17.8 | 102 | 16.7 | 94 | 11 | 11.7 | 95 | 2 | 2.1 | 95 | 14 | 14.7 | 0 | 0.0 | 81 | 1 | 1.2 | 1 | 1.2 |
| Native American | 40 | 4.9 | 33 | 5.4 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Other | 4 | 0.5 | 2 | 0.3 | 33 | 1 | 3.0 | 33 | 1 | 3.0 | 30 | 2 | 6.7 | 1 | 3.3 | 30 | 1 | 3.3 | 1 | 3.3 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 1 | 50.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 813 | 100.0 | 609 | 100.0 | 585 | 74 | 12.6 | 587 | 24 | 4.1 | 569 | 42 | 7.4 | 6 | 1.1 | 520 | 5 | 1.0 | 4 | 0.8 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 45 | 11.0 | 38 | 12.5 | 36 | 2 | 5.6 | 36 | 0 | 0.0 | 37 | 2 | 5.4 | 0 | 0.0 | 34 | 0 | 0.0 | 0 | 0.0 |
| African American | 35 | 8.6 | 27 | 8.9 | 26 | 2 | 7.7 | 26 | 0 | 0.0 | 24 | 1 | 4.2 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 187 | 45.7 | 142 | 46.7 | 132 | 12 | 9.1 | 131 | 4 | 3.1 | 119 | 5 | 4.2 | 0 | 0.0 | 108 | 0 | 0.0 | 0 | 0.0 |
| Asian | 108 | 26.4 | 75 | 24.7 | 61 | 5 | 8.2 | 61 | 0 | 0.0 | 67 | 18 | 26.9 | 0 | 0.0 | 51 | 0 | 0.0 | 0 | 0.0 |
| Native American | 3 | 0.7 | 2 | 0.7 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Other | 28 | 6.8 | 19 | 6.3 | 18 | 3 | 16.7 | 18 | 0 | 0.0 | 19 | 1 | 5.3 | 0 | 0.0 | 18 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 3 | 0.7 | 1 | 0.3 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 409 | 100.0 | 304 | 100.0 | 276 | 24 | 8.7 | 275 | 4 | 1.5 | 269 | 27 | 10.0 | 0 | 0.0 | 237 | 0 | 0.0 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 6.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Monrovia Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 6.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Monrovia Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

VII: North Hollywood Health Center

Table 7.1 Patient Visits and Tests, January - December 2010, North Hollywood Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|------------------|-----------------|-------------------------------|-----|-------------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 5,596 | 100.0 | 3,302 | 100.0 | 3,099 | 235 | 7.6 | 3,105 | 109 | 3.5 | 3,014 | 157 | 5.2 | 17 | 0.6 | 2,720 | 25 | 0.9 | 21 | 0.8 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 3,976 | 71.1 | 2,342 | 70.9 | 2,231 | 175 | 7.8 | 2,234 | 100 | 4.5 | 2,157 | 117 | 5.4 | 15 | 0.7 | 1,947 | 23 | 1.2 | 20 | 1.0 |
| Female | 1,604 | 28.7 | 951 | 28.8 | 860 | 60 | 7.0 | 862 | 6 | 0.7 | 849 | 39 | 4.6 | 2 | 0.2 | 768 | 2 | 0.3 | 1 | 0.1 |
| Transgender† | 16 | 0.3 | 9 | 0.3 | 8 | 0 | 0.0 | 9 | 3 | 33.3 | 8 | 1 | 12.5 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 854 | 15.3 | 529 | 16.0 | 493 | 74 | 15.0 | 494 | 69 | 14.0 | 485 | 66 | 13.6 | 12 | 2.5 | 384 | 18 | 4.7 | 15 | 3.9 |
| MSW | 2,889 | 51.6 | 1,704 | 51.6 | 1,653 | 95 | 5.7 | 1,655 | 29 | 1.8 | 1,572 | 42 | 2.7 | 3 | 0.2 | 1483 | 2 | 0.1 | 2 | 0.1 |
| WSM | 1,400 | 25.0 | 832 | 25.2 | 770 | 55 | 7.1 | 772 | 6 | 0.8 | 740 | 34 | 4.6 | 2 | 0.3 | 683 | 1 | 0.1 | 1 | 0.1 |
| WSW/M | 79 | 1.4 | 52 | 1.6 | 50 | 3 | 6.0 | 50 | 0 | 0.0 | 44 | 0 | 0.0 | 0 | 0.0 | 42 | 1 | 2.4 | 0 | 0.0 |
| Unknown | 374 | 6.7 | 185 | 5.6 | 133 | 8 | 6.0 | 134 | 5 | 3.7 | 173 | 15 | 8.7 | 0 | 0.0 | 128 | 3 | 2.3 | 3 | 2.3 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 6 | 0.1 | 4 | 0.1 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 200 | 3.6 | 134 | 4.1 | 128 | 19 | 14.8 | 128 | 2 | 1.6 | 116 | 5 | 4.3 | 1 | 0.9 | 109 | 1 | 0.9 | 1 | 0.9 |
| 20-24 | 1,022 | 18.3 | 659 | 20.0 | 635 | 76 | 12.0 | 635 | 29 | 4.6 | 602 | 23 | 3.8 | 3 | 0.5 | 565 | 2 | 0.4 | 2 | 0.4 |
| 25-29 | 1,248 | 22.3 | 758 | 23.0 | 729 | 60 | 8.2 | 729 | 32 | 4.4 | 700 | 27 | 3.9 | 6 | 0.9 | 654 | 8 | 1.2 | 6 | 0.9 |
| 30-34 | 997 | 17.8 | 578 | 17.5 | 548 | 31 | 5.7 | 550 | 20 | 3.6 | 525 | 16 | 3.0 | 2 | 0.4 | 481 | 5 | 1.0 | 4 | 0.8 |
| 35-44 | 1,224 | 21.9 | 688 | 20.8 | 637 | 34 | 5.3 | 640 | 18 | 2.8 | 621 | 45 | 7.2 | 3 | 0.5 | 532 | 6 | 1.1 | 6 | 1.1 |
| 45-54 | 555 | 9.9 | 300 | 9.1 | 273 | 9 | 3.3 | 274 | 7 | 2.6 | 281 | 17 | 6.0 | 1 | 0.4 | 248 | 3 | 1.2 | 2 | 0.8 |
| 55-64 | 270 | 4.8 | 147 | 4.5 | 123 | 6 | 4.9 | 123 | 1 | 0.8 | 132 | 15 | 11.4 | 1 | 0.8 | 105 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 74 | 1.3 | 34 | 1.0 | 23 | 0 | 0.0 | 23 | 0 | 0.0 | 34 | 9 | 26.5 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 2,002 | 35.8 | 1,200 | 36.3 | 1,149 | 66 | 5.7 | 1,152 | 27 | 2.3 | 1,090 | 26 | 2.4 | 4 | 0.4 | 1,021 | 5 | 0.5 | 4 | 0.4 |
| African American | 807 | 14.4 | 514 | 15.6 | 490 | 57 | 11.6 | 490 | 32 | 6.5 | 453 | 28 | 6.2 | 5 | 1.1 | 397 | 6 | 1.5 | 4 | 1.0 |
| Hispanic | 2,180 | 39.0 | 1,224 | 37.1 | 1,108 | 83 | 7.5 | 1,111 | 38 | 3.4 | 1,133 | 92 | 8.1 | 7 | 0.6 | 991 | 12 | 1.2 | 11 | 1.1 |
| Asian | 317 | 5.7 | 185 | 5.6 | 177 | 17 | 9.6 | 177 | 7 | 4.0 | 176 | 6 | 3.4 | 0 | 0.0 | 163 | 1 | 0.6 | 1 | 0.6 |
| Native American | 25 | 0.4 | 16 | 0.5 | 16 | 2 | 12.5 | 16 | 1 | 6.3 | 16 | 2 | 12.5 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 |
| Other | 238 | 4.3 | 147 | 4.5 | 143 | 10 | 7.0 | 143 | 3 | 2.1 | 131 | 3 | 2.3 | 1 | 0.8 | 122 | 1 | 0.8 | 1 | 0.8 |
| Unknown | 27 | 0.5 | 16 | 0.5 | 16 | 0 | 0.0 | 16 | 1 | 6.3 | 15 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit*** | | | | | | | | | | | | | | | | | | | | |
| New Visit | 3,185 | 56.9 | 2,981 | 90.3 | 2,936 | 217 | 7.4 | 2,937 | 106 | 3.6 | 2,788 | 137 | 4.9 | 15 | 0.5 | 2,636 | 23 | 0.9 | 20 | 0.8 |
| Return Visit | 2,411 | 43.1 | 321 | 9.7 | 163 | 18 | 11.0 | 168 | 3 | 1.8 | 226 | 20 | 8.8 | 2 | 0.9 | 84 | 2 | 2.4 | 1 | 1.2 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 3 male-to-female (MTF) and 13 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 7.2 Patient Visits and Tests by Gender and Age, January - December 2010, North Hollywood Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|--------------------------|-------|---------------|------------|------------|---------------|------------|------------|---------------|------------|------------|----------------------------|-------------|---------------|------------|------------|------------|-----|
| | Total STD Visits [†] | | No. Visits with STD Test | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | New HIV*** | |
| | | | | | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | n | % | Visits Tested | Positive n | Positive % | | |
| n | % | n | % | | | | | | | | | | | | | | | | | |
| Total | 5,596 | 100.0 | 3,302 | 100.0 | 3,099 | 235 | 7.6 | 3,105 | 109 | 3.5 | 3,014 | 157 | 5.2 | 17 | 0.6 | 2,720 | 25 | 0.9 | 21 | 0.8 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 104 | 2.6 | 68 | 2.9 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 701 | 17.6 | 443 | 18.9 | 66 | 7 | 10.6 | 66 | 2 | 3.0 | 64 | 2 | 3.1 | 0 | 0.0 | 61 | 1 | 1.6 | 1 | 1.6 |
| 20-24 | 962 | 24.2 | 581 | 24.8 | 424 | 51 | 12.0 | 424 | 28 | 6.6 | 409 | 20 | 4.9 | 3 | 0.7 | 377 | 2 | 0.5 | 2 | 0.5 |
| 25-29 | 697 | 17.5 | 408 | 17.4 | 566 | 51 | 9.0 | 566 | 32 | 5.7 | 539 | 21 | 3.9 | 6 | 1.1 | 500 | 8 | 1.6 | 6 | 1.2 |
| 30-34 | 862 | 21.7 | 492 | 21.0 | 392 | 25 | 6.4 | 394 | 15 | 3.8 | 371 | 11 | 3.0 | 2 | 0.5 | 349 | 4 | 1.1 | 3 | 0.9 |
| 35-44 | 415 | 10.4 | 224 | 9.6 | 465 | 28 | 6.0 | 466 | 16 | 3.4 | 448 | 32 | 7.1 | 3 | 0.7 | 382 | 6 | 1.6 | 6 | 1.6 |
| 45-54 | 197 | 5.0 | 108 | 4.6 | 208 | 8 | 3.8 | 208 | 6 | 2.9 | 210 | 15 | 7.1 | 1 | 0.5 | 184 | 2 | 1.1 | 2 | 1.1 |
| 55-64 | 40 | 1.0 | 19 | 0.8 | 96 | 5 | 5.2 | 96 | 1 | 1.0 | 98 | 10 | 10.2 | 0 | 0.0 | 80 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 0 | 0.0 | 0 | 0.0 | 15 | 0 | 0.0 | 15 | 0 | 0.0 | 19 | 6 | 31.6 | 0 | 0.0 | 15 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,978 | 100.0 | 2,343 | 100.0 | 2,232 | 175 | 7.8 | 2,235 | 100 | 4.5 | 2,158 | 117 | 5.4 | 15 | 0.7 | 1,948 | 23 | 1.2 | 20 | 1.0 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 6 | 0.4 | 4 | 0.4 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 96 | 6.0 | 66 | 6.9 | 62 | 12 | 19.4 | 62 | 0 | 0.0 | 52 | 3 | 5.8 | 1 | 1.9 | 48 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 320 | 19.9 | 216 | 22.7 | 211 | 25 | 11.8 | 211 | 1 | 0.5 | 193 | 3 | 1.6 | 0 | 0.0 | 188 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 283 | 17.6 | 175 | 18.4 | 161 | 9 | 5.6 | 161 | 0 | 0.0 | 159 | 5 | 3.1 | 0 | 0.0 | 152 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 297 | 18.5 | 169 | 17.8 | 155 | 6 | 3.9 | 155 | 4 | 2.6 | 153 | 5 | 3.3 | 0 | 0.0 | 131 | 1 | 0.8 | 1 | 0.8 |
| 35-44 | 356 | 22.2 | 191 | 20.1 | 168 | 6 | 3.6 | 169 | 0 | 0.0 | 169 | 13 | 7.7 | 0 | 0.0 | 149 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 140 | 8.7 | 76 | 8.0 | 65 | 1 | 1.5 | 66 | 1 | 1.5 | 71 | 2 | 2.8 | 0 | 0.0 | 64 | 1 | 1.6 | 0 | 0.0 |
| 55-64 | 73 | 4.5 | 39 | 4.1 | 27 | 1 | 3.7 | 27 | 0 | 0.0 | 34 | 5 | 14.7 | 1 | 2.9 | 25 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 34 | 2.1 | 15 | 1.6 | 8 | 0 | 0.0 | 8 | 0 | 0.0 | 15 | 3 | 20.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,605 | 100.0 | 951 | 100.0 | 860 | 60 | 7.0 | 862 | 6 | 0.7 | 849 | 39 | 4.6 | 2 | 0.2 | 768 | 2 | 0.3 | 1 | 0.1 |
| Others † | 13 | ~ | 8 | ~ | 7 | 0 | 0.0 | 8 | 3 | 37.5 | 7 | 1 | 14.3 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |

†Demographic and visit type information not available; pending resolution of logistical issues pertaining to data entry

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 7.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, North Hollywood Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 5,596 | 100.0 | 3,302 | 100.0 | 3,099 | 235 | 7.6 | 3,105 | 109 | 3.5 | 3,014 | 157 | 5.2 | 17 | 0.6 | 2,720 | 25 | 0.9 | 21 | 0.8 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 1,487 | 37.4 | 903 | 38.5 | 871 | 51 | 5.9 | 873 | 24 | 2.7 | 826 | 23 | 2.8 | 2 | 0.2 | 766 | 5 | 0.7 | 4 | 0.5 |
| African American | 583 | 14.7 | 366 | 15.6 | 350 | 45 | 12.9 | 350 | 31 | 8.9 | 323 | 21 | 6.5 | 5 | 1.5 | 274 | 4 | 1.5 | 3 | 1.1 |
| Hispanic | 1,506 | 37.9 | 840 | 35.9 | 786 | 61 | 7.8 | 787 | 34 | 4.3 | 790 | 63 | 8.0 | 7 | 0.9 | 710 | 12 | 1.7 | 11 | 1.5 |
| Asian | 197 | 5.0 | 112 | 4.8 | 107 | 12 | 11.2 | 107 | 7 | 6.5 | 109 | 5 | 4.6 | 0 | 0.0 | 99 | 1 | 1.0 | 1 | 1.0 |
| Native American | 14 | 0.4 | 10 | 0.4 | 10 | 1 | 10.0 | 10 | 0 | 0.0 | 10 | 2 | 20.0 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| Other | 169 | 4.2 | 100 | 4.3 | 96 | 5 | 5.2 | 96 | 3 | 3.1 | 89 | 3 | 3.4 | 1 | 1.1 | 82 | 1 | 1.2 | 1 | 1.2 |
| Unknown | 22 | 0.6 | 12 | 0.5 | 12 | 0 | 0.0 | 12 | 1 | 8.3 | 11 | 0 | 0.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,978 | 100.0 | 2,343 | 100.0 | 2,232 | 175 | 7.8 | 2,235 | 100 | 4.5 | 2,158 | 117 | 5.4 | 15 | 0.7 | 1,948 | 23 | 1.2 | 20 | 1.0 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 513 | 32.0 | 296 | 31.1 | 277 | 15 | 5.4 | 278 | 3 | 1.1 | 263 | 3 | 1.1 | 2 | 0.8 | 254 | 0 | 0.0 | 0 | 0.0 |
| African American | 222 | 13.8 | 146 | 15.4 | 138 | 12 | 8.7 | 138 | 0 | 0.0 | 128 | 7 | 5.5 | 0 | 0.0 | 123 | 2 | 1.6 | 1 | 0.8 |
| Hispanic | 667 | 41.6 | 380 | 40.0 | 319 | 22 | 6.9 | 320 | 2 | 0.6 | 340 | 28 | 8.2 | 0 | 0.0 | 278 | 0 | 0.0 | 0 | 0.0 |
| Asian | 120 | 7.5 | 73 | 7.7 | 70 | 5 | 7.1 | 70 | 0 | 0.0 | 67 | 1 | 1.5 | 0 | 0.0 | 64 | 0 | 0.0 | 0 | 0.0 |
| Native American | 11 | 0.7 | 6 | 0.6 | 6 | 1 | 16.7 | 6 | 1 | 16.7 | 6 | 0 | 0.0 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 |
| Other | 67 | 4.2 | 46 | 4.8 | 46 | 5 | 10.9 | 46 | 0 | 0.0 | 41 | 0 | 0.0 | 0 | 0.0 | 40 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 5 | 0.3 | 4 | 0.4 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,605 | 100.0 | 951 | 100.0 | 860 | 60 | 7.0 | 862 | 6 | 0.7 | 849 | 39 | 4.6 | 2 | 0.2 | 768 | 2 | 0.3 | 1 | 0.1 |
| Others † | 13 | ~ | 8 | ~ | 7 | 0.0 | 0.0 | 8 | 3.0 | 37.5 | 7 | 1.0 | 14.3 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 7.1a Sexually Transmitted Disease Prevalence, 2006-2010


North Hollywood Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 7.1b Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

North Hollywood Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

VIII: Pomona Health Center

Table 8.1 Patient Visits and Tests, January - December 2010, Pomona Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|------------------|-----------------|-------------------------------|------|-------------|---|-----|-------------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | New HIV*** n % | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | Visits Tested | Positive n % | n % | | | | | | |
| Total | 1,900 | 100.0 | 1,060 | 100.0 | 987 | 167 | 16.9 | 990 | 40 | 4.0 | 965 | 48 | 5.0 | 6 | 0.6 | 884 | 4 | 0.5 | 4 | 0.5 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 1,265 | 66.6 | 712 | 67.2 | 671 | 97 | 14.5 | 674 | 32 | 4.7 | 647 | 31 | 4.8 | 5 | 0.8 | 608 | 4 | 0.7 | 4 | 0.7 |
| Female | 626 | 32.9 | 348 | 32.8 | 316 | 70 | 22.2 | 316 | 8 | 2.5 | 318 | 17 | 5.3 | 1 | 0.3 | 276 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | 9 | 0.5 | 0 | 0.0 | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | -- | 0 | 0 | -- | 0 | -- |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 330 | 17.4 | 178 | 16.8 | 163 | 24 | 14.7 | 165 | 23 | 13.9 | 156 | 19 | 12.2 | 3 | 1.9 | 143 | 3 | 2.1 | 3 | 2.1 |
| MSW | 893 | 47.0 | 509 | 48.0 | 488 | 72 | 14.8 | 489 | 9 | 1.8 | 467 | 12 | 2.6 | 1 | 0.2 | 449 | 1 | 0.2 | 1 | 0.2 |
| WSM | 498 | 26.2 | 289 | 27.3 | 272 | 65 | 23.9 | 272 | 8 | 2.9 | 260 | 12 | 4.6 | 1 | 0.4 | 235 | 0 | 0.0 | 0 | 0.0 |
| WSW/M | 45 | 2.4 | 23 | 2.2 | 23 | 2 | 8.7 | 23 | 0 | 0.0 | 23 | 1 | 4.3 | 0 | 0.0 | 21 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 134 | 7.1 | 62 | 5.8 | 41 | 4 | 9.8 | 41 | 0 | 0.0 | 60 | 4 | 6.7 | 1 | 1.7 | 36 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 9 | 0.5 | 4 | 0.4 | 4 | 3 | 75.0 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 192 | 10.1 | 110 | 10.4 | 106 | 36 | 34.0 | 107 | 11 | 10.3 | 99 | 2 | 2.0 | 0 | 0.0 | 97 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 528 | 27.8 | 307 | 29.0 | 298 | 65 | 21.8 | 298 | 18 | 6.0 | 281 | 6 | 2.1 | 1 | 0.4 | 259 | 1 | 0.4 | 1 | 0.4 |
| 25-29 | 380 | 20.0 | 206 | 19.4 | 197 | 32 | 16.2 | 197 | 4 | 2.0 | 186 | 12 | 6.5 | 1 | 0.5 | 178 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 213 | 11.2 | 126 | 11.9 | 117 | 16 | 13.7 | 118 | 2 | 1.7 | 111 | 4 | 3.6 | 0 | 0.0 | 103 | 1 | 1.0 | 1 | 1.0 |
| 35-44 | 313 | 16.5 | 163 | 15.4 | 149 | 14 | 9.4 | 150 | 2 | 1.3 | 149 | 12 | 8.1 | 2 | 1.3 | 137 | 1 | 0.7 | 1 | 0.7 |
| 45-54 | 187 | 9.8 | 107 | 10.1 | 93 | 1 | 1.1 | 93 | 3 | 3.2 | 99 | 10 | 10.1 | 1 | 1.0 | 85 | 1 | 1.2 | 1 | 1.2 |
| 55-64 | 66 | 3.5 | 30 | 2.8 | 21 | 0 | 0.0 | 21 | 0 | 0.0 | 29 | 2 | 6.9 | 0 | 0.0 | 19 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 12 | 0.6 | 8 | 0.8 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 8 | 0 | 0.0 | 1 | 12.5 | 2 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 273 | 14.4 | 157 | 14.8 | 150 | 25 | 16.7 | 151 | 7 | 4.6 | 142 | 0 | 0.0 | 0 | 0.0 | 136 | 2 | 1.5 | 2 | 1.5 |
| African American | 304 | 16.0 | 177 | 16.7 | 166 | 28 | 16.9 | 167 | 6 | 3.6 | 162 | 10 | 6.2 | 1 | 0.6 | 145 | 1 | 0.7 | 1 | 0.7 |
| Hispanic | 1,115 | 58.7 | 614 | 57.9 | 565 | 103 | 18.2 | 566 | 24 | 4.2 | 557 | 35 | 6.3 | 4 | 0.7 | 505 | 1 | 0.2 | 1 | 0.2 |
| Asian | 145 | 7.6 | 77 | 7.3 | 70 | 8 | 11.4 | 70 | 1 | 1.4 | 72 | 3 | 4.2 | 1 | 1.4 | 65 | 0 | 0.0 | 0 | 0.0 |
| Native American | 5 | 0.3 | 3 | 0.3 | 3 | 1 | 33.3 | 3 | 1 | 33.3 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Other | 46 | 2.4 | 26 | 2.5 | 26 | 2 | 7.7 | 26 | 1 | 3.8 | 23 | 0 | 0.0 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 12 | 0.6 | 7 | 0.7 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 921 | 48.5 | 904 | 85.3 | 900 | 155 | 17.2 | 902 | 36 | 4.0 | 867 | 42 | 4.8 | 4 | 0.5 | 829 | 3 | 0.4 | 3 | 0.4 |
| Return Visit | 979 | 51.5 | 157 | 14.8 | 87 | 12 | 13.8 | 88 | 4 | 4.5 | 99 | 6 | 6.1 | 2 | 2.0 | 55 | 1 | 1.8 | 1 | 1.8 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 9 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 8.2 Patient Visits and Tests by Gender and Age, January - December 2010, Pomona Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|---------------|------------|------------|---------------|------------|------------|---------------|------------|------------|-----------------------------------|---------------|------------|-------------------|------------|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | HIV | | | New HIV*** n % | | | |
| | | | | | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Confirmed Early Syphilis** n % | Visits Tested | Positive n | | Positive % | | |
| Total | 1,900 | 100.0 | 1,061 | 100.0 | 987 | 167 | 16.9 | 990 | 40 | 4.0 | 966 | 48 | 5.0 | 6 | 0.6 | 884 | 4 | 0.5 | 4 | 0.5 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 1 | 0.1 | 64 | 9.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 110 | 8.7 | 195 | 27.4 | 62 | 18 | 29.0 | 63 | 8 | 12.7 | 58 | 2 | 3.4 | 0 | 0.0 | 57 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 337 | 26.6 | 158 | 22.2 | 191 | 35 | 18.3 | 191 | 15 | 7.9 | 179 | 4 | 2.2 | 1 | 0.6 | 169 | 1 | 0.6 | 1 | 0.6 |
| 25-29 | 287 | 22.7 | 91 | 12.8 | 150 | 22 | 14.7 | 150 | 4 | 2.7 | 143 | 7 | 4.9 | 1 | 0.7 | 138 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 153 | 12.1 | 106 | 14.9 | 84 | 12 | 14.3 | 85 | 2 | 2.4 | 78 | 3 | 3.8 | 0 | 0.0 | 74 | 1 | 1.4 | 1 | 1.4 |
| 35-44 | 206 | 16.3 | 78 | 11.0 | 99 | 9 | 9.1 | 100 | 2 | 2.0 | 96 | 5 | 5.2 | 1 | 1.0 | 93 | 1 | 1.1 | 1 | 1.1 |
| 45-54 | 133 | 10.5 | 16 | 2.3 | 70 | 1 | 1.4 | 70 | 1 | 1.4 | 73 | 9 | 12.3 | 1 | 1.4 | 62 | 1 | 1.6 | 1 | 1.6 |
| 55-64 | 34 | 2.7 | 3 | 0.4 | 14 | 0 | 0.0 | 14 | 0 | 0.0 | 16 | 1 | 6.3 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 4 | 0.3 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 3 | 0 | 0.0 | 1 | 33.3 | 0 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,265 | 100.0 | 711 | 100.0 | 670 | 97 | 14.5 | 673 | 32 | 4.8 | 646 | 31 | 4.8 | 5 | 0.8 | 607 | 4 | 0.7 | 4 | 0.7 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 8 | 1.3 | 4 | 1.1 | 4 | 3 | 75.0 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 82 | 13.1 | 46 | 13.2 | 44 | 18 | 40.9 | 44 | 3 | 6.8 | 41 | 0 | 0.0 | 0 | 0.0 | 40 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 191 | 30.5 | 112 | 32.2 | 107 | 30 | 28.0 | 107 | 3 | 2.8 | 102 | 2 | 2.0 | 0 | 0.0 | 90 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 85 | 13.6 | 47 | 13.5 | 46 | 10 | 21.7 | 46 | 0 | 0.0 | 42 | 5 | 11.9 | 0 | 0.0 | 39 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 59 | 9.4 | 34 | 9.8 | 33 | 4 | 12.1 | 33 | 0 | 0.0 | 32 | 1 | 3.1 | 0 | 0.0 | 29 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 107 | 17.1 | 57 | 16.4 | 50 | 5 | 10.0 | 50 | 0 | 0.0 | 53 | 7 | 13.2 | 1 | 1.9 | 44 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 54 | 8.6 | 29 | 8.3 | 23 | 0 | 0.0 | 23 | 2 | 8.7 | 26 | 1 | 3.8 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 32 | 5.1 | 14 | 4.0 | 7 | 0 | 0.0 | 7 | 0 | 0.0 | 13 | 1 | 7.7 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 8 | 1.3 | 5 | 1.4 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 626 | 100.0 | 348 | 100.0 | 316 | 70 | 22.2 | 316 | 8 | 2.5 | 318 | 17 | 5.3 | 1 | 0.3 | 276 | 0 | 0.0 | 0 | 0.0 |
| Others † | 9 | ~ | 2 | ~ | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 2 | 0.0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 8.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Pomona Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|------------------|-----------------|-------------------------------|-------------|-----|-----|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 1,900 | 100.0 | 1,061 | 100.0 | 987 | 167 | 16.9 | 990 | 40 | 4.0 | 966 | 48 | 5.0 | 6 | 0.6 | 884 | 4 | 0.5 | 4 | 0.5 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 196 | 15.5 | 117 | 16.5 | 112 | 17 | 15.2 | 113 | 6 | 5.3 | 108 | 0 | 0.0 | 0 | 0.0 | 103 | 2 | 1.9 | 2 | 1.9 |
| African American | 195 | 15.4 | 117 | 16.5 | 110 | 18 | 16.4 | 111 | 5 | 4.5 | 106 | 5 | 4.7 | 1 | 0.9 | 94 | 1 | 1.1 | 1 | 1.1 |
| Hispanic | 729 | 57.6 | 401 | 56.4 | 375 | 58 | 15.5 | 376 | 19 | 5.1 | 360 | 23 | 6.4 | 3 | 0.8 | 340 | 1 | 0.3 | 1 | 0.3 |
| Asian | 97 | 7.7 | 51 | 7.2 | 48 | 3 | 6.3 | 48 | 1 | 2.1 | 48 | 3 | 6.3 | 1 | 2.1 | 46 | 0 | 0.0 | 0 | 0.0 |
| Native American | 5 | 0.4 | 3 | 0.4 | 3 | 1 | 33.3 | 3 | 1 | 33.3 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Other | 36 | 2.8 | 18 | 2.5 | 18 | 0 | 0.0 | 18 | 0 | 0.0 | 17 | 0 | 0.0 | 0 | 0.0 | 17 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 7 | 0.6 | 4 | 0.6 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,265 | 100.0 | 711 | 100.0 | 670 | 97 | 14.5 | 673 | 32 | 4.8 | 646 | 31 | 4.8 | 5 | 0.8 | 607 | 4 | 0.7 | 4 | 0.7 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 77 | 12.3 | 40 | 11.5 | 38 | 8 | 21.1 | 38 | 1 | 2.6 | 34 | 0 | 0.0 | 0 | 0.0 | 33 | 0 | 0.0 | 0 | 0.0 |
| African American | 109 | 17.4 | 60 | 17.2 | 56 | 10 | 17.9 | 56 | 1 | 1.8 | 56 | 5 | 8.9 | 0 | 0.0 | 51 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 377 | 60.2 | 211 | 60.6 | 189 | 45 | 23.8 | 189 | 5 | 2.6 | 195 | 12 | 6.2 | 1 | 0.5 | 164 | 0 | 0.0 | 0 | 0.0 |
| Asian | 48 | 7.7 | 26 | 7.5 | 22 | 5 | 22.7 | 22 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 | 19 | 0 | 0.0 | 0 | 0.0 |
| Native American | 10 | 1.6 | 8 | 2.3 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Other | 5 | 0.8 | 3 | 0.9 | 8 | 2 | 25.0 | 8 | 1 | 12.5 | 6 | 0 | 0.0 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 626 | 100.0 | 348 | 100.0 | 316 | 70 | 22.2 | 316 | 8 | 2.5 | 318 | 17 | 5.3 | 1 | 0.3 | 276 | 0 | 0.0 | 0 | 0.0 |
| Others † | 9 | ~ | 2 | ~ | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 2 | 0.0 | 0.0 | 0 | 0.0 | 1 | 0.0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 8.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Pomona Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 8.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Pomona Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

IX: South Health Center

Table 9.1 Patient Visits and Tests, January - December 2010, South Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|-------------------------|------------------|-----------------|-------------------|-----|-------------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | Early Syphilis** n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 6,713 | 100.0 | 5,139 | 100.0 | 5,091 | 721 | 14.2 | 5,095 | 323 | 6.3 | 4,513 | 153 | 3.4 | 15 | 0.3 | 4,306 | 17 | 0.4 | 12 | 0.3 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 3,554 | 52.9 | 2,713 | 52.8 | 2,686 | 371 | 13.8 | 2,688 | 195 | 7.3 | 2,415 | 89 | 3.7 | 13 | 0.5 | 2,306 | 14 | 0.6 | 9 | 0.4 |
| Female | 3,154 | 47.0 | 2,422 | 47.1 | 2,401 | 350 | 14.6 | 2,403 | 128 | 5.3 | 2,095 | 64 | 3.1 | 2 | 0.1 | 1,997 | 3 | 0.2 | 3 | 0.2 |
| Transgender† | 5 | 0.1 | 4 | 0.1 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 261 | 3.9 | 175 | 3.4 | 172 | 14 | 8.1 | 172 | 22 | 12.8 | 165 | 32 | 19.4 | 7 | 4.2 | 148 | 8 | 5.4 | 6 | 4.1 |
| MSW | 3,173 | 47.3 | 2,472 | 48.1 | 2,453 | 355 | 14.5 | 2,455 | 172 | 7.0 | 2,187 | 45 | 2.1 | 5 | 0.2 | 2,100 | 6 | 0.3 | 3 | 0.1 |
| WSM | 2,730 | 40.7 | 2,125 | 41.4 | 2,109 | 317 | 15.0 | 2,111 | 104 | 4.9 | 1,838 | 53 | 2.9 | 2 | 0.1 | 1,751 | 3 | 0.2 | 3 | 0.2 |
| WSW/M | 281 | 4.2 | 215 | 4.2 | 214 | 20 | 9.3 | 214 | 22 | 10.3 | 177 | 4 | 2.3 | 0 | 0.0 | 169 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 268 | 4.0 | 152 | 3.0 | 143 | 15 | 10.5 | 143 | 3 | 2.1 | 146 | 19 | 13.0 | 1 | 0.7 | 138 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 15 | 0.2 | 13 | 0.3 | 13 | 3 | 23.1 | 13 | 1 | 7.7 | 12 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 923 | 13.7 | 727 | 14.1 | 724 | 213 | 29.4 | 724 | 70 | 9.7 | 642 | 6 | 0.9 | 2 | 0.3 | 612 | 1 | 0.2 | 1 | 0.2 |
| 20-24 | 1,698 | 25.3 | 1,335 | 26.0 | 1,326 | 282 | 21.3 | 1,327 | 122 | 9.2 | 1,171 | 18 | 1.5 | 4 | 0.3 | 1,129 | 6 | 0.5 | 4 | 0.4 |
| 25-29 | 1,108 | 16.5 | 873 | 17.0 | 865 | 126 | 14.6 | 866 | 52 | 6.0 | 762 | 16 | 2.1 | 3 | 0.4 | 728 | 3 | 0.4 | 3 | 0.4 |
| 30-34 | 775 | 11.5 | 594 | 11.6 | 592 | 43 | 7.3 | 593 | 34 | 5.7 | 516 | 9 | 1.7 | 1 | 0.2 | 491 | 2 | 0.4 | 2 | 0.4 |
| 35-44 | 1,083 | 16.1 | 821 | 16.0 | 814 | 38 | 4.7 | 815 | 28 | 3.4 | 726 | 35 | 4.8 | 1 | 0.1 | 693 | 2 | 0.3 | 1 | 0.1 |
| 45-54 | 822 | 12.2 | 585 | 11.4 | 574 | 13 | 2.3 | 574 | 8 | 1.4 | 510 | 36 | 7.1 | 3 | 0.6 | 480 | 1 | 0.2 | 0 | 0.0 |
| 55-64 | 226 | 3.4 | 153 | 3.0 | 147 | 3 | 2.0 | 147 | 8 | 5.4 | 139 | 27 | 19.4 | 1 | 0.7 | 128 | 2 | 1.6 | 1 | 0.8 |
| 65+ | 63 | 0.9 | 38 | 0.7 | 36 | 0 | 0.0 | 36 | 0 | 0.0 | 35 | 6 | 17.1 | 0 | 0.0 | 33 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 148 | 2.2 | 94 | 1.8 | 93 | 5 | 5.4 | 93 | 3 | 3.2 | 91 | 5 | 5.5 | 1 | 1.1 | 87 | 0 | 0.0 | 0 | 0.0 |
| African American | 4,761 | 70.9 | 3,795 | 73.8 | 3,769 | 573 | 15.2 | 3,772 | 289 | 7.7 | 3,282 | 67 | 2.0 | 9 | 0.3 | 3,128 | 13 | 0.4 | 8 | 0.3 |
| Hispanic | 1,708 | 25.4 | 1,178 | 22.9 | 1,159 | 132 | 11.4 | 1,160 | 26 | 2.2 | 1,076 | 75 | 7.0 | 4 | 0.4 | 1,030 | 4 | 0.4 | 4 | 0.4 |
| Asian | 32 | 0.5 | 23 | 0.4 | 21 | 5 | 23.8 | 21 | 0 | 0.0 | 21 | 3 | 14.3 | 0 | 0.0 | 20 | 0 | 0.0 | 0 | 0.0 |
| Native American | 15 | 0.2 | 11 | 0.2 | 11 | 1 | 9.1 | 11 | 1 | 9.1 | 10 | 2 | 20.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Other | 42 | 0.6 | 34 | 0.7 | 34 | 5 | 14.7 | 34 | 2 | 5.9 | 29 | 1 | 3.4 | 1 | 3.4 | 28 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 7 | 0.1 | 4 | 0.1 | 4 | 0 | 0.0 | 4 | 2 | 50.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 5,128 | 76.4 | 4,996 | 97.2 | 4,968 | 714 | 14.4 | 4,970 | 318 | 6.4 | 4,487 | 145 | 3.2 | 15 | 0.3 | 4,288 | 16 | 0.4 | 12 | 0.3 |
| Return Visit | 1,585 | 23.6 | 143 | 2.8 | 123 | 7 | 5.7 | 125 | 5 | 4.0 | 26 | 8 | 30.8 | 0 | 0.0 | 18 | 1 | 5.6 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 3 male-to-female (MTF) and 2 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 9.2 Patient Visits and Tests by Gender and Age, January - December 2010, South Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|------------------|-----------------|-------------------------------|-------------|-------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | | |
| Total | 6,713 | 100.0 | 5,139 | 100.0 | 5,091 | 721 | 14.2 | 5,095 | 323 | 6.3 | 4,513 | 153 | 3.4 | 15 | 0.3 | 4,306 | 17 | 0.4 | 12 | 0.3 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 4 | 0.1 | 4 | 0.1 | 4 | 1 | 25.0 | 4 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 317 | 8.9 | 253 | 9.3 | 252 | 73 | 29.0 | 252 | 34 | 13.5 | 230 | 5 | 2.2 | 2 | 0.9 | 222 | 1 | 0.5 | 1 | 0.5 |
| 20-24 | 850 | 23.9 | 674 | 24.8 | 670 | 151 | 22.5 | 670 | 67 | 10.0 | 596 | 13 | 2.2 | 4 | 0.7 | 575 | 5 | 0.9 | 3 | 0.5 |
| 25-29 | 618 | 17.4 | 477 | 17.6 | 475 | 80 | 16.8 | 475 | 36 | 7.6 | 424 | 10 | 2.4 | 3 | 0.7 | 406 | 3 | 0.7 | 3 | 0.7 |
| 30-34 | 491 | 13.8 | 369 | 13.6 | 368 | 28 | 7.6 | 369 | 23 | 6.2 | 322 | 7 | 2.2 | 1 | 0.3 | 308 | 1 | 0.3 | 1 | 0.3 |
| 35-44 | 605 | 17.0 | 467 | 17.2 | 461 | 26 | 5.6 | 462 | 22 | 4.8 | 416 | 16 | 3.8 | 0 | 0.0 | 394 | 2 | 0.5 | 1 | 0.3 |
| 45-54 | 471 | 13.2 | 336 | 12.4 | 329 | 9 | 2.7 | 329 | 6 | 1.8 | 302 | 19 | 6.3 | 2 | 0.7 | 285 | 1 | 0.4 | 0 | 0.0 |
| 55-64 | 147 | 4.1 | 104 | 3.8 | 100 | 3 | 3.0 | 100 | 7 | 7.0 | 93 | 15 | 16.1 | 1 | 1.1 | 86 | 1 | 1.2 | 0 | 0.0 |
| 65+ | 54 | 1.5 | 32 | 1.2 | 30 | 0 | 0.0 | 30 | 0 | 0.0 | 30 | 4 | 13.3 | 0 | 0.0 | 28 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,557 | 100.0 | 2,716 | 100.0 | 2,689 | 371 | 13.8 | 2,691 | 195 | 7.2 | 2,417 | 89 | 3.7 | 13 | 0.5 | 2,308 | 14 | 0.6 | 9 | 0.4 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 11 | 0.3 | 9 | 0.4 | 9 | 2 | 22.2 | 9 | 1 | 11.1 | 8 | 0 | 0.0 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 606 | 19.2 | 474 | 19.6 | 472 | 140 | 29.7 | 472 | 36 | 7.6 | 412 | 1 | 0.2 | 0 | 0.0 | 390 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 848 | 26.9 | 661 | 27.3 | 656 | 131 | 20.0 | 657 | 55 | 8.4 | 575 | 5 | 0.9 | 0 | 0.0 | 554 | 1 | 0.2 | 1 | 0.2 |
| 25-29 | 490 | 15.5 | 396 | 16.4 | 390 | 46 | 11.8 | 391 | 16 | 4.1 | 338 | 6 | 1.8 | 0 | 0.0 | 322 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 284 | 9.0 | 225 | 9.3 | 224 | 15 | 6.7 | 224 | 11 | 4.9 | 194 | 2 | 1.0 | 0 | 0.0 | 183 | 1 | 0.5 | 1 | 0.5 |
| 35-44 | 476 | 15.1 | 353 | 14.6 | 352 | 12 | 3.4 | 352 | 6 | 1.7 | 309 | 19 | 6.1 | 1 | 0.3 | 298 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 351 | 11.1 | 249 | 10.3 | 245 | 4 | 1.6 | 245 | 2 | 0.8 | 208 | 17 | 8.2 | 1 | 0.5 | 195 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 79 | 2.5 | 49 | 2.0 | 47 | 0 | 0.0 | 47 | 1 | 2.1 | 46 | 12 | 26.1 | 0 | 0.0 | 42 | 1 | 2.4 | 1 | 2.4 |
| 65+ | 9 | 0.3 | 6 | 0.2 | 6 | 0 | 0.0 | 6 | 0 | 0.0 | 5 | 2 | 40.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,154 | 100.0 | 2,422 | 100.0 | 2,401 | 350 | 14.6 | 2,403 | 128 | 5.3 | 2,095 | 64 | 3.1 | 2 | 0.1 | 1,997 | 3 | 0.2 | 3 | 0.2 |
| Others † | 2 | ~ | 1 | ~ | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 9.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, South Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 6,713 | 100.0 | 5,139 | 100.0 | 5,091 | 721 | 14.2 | 5,095 | 323 | 6.3 | 4,513 | 153 | 3.4 | 15 | 0.3 | 4,306 | 17 | 0.4 | 12 | 0.3 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 75 | 2.1 | 46 | 1.7 | 45 | 1 | 2.2 | 45 | 1 | 2.2 | 46 | 3 | 6.5 | 1 | 2.2 | 44 | 0 | 0.0 | 0 | 0.0 |
| African American | 2,372 | 66.7 | 1,896 | 69.8 | 1,882 | 299 | 15.9 | 1,883 | 174 | 9.2 | 1,664 | 33 | 2.0 | 7 | 0.4 | 1,590 | 10 | 0.6 | 5 | 0.3 |
| Hispanic | 1,062 | 29.9 | 735 | 27.1 | 723 | 66 | 9.1 | 724 | 16 | 2.2 | 674 | 52 | 7.7 | 4 | 0.6 | 642 | 4 | 0.6 | 4 | 0.6 |
| Asian | 17 | 0.5 | 14 | 0.5 | 14 | 3 | 21.4 | 14 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| Native American | 6 | 0.2 | 5 | 0.2 | 5 | 0 | 0.0 | 5 | 1 | 20.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Other | 18 | 0.5 | 16 | 0.6 | 16 | 2 | 12.5 | 16 | 1 | 6.3 | 13 | 1 | 7.7 | 1 | 7.7 | 12 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 7 | 0.2 | 4 | 0.1 | 4 | 0 | 0.0 | 4 | 2 | 50.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,557 | 100.0 | 2,716 | 100.0 | 2,689 | 371 | 13.8 | 2,691 | 195 | 7.2 | 2,417 | 89 | 3.7 | 13 | 0.5 | 2,308 | 14 | 0.6 | 9 | 0.4 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 73 | 2.3 | 48 | 2.0 | 48 | 4 | 8.3 | 48 | 2 | 4.2 | 45 | 2 | 4.4 | 0 | 0.0 | 43 | 0 | 0.0 | 0 | 0.0 |
| African American | 2,389 | 75.7 | 1,899 | 78.4 | 1,887 | 274 | 14.5 | 1,889 | 115 | 6.1 | 1,618 | 34 | 2.1 | 2 | 0.1 | 1,538 | 3 | 0.2 | 3 | 0.2 |
| Hispanic | 644 | 20.4 | 442 | 18.2 | 435 | 66 | 15.2 | 435 | 10 | 2.3 | 401 | 23 | 5.7 | 0 | 0.0 | 387 | 0 | 0.0 | 0 | 0.0 |
| Asian | 15 | 0.5 | 9 | 0.4 | 7 | 2 | 28.6 | 7 | 0 | 0.0 | 9 | 3 | 33.3 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| Native American | 9 | 0.3 | 6 | 0.2 | 6 | 1 | 16.7 | 6 | 0 | 0.0 | 6 | 2 | 33.3 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Other | 24 | 0.8 | 18 | 0.7 | 18 | 3 | 16.7 | 18 | 1 | 5.6 | 16 | 0 | 0.0 | 0 | 0.0 | 16 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,154 | 100.0 | 2,422 | 100.0 | 2,401 | 350 | 14.6 | 2,403 | 128 | 5.3 | 2,095 | 64 | 3.1 | 2 | 0.1 | 1,997 | 3 | 0.2 | 3 | 0.2 |
| Others † | 2 | ~ | 1 | ~ | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 1 | 0.0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 9.1a. Sexually Transmitted Disease Prevalence, 2006-2010


South Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 9.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

South Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

X: Ruth Temple Health Center

Table 10.1 Patient Visits and Tests, January - December 2010, Ruth Temple Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-------------------|-------|----|------|----|------|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 5,471 | 100.0 | 3,569 | 100.0 | 3,507 | 444 | 12.7 | 3,523 | 229 | 6.5 | 3,158 | 210 | 6.6 | 14 | 0.4 | 2,890 | 28 | 1.0 | 16 | 0.6 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 3,176 | 58.1 | 2,029 | 56.9 | 2,005 | 262 | 13.1 | 2,011 | 155 | 7.7 | 1,832 | 110 | 6.0 | 9 | 0.5 | 1,674 | 24 | 1.4 | 15 | 0.9 |
| Female | 2,292 | 41.9 | 1,538 | 43.1 | 1,500 | 181 | 12.1 | 1,510 | 74 | 4.9 | 1,324 | 99 | 7.5 | 5 | 0.4 | 1,214 | 4 | 0.3 | 1 | 0.1 |
| Transgender† | 3 | 0.1 | 2 | 0.1 | 2 | 1 | 50.0 | 2 | 0 | 0.0 | 2 | 1 | 50.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 235 | 4.3 | 141 | 4.0 | 140 | 20 | 14.3 | 140 | 22 | 15.7 | 134 | 22 | 16.4 | 4 | 3.0 | 111 | 17 | 15.3 | 12 | 10.8 |
| MSW | 2,849 | 52.1 | 1,840 | 51.6 | 1,818 | 238 | 13.1 | 1,824 | 129 | 7.1 | 1,651 | 80 | 4.8 | 5 | 0.3 | 1,519 | 7 | 0.5 | 3 | 0.2 |
| WSM | 2,018 | 36.9 | 1,356 | 38.0 | 1,325 | 166 | 12.5 | 1,333 | 64 | 4.8 | 1,164 | 91 | 7.8 | 5 | 0.4 | 1,072 | 2 | 0.2 | 0 | 0.0 |
| WSW/M | 176 | 3.2 | 130 | 3.6 | 126 | 13 | 10.3 | 128 | 9 | 7.0 | 113 | 1 | 0.9 | 0 | 0.0 | 102 | 1 | 1.0 | 0 | 0.0 |
| Unknown | 193 | 3.5 | 102 | 2.9 | 98 | 6 | 6.1 | 98 | 5 | 5.1 | 96 | 16 | 16.7 | 0 | 0.0 | 86 | 1 | 1.2 | 1 | 1.2 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 8 | 0.1 | 5 | 0.1 | 5 | 1 | 20.0 | 5 | 1 | 20.0 | 5 | 0 | 0.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 522 | 9.5 | 353 | 9.9 | 346 | 100 | 28.9 | 346 | 52 | 15.0 | 312 | 8 | 2.6 | 0 | 0.0 | 294 | 4 | 1.4 | 2 | 0.7 |
| 20-24 | 1,234 | 22.6 | 831 | 23.3 | 822 | 166 | 20.2 | 826 | 75 | 9.1 | 726 | 17 | 2.3 | 5 | 0.7 | 684 | 6 | 0.9 | 6 | 0.9 |
| 25-29 | 983 | 18.0 | 659 | 18.5 | 647 | 71 | 11.0 | 652 | 35 | 5.4 | 569 | 17 | 3.0 | 1 | 0.2 | 527 | 3 | 0.6 | 2 | 0.4 |
| 30-34 | 706 | 12.9 | 445 | 12.5 | 438 | 51 | 11.6 | 442 | 20 | 4.5 | 398 | 23 | 5.8 | 2 | 0.5 | 363 | 7 | 1.9 | 3 | 0.8 |
| 35-44 | 909 | 16.6 | 584 | 16.4 | 573 | 38 | 6.6 | 574 | 22 | 3.8 | 516 | 47 | 9.1 | 1 | 0.2 | 456 | 4 | 0.9 | 2 | 0.4 |
| 45-54 | 800 | 14.6 | 505 | 14.1 | 498 | 11 | 2.2 | 500 | 18 | 3.6 | 457 | 53 | 11.6 | 2 | 0.4 | 410 | 2 | 0.5 | 1 | 0.2 |
| 55-64 | 267 | 4.9 | 164 | 4.6 | 156 | 5 | 3.2 | 156 | 5 | 3.2 | 152 | 36 | 23.7 | 1 | 0.7 | 132 | 1 | 0.8 | 0 | 0.0 |
| 65+ | 42 | 0.8 | 23 | 0.6 | 22 | 0 | 0.0 | 22 | 1 | 4.5 | 23 | 9 | 39.1 | 2 | 8.7 | 19 | 1 | 5.3 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 135 | 2.5 | 89 | 2.5 | 89 | 8 | 9.0 | 89 | 3 | 3.4 | 86 | 3 | 3.5 | 0 | 0.0 | 80 | 1 | 1.3 | 1 | 1.3 |
| African American | 4,312 | 78.8 | 2,862 | 80.2 | 2,815 | 369 | 13.1 | 2,828 | 205 | 7.2 | 2,513 | 145 | 5.8 | 12 | 0.5 | 2,310 | 21 | 0.9 | 13 | 0.6 |
| Hispanic | 881 | 16.1 | 530 | 14.9 | 518 | 56 | 10.8 | 520 | 14 | 2.7 | 486 | 53 | 10.9 | 0 | 0.0 | 431 | 5 | 1.2 | 1 | 0.2 |
| Asian | 61 | 1.1 | 36 | 1.0 | 35 | 3 | 8.6 | 35 | 2 | 5.7 | 30 | 4 | 13.3 | 1 | 3.3 | 29 | 0 | 0.0 | 0 | 0.0 |
| Native American | 4 | 0.1 | 2 | 0.1 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Other | 72 | 1.3 | 47 | 1.3 | 45 | 7 | 15.6 | 46 | 4 | 8.7 | 38 | 5 | 13.2 | 1 | 2.6 | 35 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 6 | 0.1 | 3 | 0.1 | 3 | 0 | 0.0 | 3 | 1 | 33.3 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 1 | 33.3 | 1 | 33.3 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 3,373 | 61.7 | 3,301 | 92.5 | 3,270 | 411 | 12.6 | 3,282 | 216 | 6.6 | 2,994 | 183 | 6.1 | 14 | 0.5 | 2,754 | 24 | 0.9 | 15 | 0.5 |
| Return Visit | 2,098 | 38.3 | 268 | 7.5 | 237 | 32 | 13.5 | 241 | 13 | 5.4 | 164 | 27 | 16.5 | 0 | 0.0 | 136 | 4 | 2.9 | 1 | 0.7 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 3 unspecified

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested; similarly for RPR/TPPA, EIA/WB, etc.)

Table 10.2 Patient Visits and Tests by Gender and Age, January - December 2010, Ruth Temple Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|---------------|------|------------------|---------------|------|------------------|---------------|------|-------------------------------|-------------|------------------|---------------|-----|-----------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | n | % | Visits Tested | Positive n | % | New HIV*** n | % |
| Total | 5,471 | 100.0 | 3,569 | 100.0 | 3,507 | 443 | 12.6 | 3,523 | 229 | 6.5 | 3,158 | 210 | 6.6 | 14 | 0.4 | 2,890 | 28 | 1.0 | 16 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 1 | 0.0 | 1 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 237 | 7.5 | 156 | 7.7 | 153 | 49 | 32.0 | 153 | 25 | 16.3 | 143 | 4 | 2.8 | 0 | 0.0 | 134 | 3 | 2.2 | 2 | 1.5 |
| 20-24 | 625 | 19.7 | 405 | 20.0 | 404 | 87 | 21.5 | 405 | 47 | 11.6 | 370 | 7 | 1.9 | 3 | 0.8 | 348 | 6 | 1.7 | 6 | 1.7 |
| 25-29 | 580 | 18.3 | 378 | 18.6 | 374 | 49 | 13.1 | 376 | 29 | 7.7 | 329 | 9 | 2.7 | 1 | 0.3 | 308 | 3 | 1.0 | 2 | 0.6 |
| 30-34 | 436 | 13.7 | 271 | 13.4 | 267 | 33 | 12.4 | 269 | 12 | 4.5 | 242 | 15 | 6.2 | 2 | 0.8 | 223 | 7 | 3.1 | 3 | 1.3 |
| 35-44 | 568 | 17.9 | 363 | 17.9 | 359 | 32 | 8.9 | 360 | 22 | 6.1 | 326 | 16 | 4.9 | 0 | 0.0 | 293 | 3 | 1.0 | 2 | 0.7 |
| 45-54 | 512 | 16.1 | 320 | 15.8 | 317 | 10 | 3.2 | 317 | 14 | 4.4 | 295 | 30 | 10.2 | 1 | 0.3 | 258 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 184 | 5.8 | 116 | 5.7 | 112 | 2 | 1.8 | 112 | 5 | 4.5 | 107 | 21 | 19.6 | 0 | 0.0 | 94 | 1 | 1.1 | 0 | 0.0 |
| 65+ | 33 | 1.0 | 19 | 0.9 | 18 | 0 | 0.0 | 18 | 1 | 5.6 | 19 | 8 | 42.1 | 2 | 10.5 | 15 | 1 | 6.7 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,176 | 100.0 | 2,029 | 100.0 | 2,005 | 262 | 13.1 | 2,011 | 155 | 7.7 | 1,832 | 110 | 6.0 | 9 | 0.5 | 1,674 | 24 | 1.4 | 15 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 7 | 0.3 | 4 | 0.3 | 4 | 1 | 25.0 | 4 | 1 | 25.0 | 4 | 0 | 0.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 285 | 12.4 | 197 | 12.8 | 193 | 51 | 26.4 | 193 | 27 | 14.0 | 169 | 4 | 2.4 | 0 | 0.0 | 160 | 1 | 0.6 | 0 | 0.0 |
| 20-24 | 609 | 26.6 | 426 | 27.7 | 418 | 79 | 18.9 | 421 | 28 | 6.7 | 356 | 10 | 2.8 | 2 | 0.6 | 336 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 403 | 17.6 | 281 | 18.3 | 273 | 22 | 8.1 | 276 | 6 | 2.2 | 240 | 8 | 3.3 | 0 | 0.0 | 219 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 270 | 11.8 | 174 | 11.3 | 171 | 18 | 10.5 | 173 | 8 | 4.6 | 156 | 8 | 5.1 | 0 | 0.0 | 140 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 338 | 14.7 | 219 | 14.2 | 212 | 6 | 2.8 | 212 | 0 | 0.0 | 188 | 30 | 16.0 | 1 | 0.5 | 161 | 1 | 0.6 | 0 | 0.0 |
| 45-54 | 288 | 12.6 | 185 | 12.0 | 181 | 1 | 0.6 | 183 | 4 | 2.2 | 162 | 23 | 14.2 | 1 | 0.6 | 152 | 2 | 1.3 | 1 | 0.7 |
| 55-64 | 83 | 3.6 | 48 | 3.1 | 44 | 3 | 6.8 | 44 | 0 | 0.0 | 45 | 15 | 33.3 | 1 | 2.2 | 38 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 9 | 0.4 | 4 | 0.3 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 4 | 1 | 25.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 2,292 | 100.0 | 1,538 | 100.0 | 1,500 | 181 | 12.1 | 1,510 | 74 | 4.9 | 1,324 | 99 | 7.5 | 5 | 0.4 | 1,214 | 4 | 0.3 | 1 | 0.1 |
| Others † | 3 | ~ | 2 | ~ | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 1 | 50.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 10.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Ruth Temple Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|-----------------------|--|-------|--------------------------|-------|---------------|------------|------------|---------------|------------|------------|---------------|------------|------------|----------------------------|-----|---------------|------------|------------|--------------|------|
| | Total STD Visits | | No. Visits with STD Test | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | n | % | Visits Tested | Positive n | Positive % | New HIV*** n | % |
| n | % | n | % | | | | | | | | | | | | | | | | | |
| Total | 5,471 | 100.0 | 3,569 | 100.0 | 3,507 | 443 | 12.6 | 3,523 | 229 | 6.5 | 3,158 | 210 | 6.6 | 14 | 0.4 | 2,890 | 28 | 1.0 | 16 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 81 | 2.6 | 55 | 2.7 | 55 | 3 | 5.5 | 55 | 3 | 5.5 | 52 | 1 | 1.9 | 0 | 0.0 | 48 | 1 | 2.1 | 1 | 2.1 |
| African American | 2,484 | 78.2 | 1,604 | 79.1 | 1,586 | 224 | 14.1 | 1,590 | 137 | 8.6 | 1,437 | 79 | 5.5 | 8 | 0.6 | 1,321 | 19 | 1.4 | 13 | 1.0 |
| Hispanic | 544 | 17.1 | 327 | 16.1 | 322 | 29 | 9.0 | 324 | 11 | 3.4 | 304 | 26 | 8.6 | 0 | 0.0 | 268 | 3 | 1.1 | 0 | 0.0 |
| Asian | 30 | 0.9 | 19 | 0.9 | 18 | 2 | 11.1 | 18 | 2 | 11.1 | 17 | 4 | 23.5 | 1 | 5.9 | 16 | 0 | 0.0 | 0 | 0.0 |
| Native American | 2 | 0.1 | 1 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 29 | 0.9 | 20 | 1.0 | 20 | 4 | 20.0 | 20 | 1 | 5.0 | 18 | 0 | 0.0 | 0 | 0.0 | 17 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 6 | 0.2 | 3 | 0.1 | 3 | 0 | 0.0 | 3 | 1 | 33.3 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 1 | 33.3 | 1 | 33.3 |
| Subtotal | 3,176 | 100.0 | 2,029 | 100.0 | 2,005 | 262 | 13.1 | 2,011 | 155 | 7.7 | 1,832 | 110 | 6.0 | 9 | 0.5 | 1,674 | 24 | 1.4 | 15 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 54 | 2.4 | 34 | 2.2 | 34 | 5 | 14.7 | 34 | 0 | 0.0 | 34 | 2 | 5.9 | 0 | 0.0 | 32 | 0 | 0.0 | 0 | 0.0 |
| African American | 1,828 | 79.8 | 1,258 | 81.8 | 1,229 | 145 | 11.8 | 1,238 | 68 | 5.5 | 1,076 | 66 | 6.1 | 4 | 0.4 | 989 | 2 | 0.2 | 0 | 0.0 |
| Hispanic | 336 | 14.7 | 202 | 13.1 | 195 | 27 | 13.8 | 195 | 3 | 1.5 | 181 | 26 | 14.4 | 0 | 0.0 | 162 | 2 | 1.2 | 1 | 0.6 |
| Asian | 29 | 1.3 | 16 | 1.0 | 16 | 1 | 6.3 | 16 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 | 12 | 0 | 0.0 | 0 | 0.0 |
| Native American | 2 | 0.1 | 1 | 0.1 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 43 | 1.9 | 27 | 1.8 | 25 | 3 | 12.0 | 26 | 3 | 11.5 | 20 | 5 | 25.0 | 1 | 5.0 | 18 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 2,292 | 100.0 | 1,538 | 100.0 | 1,500 | 181 | 12.1 | 1,510 | 74 | 4.9 | 1,324 | 99 | 7.5 | 5 | 0.4 | 1,214 | 4 | 0.3 | 1 | 0.1 |
| Others † | 3 | ~ | 2 | ~ | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 1 | 50.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 10.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Ruth Temple Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 10.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Ruth Temple Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

XI: Torrance Health Center

Table 11.1 Patient Visits and Tests, January - December 2010, Torrance Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-------------------|-----|-----|------|---|------|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | | HIV | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 793 | 100.0 | 613 | 100.0 | 580 | 66 | 11.4 | 583 | 31 | 5.3 | 579 | 37 | 6.4 | 2 | 0.3 | 526 | 8 | 1.5 | 6 | 1.1 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 555 | 70.0 | 426 | 69.5 | 410 | 48 | 11.7 | 412 | 29 | 7.0 | 403 | 19 | 4.7 | 2 | 0.5 | 368 | 8 | 2.2 | 6 | 1.6 |
| Female | 238 | 30.0 | 187 | 30.5 | 170 | 18 | 10.6 | 171 | 2 | 1.2 | 176 | 18 | 10.2 | 0 | 0.0 | 158 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | 0 | 0.0 | 0 | 0.0 | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | -- | 0 | 0 | -- | 0 | -- |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 79 | 10.0 | 59 | 9.6 | 54 | 9 | 16.7 | 54 | 12 | 22.2 | 57 | 12 | 21.1 | 2 | 3.5 | 47 | 8 | 17.0 | 6 | 12.8 |
| MSW | 444 | 56.0 | 346 | 56.4 | 336 | 36 | 10.7 | 338 | 15 | 4.4 | 325 | 6 | 1.8 | 0 | 0.0 | 303 | 0 | 0.0 | 0 | 0.0 |
| WSM | 181 | 22.8 | 150 | 24.5 | 136 | 14 | 10.3 | 137 | 1 | 0.7 | 140 | 13 | 9.3 | 0 | 0.0 | 124 | 0 | 0.0 | 0 | 0.0 |
| WSW/M | 21 | 2.6 | 15 | 2.4 | 15 | 2 | 13.3 | 15 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 68 | 8.6 | 43 | 7.0 | 39 | 5 | 12.8 | 39 | 3 | 7.7 | 43 | 6 | 14.0 | 0 | 0.0 | 38 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 2 | 0.3 | 2 | 0.3 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 45 | 5.7 | 38 | 6.2 | 38 | 11 | 28.9 | 38 | 5 | 13.2 | 36 | 0 | 0.0 | 0 | 0.0 | 36 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 175 | 22.1 | 142 | 23.2 | 135 | 21 | 15.6 | 136 | 11 | 8.1 | 133 | 7 | 5.3 | 1 | 0.8 | 122 | 4 | 3.3 | 3 | 2.5 |
| 25-29 | 198 | 25.0 | 149 | 24.3 | 143 | 9 | 6.3 | 145 | 8 | 5.5 | 140 | 5 | 3.6 | 1 | 0.7 | 132 | 1 | 0.8 | 1 | 0.8 |
| 30-34 | 102 | 12.9 | 84 | 13.7 | 80 | 13 | 16.3 | 80 | 1 | 1.3 | 77 | 3 | 3.9 | 0 | 0.0 | 71 | 1 | 1.4 | 0 | 0.0 |
| 35-44 | 124 | 15.6 | 98 | 16.0 | 90 | 5 | 5.6 | 90 | 4 | 4.4 | 94 | 6 | 6.4 | 0 | 0.0 | 79 | 1 | 1.3 | 1 | 1.3 |
| 45-54 | 84 | 10.6 | 62 | 10.1 | 60 | 6 | 10.0 | 60 | 2 | 3.3 | 59 | 5 | 8.5 | 0 | 0.0 | 54 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 39 | 4.9 | 24 | 3.9 | 22 | 0 | 0.0 | 22 | 0 | 0.0 | 24 | 7 | 29.2 | 0 | 0.0 | 19 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 24 | 3.0 | 14 | 2.3 | 10 | 1 | 10.0 | 10 | 0 | 0.0 | 14 | 4 | 28.6 | 0 | 0.0 | 11 | 1 | 9.1 | 1 | 9.1 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 160 | 20.2 | 127 | 20.7 | 123 | 7 | 5.7 | 124 | 6 | 4.8 | 123 | 3 | 2.4 | 0 | 0.0 | 118 | 1 | 0.8 | 1 | 0.8 |
| African American | 233 | 29.4 | 183 | 29.9 | 178 | 33 | 18.5 | 178 | 12 | 6.7 | 165 | 6 | 3.6 | 1 | 0.6 | 153 | 2 | 1.3 | 2 | 1.3 |
| Hispanic | 261 | 32.9 | 203 | 33.1 | 181 | 15 | 8.3 | 183 | 9 | 4.9 | 196 | 24 | 12.2 | 1 | 0.5 | 166 | 4 | 2.4 | 2 | 1.2 |
| Asian | 98 | 12.4 | 68 | 11.1 | 66 | 8 | 12.1 | 66 | 2 | 3.0 | 64 | 3 | 4.7 | 0 | 0.0 | 60 | 0 | 0.0 | 0 | 0.0 |
| Native American | 1 | 0.1 | 1 | 0.2 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 27 | 3.4 | 21 | 3.4 | 21 | 2 | 9.5 | 21 | 1 | 4.8 | 20 | 1 | 5.0 | 0 | 0.0 | 19 | 1 | 5.3 | 1 | 5.3 |
| Unknown | 13 | 1.6 | 10 | 1.6 | 10 | 1 | 10.0 | 10 | 1 | 10.0 | 10 | 0 | 0.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 566 | 71.4 | 546 | 89.1 | 541 | 62 | 11.5 | 542 | 30 | 5.5 | 531 | 19 | 3.6 | 1 | 0.2 | 508 | 7 | 1.4 | 6 | 1.2 |
| Return Visit | 227 | 28.6 | 67 | 10.9 | 39 | 4 | 10.3 | 41 | 1 | 2.4 | 48 | 18 | 37.5 | 1 | 2.1 | 18 | 1 | 5.6 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include both male-to-female (MTF) and female-to-male (FTM)

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested; similarly for RPR/TPPA, EIA/WB, etc.)

Table 11.2 Patient Visits and Tests by Gender and Age, January - December 2010, Torrance Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------|-------|----------------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|---------------|-----------------|----------------------------|-------------|---------------|-----------------|------|------------|------|
| | Total STD Visits n % | | No. Visits | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | New HIV*** | |
| | | | n | % | with STD Test n % | Visits Tested | Positive n % | n | % | Visits Tested | Positive n % | n | % | |
| Total | 793 | 100.0 | 613 | 100.0 | 580 | 66 | 11.4 | 583 | 31 | 5.3 | 579 | 37 | 6.4 | 2 | 0.3 | 526 | 8 | 1.5 | 6 | 1.1 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 1 | 0.2 | 1 | 0.2 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 23 | 4.1 | 20 | 4.7 | 20 | 6 | 30.0 | 20 | 4 | 20.0 | 19 | 0 | 0.0 | 0 | 0.0 | 19 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 118 | 21.3 | 95 | 22.3 | 92 | 15 | 16.3 | 92 | 10 | 10.9 | 90 | 6 | 6.7 | 1 | 1.1 | 82 | 4 | 4.9 | 3 | 3.7 |
| 25-29 | 148 | 26.7 | 110 | 25.8 | 106 | 6 | 5.7 | 108 | 8 | 7.4 | 103 | 3 | 2.9 | 1 | 1.0 | 98 | 1 | 1.0 | 1 | 1.0 |
| 30-34 | 85 | 15.3 | 69 | 16.2 | 65 | 12 | 18.5 | 65 | 1 | 1.5 | 63 | 3 | 4.8 | 0 | 0.0 | 57 | 1 | 1.8 | 0 | 0.0 |
| 35-44 | 80 | 14.4 | 62 | 14.6 | 60 | 4 | 6.7 | 60 | 4 | 6.7 | 60 | 1 | 1.7 | 0 | 0.0 | 51 | 1 | 2.0 | 1 | 2.0 |
| 45-54 | 58 | 10.5 | 43 | 10.1 | 43 | 4 | 9.3 | 43 | 2 | 4.7 | 41 | 2 | 4.9 | 0 | 0.0 | 39 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 26 | 4.7 | 18 | 4.2 | 16 | 0 | 0.0 | 16 | 0 | 0.0 | 18 | 3 | 16.7 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 16 | 2.9 | 8 | 1.9 | 7 | 1 | 14.3 | 7 | 0 | 0.0 | 8 | 1 | 12.5 | 0 | 0.0 | 7 | 1 | 14.3 | 1 | 14.3 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 555 | 100.0 | 426 | 100.0 | 410 | 48 | 11.7 | 412 | 29 | 7.0 | 403 | 19 | 4.7 | 2 | 0.5 | 368 | 8 | 2.2 | 6 | 1.6 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 1 | 0.4 | 1 | 0.5 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 22 | 9.2 | 18 | 9.6 | 18 | 5 | 27.8 | 18 | 1 | 5.6 | 17 | 0 | 0.0 | 0 | 0.0 | 17 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 57 | 23.9 | 47 | 25.1 | 43 | 6 | 14.0 | 44 | 1 | 2.3 | 43 | 1 | 2.3 | 0 | 0.0 | 40 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 50 | 21.0 | 39 | 20.9 | 37 | 3 | 8.1 | 37 | 0 | 0.0 | 37 | 2 | 5.4 | 0 | 0.0 | 34 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 17 | 7.1 | 15 | 8.0 | 15 | 1 | 6.7 | 15 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 | 14 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 44 | 18.5 | 36 | 19.3 | 30 | 1 | 3.3 | 30 | 0 | 0.0 | 34 | 5 | 14.7 | 0 | 0.0 | 28 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 26 | 10.9 | 19 | 10.2 | 17 | 2 | 11.8 | 17 | 0 | 0.0 | 18 | 3 | 16.7 | 0 | 0.0 | 15 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 13 | 5.5 | 6 | 3.2 | 6 | 0 | 0.0 | 6 | 0 | 0.0 | 6 | 4 | 66.7 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 8 | 3.4 | 6 | 3.2 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 6 | 3 | 50.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 238 | 100.0 | 187 | 100.0 | 170 | 18 | 10.6 | 171 | 2 | 1.2 | 176 | 18 | 10.2 | 0 | 0.0 | 158 | 0 | 0.0 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 11.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Torrance Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-----|---|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 793 | 100.0 | 613 | 100.0 | 580 | 66 | 11.4 | 583 | 31 | 5.3 | 579 | 37 | 6.4 | 2 | 0.3 | 526 | 8 | 1.5 | 6 | 1.1 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 131 | 23.6 | 105 | 24.6 | 104 | 4 | 3.8 | 104 | 4 | 3.8 | 103 | 1 | 1.0 | 0 | 0.0 | 99 | 1 | 1.0 | 1 | 1.0 |
| African American | 157 | 28.3 | 123 | 28.9 | 119 | 25 | 21.0 | 119 | 12 | 10.1 | 111 | 4 | 3.6 | 1 | 0.9 | 101 | 2 | 2.0 | 2 | 2.0 |
| Hispanic | 175 | 31.5 | 133 | 31.2 | 122 | 11 | 9.0 | 124 | 9 | 7.3 | 128 | 12 | 9.4 | 1 | 0.8 | 111 | 4 | 3.6 | 2 | 1.8 |
| Asian | 62 | 11.2 | 43 | 10.1 | 43 | 6 | 14.0 | 43 | 2 | 4.7 | 39 | 1 | 2.6 | 0 | 0.0 | 37 | 0 | 0.0 | 0 | 0.0 |
| Native American | 1 | 0.2 | 1 | 0.2 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 18 | 3.2 | 13 | 3.1 | 13 | 1 | 7.7 | 13 | 1 | 7.7 | 13 | 1 | 7.7 | 0 | 0.0 | 12 | 1 | 8.3 | 1 | 8.3 |
| Unknown | 11 | 2.0 | 8 | 1.9 | 8 | 1 | 12.5 | 8 | 1 | 12.5 | 8 | 0 | 0.0 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 555 | 100.0 | 426 | 100.0 | 410 | 48 | 11.7 | 412 | 29 | 7.0 | 403 | 19 | 4.7 | 2 | 0.5 | 368 | 8 | 2.2 | 6 | 1.6 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 29 | 12.2 | 22 | 11.8 | 19 | 3 | 15.8 | 20 | 2 | 10.0 | 20 | 2 | 10.0 | 0 | 0.0 | 19 | 0 | 0.0 | 0 | 0.0 |
| African American | 76 | 31.9 | 60 | 32.1 | 59 | 8 | 13.6 | 59 | 0 | 0.0 | 54 | 2 | 3.7 | 0 | 0.0 | 52 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 86 | 36.1 | 70 | 37.4 | 59 | 4 | 6.8 | 59 | 0 | 0.0 | 68 | 12 | 17.6 | 0 | 0.0 | 55 | 0 | 0.0 | 0 | 0.0 |
| Asian | 36 | 15.1 | 25 | 13.4 | 23 | 2 | 8.7 | 23 | 0 | 0.0 | 25 | 2 | 8.0 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 |
| Native American | 9 | 3.8 | 8 | 4.3 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Other | 2 | 0.8 | 2 | 1.1 | 8 | 1 | 12.5 | 8 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 238 | 100.0 | 187 | 100.0 | 170 | 18 | 10.6 | 171 | 2 | 1.2 | 176 | 18 | 10.2 | 0 | 0.0 | 158 | 0 | 0.0 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 11.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Torrance Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 11.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Torrance Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

XII: Curtis Tucker Health Center

Table 12.1 Patient Visits and Tests, January - December 2010, Curtis Tucker Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-------------------|-------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 5,608 | 100.0 | 3,554 | 100.0 | 3,399 | 346 | 10.2 | 3,431 | 180 | 5.2 | 3,091 | 121 | 3.9 | 10 | 0.3 | 2,879 | 19 | 0.7 | 17 | 0.6 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 3,378 | 60.2 | 2,097 | 59.0 | 2,027 | 228 | 11.2 | 2,039 | 134 | 6.6 | 1,870 | 83 | 4.4 | 10 | 0.5 | 1,733 | 19 | 1.1 | 17 | 1.0 |
| Female | 2,229 | 39.7 | 1,456 | 41.0 | 1,371 | 118 | 8.6 | 1,391 | 46 | 3.3 | 1,220 | 38 | 3.1 | 0 | 0.0 | 1,145 | 0 | 0.0 | 0 | 0.0 |
| Transgender† | 1 | 0.0 | 1 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 329 | 5.9 | 202 | 5.7 | 188 | 18 | 9.6 | 189 | 33 | 17.5 | 180 | 37 | 20.6 | 5 | 2.8 | 150 | 12 | 8.0 | 10 | 6.7 |
| MSW | 2,942 | 52.5 | 1,847 | 52.0 | 1,795 | 207 | 11.5 | 1,805 | 98 | 5.4 | 1,644 | 41 | 2.5 | 4 | 0.2 | 1543 | 5 | 0.3 | 5 | 0.3 |
| WSM | 1,951 | 34.8 | 1,300 | 36.6 | 1,229 | 106 | 8.6 | 1,246 | 40 | 3.2 | 1,092 | 32 | 2.9 | 0 | 0.0 | 1029 | 0 | 0.0 | 0 | 0.0 |
| WSW/M | 189 | 3.4 | 117 | 3.3 | 109 | 6 | 5.5 | 112 | 6 | 5.4 | 95 | 3 | 3.2 | 0 | 0.0 | 90 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 197 | 3.5 | 88 | 2.5 | 78 | 9 | 11.5 | 79 | 3 | 3.8 | 80 | 8 | 10.0 | 1 | 1.3 | 67 | 2 | 3.0 | 2 | 3.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 18 | 0.3 | 10 | 0.3 | 10 | 6 | 60.0 | 10 | 1 | 10.0 | 8 | 0 | 0.0 | 0 | 0.0 | 8 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 604 | 10.8 | 397 | 11.2 | 377 | 85 | 22.5 | 380 | 43 | 11.3 | 343 | 5 | 1.5 | 2 | 0.6 | 329 | 1 | 0.3 | 1 | 0.3 |
| 20-24 | 1,300 | 23.2 | 852 | 24.0 | 820 | 126 | 15.4 | 830 | 51 | 6.1 | 732 | 17 | 2.3 | 1 | 0.1 | 699 | 7 | 1.0 | 5 | 0.7 |
| 25-29 | 1,052 | 18.8 | 682 | 19.2 | 663 | 65 | 9.8 | 667 | 33 | 4.9 | 596 | 17 | 2.9 | 2 | 0.3 | 558 | 2 | 0.4 | 2 | 0.4 |
| 30-34 | 758 | 13.5 | 480 | 13.5 | 459 | 34 | 7.4 | 466 | 20 | 4.3 | 428 | 12 | 2.8 | 2 | 0.5 | 397 | 4 | 1.0 | 4 | 1.0 |
| 35-44 | 959 | 17.1 | 590 | 16.6 | 566 | 23 | 4.1 | 567 | 20 | 3.5 | 508 | 25 | 4.9 | 0 | 0.0 | 462 | 3 | 0.6 | 3 | 0.6 |
| 45-54 | 689 | 12.3 | 417 | 11.7 | 392 | 5 | 1.3 | 398 | 9 | 2.3 | 365 | 30 | 8.2 | 3 | 0.8 | 331 | 2 | 0.6 | 2 | 0.6 |
| 55-64 | 184 | 3.3 | 102 | 2.9 | 90 | 1 | 1.1 | 91 | 2 | 2.2 | 91 | 9 | 9.9 | 0 | 0.0 | 78 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 44 | 0.8 | 24 | 0.7 | 22 | 1 | 4.5 | 22 | 1 | 4.5 | 20 | 6 | 30.0 | 0 | 0.0 | 17 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 301 | 5.4 | 184 | 5.2 | 173 | 3 | 1.7 | 175 | 10 | 5.7 | 165 | 4 | 2.4 | 0 | 0.0 | 153 | 0 | 0.0 | 0 | 0.0 |
| African American | 3,742 | 66.7 | 2,485 | 69.9 | 2,400 | 273 | 11.4 | 2,427 | 147 | 6.1 | 2,131 | 63 | 3.0 | 7 | 0.3 | 1,992 | 16 | 0.8 | 14 | 0.7 |
| Hispanic | 1,250 | 22.3 | 684 | 19.2 | 636 | 49 | 7.7 | 639 | 15 | 2.3 | 624 | 46 | 7.4 | 3 | 0.5 | 571 | 3 | 0.5 | 3 | 0.5 |
| Asian | 111 | 2.0 | 70 | 2.0 | 67 | 7 | 10.4 | 67 | 2 | 3.0 | 64 | 1 | 1.6 | 0 | 0.0 | 60 | 0 | 0.0 | 0 | 0.0 |
| Native American | 14 | 0.2 | 11 | 0.3 | 10 | 0 | 0.0 | 10 | 1 | 10.0 | 9 | 2 | 22.2 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Other | 123 | 2.2 | 80 | 2.3 | 73 | 6 | 8.2 | 73 | 4 | 5.5 | 63 | 4 | 6.3 | 0 | 0.0 | 59 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 67 | 1.2 | 40 | 1.1 | 40 | 8 | 20.0 | 40 | 1 | 2.5 | 35 | 1 | 2.9 | 0 | 0.0 | 35 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 3,758 | 67.0 | 3,408 | 95.9 | 3,303 | 336 | 10.2 | 3,329 | 173 | 5.2 | 3,020 | 106 | 3.5 | 8 | 0.3 | 2,839 | 19 | 0.7 | 17 | 0.6 |
| Return Visit | 1,850 | 33.0 | 146 | 4.1 | 96 | 10 | 10.4 | 102 | 7 | 6.9 | 71 | 15 | 21.1 | 2 | 2.8 | 40 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include 1 male-to-female (MTF)

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 12.2 Patient Visits and Tests by Gender and Age, January - December 2010, Curtis Tucker Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------|-------|----------------------|---------------|--------------|---------------|--------------|---------------|--------------|-----|------|----------------------------|--------------|-------------------|----|-----|----|-----|
| | Total STD Visits n % | | No. Visits | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | n | % | with STD Test n % | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n | % | Visits Tested | Positive n % | New HIV*** n % | | | | |
| Total | 5,608 | 100.0 | 3,554 | 100.0 | 3,399 | 346 | 10.2 | 3,431 | 180 | 5.2 | 3,091 | 121 | 3.9 | 10 | 0.3 | 2,879 | 19 | 0.7 | 17 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 7 | 0.2 | 3 | 0.1 | 3 | 2.0 | 66.7 | 3 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 264 | 7.8 | 174 | 8.3 | 170 | 39 | 22.9 | 171 | 26 | 15.2 | 159 | 5 | 3.1 | 2 | 1.3 | 153 | 1 | 0.7 | 1 | 0.7 |
| 20-24 | 741 | 21.9 | 478 | 22.8 | 463 | 90 | 19.4 | 469 | 37 | 7.9 | 420 | 12 | 2.9 | 1 | 0.2 | 401 | 7 | 1.7 | 5 | 1.2 |
| 25-29 | 678 | 20.1 | 430 | 20.5 | 423 | 39 | 9.2 | 423 | 26 | 6.1 | 381 | 13 | 3.4 | 2 | 0.5 | 359 | 2 | 0.6 | 2 | 0.6 |
| 30-34 | 472 | 14.0 | 299 | 14.3 | 289 | 30 | 10.4 | 291 | 13 | 4.5 | 269 | 8 | 3.0 | 2 | 0.7 | 247 | 4 | 1.6 | 4 | 1.6 |
| 35-44 | 633 | 18.7 | 377 | 18.0 | 367 | 23 | 6.3 | 368 | 20 | 5.4 | 338 | 14 | 4.1 | 0 | 0.0 | 307 | 3 | 1.0 | 3 | 1.0 |
| 45-54 | 436 | 12.9 | 253 | 12.1 | 241 | 3 | 1.2 | 242 | 9 | 3.7 | 227 | 24 | 10.6 | 3 | 1.3 | 200 | 2 | 1.0 | 2 | 1.0 |
| 55-64 | 122 | 3.6 | 70 | 3.3 | 60 | 1 | 1.7 | 61 | 2 | 3.3 | 62 | 4 | 6.5 | 0 | 0.0 | 53 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 26 | 0.8 | 14 | 0.7 | 12 | 1 | 8.3 | 12 | 1 | 8.3 | 12 | 3 | 25.0 | 0 | 0.0 | 11 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,379 | 100.0 | 2,098 | 100.0 | 2,028 | 228 | 11.2 | 2,040 | 134 | 6.6 | 1,871 | 83 | 4.4 | 10 | 0.5 | 1,734 | 19 | 1.1 | 17 | 1.0 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 11 | 0.5 | 7 | 0.5 | 7 | 4 | 57.1 | 7 | 1 | 14.3 | 5 | 0 | 0.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 340 | 15.3 | 223 | 15.3 | 207 | 46 | 22.2 | 209 | 17 | 8.1 | 184 | 0 | 0.0 | 0 | 0.0 | 176 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 559 | 25.1 | 374 | 25.7 | 357 | 36 | 10.1 | 361 | 14 | 3.9 | 312 | 5 | 1.6 | 0 | 0.0 | 298 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 374 | 16.8 | 252 | 17.3 | 240 | 26 | 10.8 | 244 | 7 | 2.9 | 215 | 4 | 1.9 | 0 | 0.0 | 199 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 286 | 12.8 | 181 | 12.4 | 170 | 4 | 2.4 | 175 | 7 | 4.0 | 159 | 4 | 2.5 | 0 | 0.0 | 150 | 0 | 0.0 | 0 | 0.0 |
| 35-44 | 326 | 14.6 | 213 | 14.6 | 199 | 0 | 0.0 | 199 | 0 | 0.0 | 170 | 11 | 6.5 | 0 | 0.0 | 155 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 253 | 11.4 | 164 | 11.3 | 151 | 2 | 1.3 | 156 | 0 | 0.0 | 138 | 6 | 4.3 | 0 | 0.0 | 131 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 62 | 2.8 | 32 | 2.2 | 30 | 0 | 0.0 | 30 | 0 | 0.0 | 29 | 5 | 17.2 | 0 | 0.0 | 25 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 18 | 0.8 | 10 | 0.7 | 10 | 0 | 0.0 | 10 | 0 | 0.0 | 8 | 3 | 37.5 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 2,229 | 100.0 | 1,456 | 100.0 | 1,371 | 118 | 8.6 | 1,391 | 46 | 3.3 | 1,220 | 38 | 3.1 | 0 | 0.0 | 1,145 | 0 | 0.0 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 17 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 12.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Curtis Tucker Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | | HIV Testing | | | | |
|-----------------------|--|-------|--------------------------|-------|---------------|------------|------------|---------------|------------|------------|---------------|------------|------------|----------------------------|-----|---------------|------------|------------|--------------|-------|
| | Total STD Visits | | No. Visits with STD Test | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | Visits Tested | Positive n | Positive % | n | % | Visits Tested | Positive n | Positive % | New HIV*** n | *** % |
| n | % | n | % | | | | | | | | | | | | | | | | | |
| Total | 5,608 | 100.0 | 3,554 | 100.0 | 3,399 | 346 | 10.2 | 3,431 | 180 | 5.2 | 3,091 | 121 | 3.9 | 10 | 0.3 | 2,879 | 19 | 0.7 | 17 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 212 | 6.3 | 126 | 6.0 | 119 | 2 | 1.7 | 120 | 6 | 5.0 | 117 | 3 | 2.6 | 0 | 0.0 | 108 | 0 | 0.0 | 0 | 0.0 |
| African American | 2,129 | 63.0 | 1,402 | 66.8 | 1,369 | 179 | 13.1 | 1,378 | 113 | 8.2 | 1,233 | 42 | 3.4 | 7 | 0.6 | 1,144 | 16 | 1.4 | 14 | 1.2 |
| Hispanic | 858 | 25.4 | 458 | 21.8 | 430 | 39 | 9.1 | 432 | 11 | 2.5 | 421 | 32 | 7.6 | 3 | 0.7 | 385 | 3 | 0.8 | 3 | 0.8 |
| Asian | 65 | 1.9 | 38 | 1.8 | 37 | 2 | 5.4 | 37 | 1 | 2.7 | 38 | 1 | 2.6 | 0 | 0.0 | 36 | 0 | 0.0 | 0 | 0.0 |
| Native American | 6 | 0.2 | 6 | 0.3 | 6 | 0 | 0.0 | 6 | 1 | 16.7 | 4 | 2 | 50.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Other | 58 | 1.7 | 39 | 1.9 | 38 | 1 | 2.6 | 38 | 2 | 5.3 | 32 | 2 | 6.3 | 0 | 0.0 | 31 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 51 | 1.5 | 29 | 1.4 | 29 | 5 | 17.2 | 29 | 0 | 0.0 | 26 | 1 | 3.8 | 0 | 0.0 | 26 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 3,379 | 100.0 | 2,098 | 100.0 | 2,028 | 228 | 11.2 | 2,040 | 134 | 6.6 | 1,871 | 83 | 4.4 | 10 | 0.5 | 1,734 | 19 | 1.1 | 17 | 1.0 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 89 | 4.0 | 58 | 4.0 | 54 | 1 | 1.9 | 55 | 4 | 7.3 | 48 | 1 | 2.1 | 0 | 0.0 | 45 | 0 | 0.0 | 0 | 0.0 |
| African American | 1,613 | 72.4 | 1,083 | 74.4 | 1,031 | 94 | 9.1 | 1,049 | 34 | 3.2 | 898 | 21 | 2.3 | 0 | 0.0 | 848 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 392 | 17.6 | 226 | 15.5 | 206 | 10 | 4.9 | 207 | 4 | 1.9 | 203 | 14 | 6.9 | 0 | 0.0 | 186 | 0 | 0.0 | 0 | 0.0 |
| Asian | 46 | 2.1 | 32 | 2.2 | 30 | 5 | 16.7 | 30 | 1 | 3.3 | 26 | 0 | 0.0 | 0 | 0.0 | 24 | 0 | 0.0 | 0 | 0.0 |
| Native American | 8 | 0.4 | 5 | 0.3 | 4 | 0 | 0.0 | 4 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Other | 65 | 2.9 | 41 | 2.8 | 35 | 5 | 14.3 | 35 | 2 | 5.7 | 31 | 2 | 6.5 | 0 | 0.0 | 28 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 16 | 0.7 | 11 | 0.8 | 11 | 3 | 27.3 | 11 | 1 | 9.1 | 9 | 0 | 0.0 | 0 | 0.0 | 9 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 2,229 | 100.0 | 1,456 | 100.0 | 1,371 | 118 | 8.6 | 1,391 | 46 | 3.3 | 1,220 | 38 | 3.1 | 0 | 0.0 | 1,145 | 0 | 0.0 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 12.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Curtis Tucker Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 12.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Curtis Tucker Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

XIII: Whittier Health Center

Table 13.1 Patient Visits and Tests, January - December 2010, Whittier Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|--------------------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|------------------|----------|-----|------------------|-------------------------------|-------------------|-------|----|-----|---|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive* n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 2,164 | 100.0 | 1,333 | 100.0 | 1,268 | 154 | 12.1 | 1,276 | 41 | 3.2 | 1,228 | 95 | 7.7 | 8 | 0.7 | 1,179 | 12 | 1.0 | 7 | 0.6 |
| Gender | | | | | | | | | | | | | | | | | | | | |
| Male | 1,501 | 69.4 | 909 | 68.2 | 866 | 96 | 11.1 | 872 | 31 | 3.6 | 843 | 70 | 8.3 | 8 | 0.9 | 813 | 11 | 1.4 | 7 | 0.9 |
| Female | 663 | 30.6 | 424 | 31.8 | 402 | 58 | 14.4 | 404 | 10 | 2.5 | 385 | 25 | 6.5 | 0 | 0.0 | 366 | 1 | 0.3 | 0 | 0.0 |
| Transgender† | 0 | 0.0 | 0 | 0.0 | 0 | 0 | -- | 0 | 0 | -- | 0 | 0 | -- | 0 | -- | 0 | 0 | -- | 0 | -- |
| Gender of Sex Partner†† | | | | | | | | | | | | | | | | | | | | |
| MSM/W | 290 | 13.4 | 177 | 13.3 | 168 | 18 | 10.7 | 169 | 22 | 13.0 | 163 | 46 | 28.2 | 6 | 3.7 | 145 | 11 | 7.6 | 7 | 4.8 |
| MSW | 1,114 | 51.5 | 699 | 52.4 | 665 | 76 | 11.4 | 670 | 9 | 1.3 | 647 | 19 | 2.9 | 2 | 0.3 | 636 | 0 | 0.0 | 0 | 0.0 |
| WSM | 578 | 26.7 | 386 | 29.0 | 366 | 54 | 14.8 | 368 | 9 | 2.4 | 350 | 23 | 6.6 | 0 | 0.0 | 333 | 1 | 0.3 | 0 | 0.0 |
| WSW/M | 30 | 1.4 | 22 | 1.7 | 22 | 2 | 9.1 | 22 | 0 | 0.0 | 19 | 0 | 0.0 | 0 | 0.0 | 18 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 152 | 7.0 | 49 | 3.7 | 47 | 4 | 8.5 | 47 | 1 | 2.1 | 49 | 7 | 14.3 | 0 | 0.0 | 47 | 0 | 0.0 | 0 | 0.0 |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 6 | 0.3 | 3 | 0.2 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 197 | 9.1 | 145 | 10.9 | 143 | 33 | 23.1 | 145 | 6 | 4.1 | 128 | 6 | 4.7 | 0 | 0.0 | 124 | 1 | 0.8 | 1 | 0.8 |
| 20-24 | 541 | 25.0 | 346 | 26.0 | 339 | 65 | 19.2 | 339 | 12 | 3.5 | 320 | 11 | 3.4 | 2 | 0.6 | 313 | 2 | 0.6 | 2 | 0.6 |
| 25-29 | 451 | 20.8 | 269 | 20.2 | 256 | 32 | 12.5 | 259 | 12 | 4.6 | 252 | 17 | 6.7 | 2 | 0.8 | 246 | 4 | 1.6 | 1 | 0.4 |
| 30-34 | 292 | 13.5 | 179 | 13.4 | 172 | 10 | 5.8 | 174 | 6 | 3.4 | 164 | 16 | 9.8 | 1 | 0.6 | 156 | 3 | 1.9 | 1 | 0.6 |
| 35-44 | 366 | 16.9 | 221 | 16.6 | 209 | 9 | 4.3 | 209 | 4 | 1.9 | 209 | 21 | 10.0 | 3 | 1.4 | 198 | 2 | 1.0 | 2 | 1.0 |
| 45-54 | 225 | 10.4 | 121 | 9.1 | 106 | 3 | 2.8 | 107 | 0 | 0.0 | 110 | 14 | 12.7 | 0 | 0.0 | 103 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 65 | 3.0 | 39 | 2.9 | 33 | 2 | 6.1 | 33 | 0 | 0.0 | 34 | 7 | 20.6 | 0 | 0.0 | 29 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 21 | 1.0 | 10 | 0.8 | 7 | 0 | 0.0 | 7 | 1 | 14.3 | 9 | 3 | 33.3 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 250 | 11.6 | 151 | 11.3 | 145 | 8 | 5.5 | 146 | 5 | 3.4 | 137 | 6 | 4.4 | 0 | 0.0 | 132 | 0 | 0.0 | 0 | 0.0 |
| African American | 90 | 4.2 | 61 | 4.6 | 58 | 10 | 17.2 | 58 | 1 | 1.7 | 55 | 4 | 7.3 | 0 | 0.0 | 53 | 3 | 5.7 | 1 | 1.9 |
| Hispanic | 1,648 | 76.2 | 1,001 | 75.1 | 948 | 129 | 13.6 | 954 | 34 | 3.6 | 928 | 80 | 8.6 | 7 | 0.8 | 890 | 9 | 1.0 | 6 | 0.7 |
| Asian | 113 | 5.2 | 77 | 5.8 | 74 | 2 | 2.7 | 75 | 0 | 0.0 | 68 | 2 | 2.9 | 1 | 1.5 | 64 | 0 | 0.0 | 0 | 0.0 |
| Native American | 10 | 0.5 | 7 | 0.5 | 7 | 0 | 0.0 | 7 | 1 | 14.3 | 7 | 2 | 28.6 | 0 | 0.0 | 7 | 0 | 0.0 | 0 | 0.0 |
| Other | 44 | 2.0 | 31 | 2.3 | 31 | 4 | 12.9 | 31 | 0 | 0.0 | 28 | 1 | 3.6 | 0 | 0.0 | 28 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 9 | 0.4 | 5 | 0.4 | 5 | 1 | 20.0 | 5 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Type of Visit | | | | | | | | | | | | | | | | | | | | |
| New Visit | 1,468 | 67.8 | 1,226 | 92.0 | 1,201 | 147 | 12.2 | 1,207 | 37 | 3.1 | 1,161 | 83 | 7.1 | 8 | 0.7 | 1,135 | 11 | 1.0 | 6 | 0.5 |
| Return Visit | 696 | 32.2 | 107 | 8.0 | 67 | 7 | 10.4 | 69 | 4 | 5.8 | 67 | 12 | 17.9 | 0 | 0.0 | 44 | 1 | 2.3 | 1 | 2.3 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Transgendered in total visits include both male-to-female (MTF) and female-to-male (FTM)

††Based on self-reported gender of sexual partners; "MSM/W"=men who have sex with men, or men who have sex with men and women; "MSW"=men who have sex with women; "WSM"=women who have sex with men; "WSW/M"=women who have sex with women, or women who have sex with men and women

Note: "Visits Tested" for chlamydia, gonorrhea, syphilis and HIV refers to number of individual visits and not individual tests (ie. a patient who is tested for urethral and rectal chlamydia on the same visit is only counted as one visit tested: similarly for RPR/TPPA, EIA/WB, etc.)

Table 13.2 Patient Visits and Tests by Gender and Age, January - December 2010, Whittier Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------|--|-------|------------------------------------|-------|------------------|---------------|------|------------------|---------------|------|------------------|---------------|------|--------------------------------------|-------------|------------------|---------------|-----|-------------------|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** n % | | HIV | | | | |
| | | | | | Visits Tested | Positive n | % | Visits Tested | Positive n | % | Visits Tested | Positive n | % | | | Visits Tested | Positive n | % | New HIV*** n % | |
| Total | 2,164 | 100.0 | 1,333 | 100.0 | 1,268 | 154 | 12.1 | 1,276 | 41 | 3.2 | 1,229 | 95 | 7.7 | 8 | 0.7 | 1,179 | 12 | 1.0 | 7 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 4 | 0.3 | 2 | 0.2 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 106 | 7.1 | 77 | 8.5 | 75 | 12 | 16.0 | 77 | 3 | 3.9 | 69 | 4 | 5.8 | 0 | 0.0 | 67 | 1 | 1.5 | 1 | 1.5 |
| 20-24 | 384 | 25.6 | 238 | 26.2 | 232 | 40 | 17.2 | 232 | 9 | 3.9 | 223 | 8 | 3.6 | 2 | 0.9 | 219 | 2 | 0.9 | 2 | 0.9 |
| 25-29 | 351 | 23.4 | 207 | 22.8 | 199 | 27 | 13.6 | 201 | 9 | 4.5 | 193 | 13 | 6.7 | 2 | 1.0 | 190 | 4 | 2.1 | 4 | 2.1 |
| 30-34 | 205 | 13.7 | 128 | 14.1 | 124 | 8 | 6.5 | 125 | 5 | 4.0 | 118 | 10 | 8.5 | 1 | 0.8 | 113 | 2 | 1.8 | 2 | 1.8 |
| 35-44 | 272 | 18.1 | 150 | 16.5 | 142 | 7 | 4.9 | 142 | 4 | 2.8 | 142 | 19 | 13.4 | 3 | 2.1 | 134 | 2 | 1.5 | 2 | 1.5 |
| 45-54 | 135 | 9.0 | 78 | 8.6 | 67 | 2 | 3.0 | 68 | 0 | 0.0 | 72 | 9 | 12.5 | 0 | 0.0 | 66 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 35 | 2.3 | 24 | 2.6 | 20 | 0 | 0.0 | 20 | 0 | 0.0 | 21 | 6 | 28.6 | 0 | 0.0 | 18 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 9 | 0.6 | 5 | 0.6 | 5 | 0 | 0.0 | 5 | 1 | 20.0 | 4 | 1 | 25.0 | 0 | 0.0 | 4 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,501 | 100.0 | 909 | 100.0 | 866 | 96 | 11.1 | 872 | 31 | 3.6 | 844 | 70 | 8.3 | 8 | 0.9 | 813 | 11 | 1.4 | 7 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Age | | | | | | | | | | | | | | | | | | | | |
| 0-14 | 2 | 0.3 | 1 | 0.2 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| 15-19 | 91 | 13.7 | 68 | 16.0 | 68 | 21 | 30.9 | 68 | 3 | 4.4 | 59 | 2 | 3.4 | 0 | 0.0 | 57 | 0 | 0.0 | 0 | 0.0 |
| 20-24 | 157 | 23.7 | 108 | 25.5 | 107 | 25 | 23.4 | 107 | 3 | 2.8 | 97 | 3 | 3.1 | 0 | 0.0 | 94 | 0 | 0.0 | 0 | 0.0 |
| 25-29 | 100 | 15.1 | 62 | 14.6 | 57 | 5 | 8.8 | 58 | 3 | 5.2 | 59 | 4 | 6.8 | 0 | 0.0 | 56 | 0 | 0.0 | 0 | 0.0 |
| 30-34 | 87 | 13.1 | 51 | 12.0 | 48 | 2 | 4.2 | 49 | 1 | 2.0 | 46 | 6 | 13.0 | 0 | 0.0 | 43 | 1 | 2.3 | 0 | 0.0 |
| 35-44 | 94 | 14.2 | 71 | 16.7 | 67 | 2 | 3.0 | 67 | 0 | 0.0 | 67 | 2 | 3.0 | 0 | 0.0 | 64 | 0 | 0.0 | 0 | 0.0 |
| 45-54 | 90 | 13.6 | 43 | 10.1 | 39 | 1 | 2.6 | 39 | 0 | 0.0 | 38 | 5 | 13.2 | 0 | 0.0 | 37 | 0 | 0.0 | 0 | 0.0 |
| 55-64 | 30 | 4.5 | 15 | 3.5 | 13 | 2 | 15.4 | 13 | 0 | 0.0 | 13 | 1 | 7.7 | 0 | 0.0 | 11 | 0 | 0.0 | 0 | 0.0 |
| 65+ | 12 | 1.8 | 5 | 1.2 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 5 | 2 | 40.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 663 | 100.0 | 424 | 100.0 | 402 | 58 | 14.4 | 404 | 10 | 2.5 | 385 | 25 | 6.5 | 0 | 0.0 | 366 | 1 | 0.3 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 | 0 | 0 | 0.0 | 0 | 0.0 |

*Screening plus confirmatory test

**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Table 13.3 Patient Visits and Tests by Gender and Race/Ethnicity, January - December 2010, Whittier Health Center

| | Chlamydia, Gonorrhea, and Syphilis Testing | | | | | | | | | | | | | | HIV Testing | | | | | |
|-----------------------|--|-------|------------------------------------|-------|------------------|-----------------|------------------|-----------------|------------------|-----------------|----------|-----|------------------|-------------------------------|-------------------|-------|-----|-----|-----|-----|
| | Total STD Visits n % | | No. Visits with STD Test n % | | Chlamydia | | | Gonorrhea | | | Syphilis | | | Confirmed Early Syphilis** | | HIV | | | | |
| | | | | | Visits Tested | Positive n % | Visits Tested | Positive n % | Visits Tested | Positive n % | n % | n % | Visits Tested | Positive n % | New HIV*** n % | | | | | |
| Total | 2,164 | 100.0 | 1,333 | 100.0 | 1,268 | 154 | 12.1 | 1,276 | 41 | 3.2 | 1,229 | 95 | 7.7 | 8 | 0.7 | 1,179 | 12 | 1.0 | 7 | 0.6 |
| Male | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 158 | 10.5 | 98 | 10.8 | 94 | 6 | 6.4 | 95 | 4 | 4.2 | 90 | 6 | 6.7 | 0 | 0.0 | 86 | 0 | 0.0 | 0 | 0.0 |
| African American | 66 | 4.4 | 43 | 4.7 | 41 | 7 | 17.1 | 41 | 1 | 2.4 | 39 | 4 | 10.3 | 0 | 0.0 | 37 | 3 | 8.1 | 1 | 2.7 |
| Hispanic | 1,148 | 76.5 | 684 | 75.2 | 647 | 79 | 12.2 | 652 | 25 | 3.8 | 639 | 56 | 8.8 | 7 | 1.1 | 616 | 8 | 1.3 | 6 | 1.0 |
| Asian | 76 | 5.1 | 51 | 5.6 | 51 | 1 | 2.0 | 51 | 0 | 0.0 | 43 | 1 | 2.3 | 1 | 2.3 | 42 | 0 | 0.0 | 0 | 0.0 |
| Native American | 9 | 0.6 | 6 | 0.7 | 6 | 0 | 0.0 | 6 | 1 | 16.7 | 6 | 2 | 33.3 | 0 | 0.0 | 6 | 0 | 0.0 | 0 | 0.0 |
| Other | 37 | 2.5 | 24 | 2.6 | 24 | 2 | 8.3 | 24 | 0 | 0.0 | 24 | 1 | 4.2 | 0 | 0.0 | 23 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 7 | 0.5 | 3 | 0.3 | 3 | 1 | 33.3 | 3 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 | 3 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 1,501 | 100.0 | 909 | 100.0 | 866 | 96 | 11.1 | 872 | 31 | 3.6 | 844 | 70 | 8.3 | 8 | 0.9 | 813 | 11 | 1.4 | 7 | 0.9 |
| Female | | | | | | | | | | | | | | | | | | | | |
| Race/Ethnicity | | | | | | | | | | | | | | | | | | | | |
| White | 92 | 13.9 | 53 | 12.5 | 51 | 2 | 3.9 | 51 | 1 | 2.0 | 47 | 0 | 0.0 | 0 | 0.0 | 46 | 0 | 0.0 | 0 | 0.0 |
| African American | 24 | 3.6 | 18 | 4.2 | 17 | 3 | 17.6 | 17 | 0 | 0.0 | 16 | 0 | 0.0 | 0 | 0.0 | 16 | 0 | 0.0 | 0 | 0.0 |
| Hispanic | 500 | 75.4 | 317 | 74.8 | 301 | 50 | 16.6 | 302 | 9 | 3.0 | 289 | 24 | 8.3 | 0 | 0.0 | 274 | 1 | 0.4 | 0 | 0.0 |
| Asian | 37 | 5.6 | 26 | 6.1 | 23 | 1 | 4.3 | 24 | 0 | 0.0 | 25 | 1 | 4.0 | 0 | 0.0 | 22 | 0 | 0.0 | 0 | 0.0 |
| Native American | 1 | 0.2 | 1 | 0.2 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 | 1 | 0 | 0.0 | 0 | 0.0 |
| Other | 7 | 1.1 | 7 | 1.7 | 7 | 2 | 28.6 | 7 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 | 5 | 0 | 0.0 | 0 | 0.0 |
| Unknown | 2 | 0.3 | 2 | 0.5 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 | 2 | 0 | 0.0 | 0 | 0.0 |
| Subtotal | 663 | 100.0 | 424 | 100.0 | 402 | 58 | 14.4 | 404 | 10 | 2.5 | 385 | 25 | 6.5 | 0 | 0.0 | 366 | 1 | 0.3 | 0 | 0.0 |
| Others † | 0 | ~ | 0 | ~ | 0 | 0.0 | 0.0 | 0 | 0.0 | 0.0 | 0 | 0.0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0.0 | 0.0 | 0.0 |

*Screening plus confirmatory test


**Excludes cases pending review by Public Health Investigator

***Positive HIV cases with no record of a previous positive HIV test in STDP Casewatch, Public Health Laboratory data, or through self-report on Clinic Intake Form

†Includes transgender and gender unspecified

Figure 13.1a. Sexually Transmitted Disease Prevalence, 2006-2010


Whittier Health Center



Note: Early Syphilis, HIV and New HIV trends shown in greater detail on next figure

Figure 13.1b. Early Syphilis, HIV, and New HIV Prevalence, 2006-2010

Whittier Health Center



Sexually Transmitted Disease Clinic Morbidity Report, 2010

XIV: Appendices

APPENDIX A: LABORATORY DIAGNOSTIC TESTS USED FOR THE DIAGNOSIS OF SEXUALLY TRANSMITTED DISEASES IN LAC STD CLINICS IN 2010

All specimens sent from the Los Angeles County STD Clinics for gonorrhea, chlamydia, syphilis, herpes, and HIV were tested at the Department of Public Health, Public Health Laboratory. Cervical cytology specimens were tested by the Laboratory Corporation of America (LabCorp).

Testing for *Neisseria gonorrhoeae* and *Chlamydia trachomatis*

Chlamydia trachomatis and *Neisseria gonorrhoeae* were detected from urine, vaginal, and endocervical swabs using a nucleic acid amplification test (NAAT) Aptima Combo 2 Assay (Gen-Probe, Inc., San Diego, CA). This same test technology was validated by the Public Health Laboratory to test for chlamydia and gonorrhea in rectal specimens and gonorrhea in pharyngeal specimens. In this report, urogenital testing refers to any urine, urethral, cervical or vaginal specimens.

Testing for syphilis

Blood was screened for syphilis using the Rapid Plasma Reagin (RPR) test which is a nontreponemal assay. The RPR antigen detects reagin, a nonspecific antibody present in the serum of persons with syphilis. Positive RPR tests were followed by the treponemal, confirmatory Treponema pallidum-particle agglutination (TP-PA) assay. The TP-PA detects *Treponema pallidum* antibodies in the serum as an aid in the diagnosis of syphilis and provides useful epidemiological information.

Testing for HIV

Until March 2008, blood was screened for antibody to HIV-1 using a first generation enzyme-linked immunosorbent assay (ELISA), Vironostika HIV-1 Microelisa System (Organon Teknika). In March 2008 the laboratory changed to use a more sensitive third generation test ADVIA Centaur XP, (Siemens Medical Solutions Diagnostics, East Walpole, MA). This is a chemiluminescent assay for HIV 1&2 plus GpO. Positive sera were confirmed using a Western Blot test (Genetic Systems Bio-Rad Laboratories).

Testing for Herpes

Specimens were collected from genital ulcers and other symptomatic sites and tested for Herpes Simplex Virus using a real time polymerase chain reaction method (PCR) (Real Art HSV ½ Light Cycler, Roche Diagnostics) that has been validated by the Public Health Laboratory for performance.

Cervical Cytology

Cervical cytology was performed using a liquid based Pap test (ThinPrep Pap Test, Hologic, MA). A reflex High-Risk HPV DNA test (Digene Hybrid Capture, Qiagen, MD) was performed if the pap result showed Atypical Squamous Cells of Undetermined Significance (AS-CUS).

APPENDIX B: DATA MANAGEMENT

Data Management Methods

With the exception of Antelope Valley Health Center visits, the Public Health Information System (PHIS) was used to examine all visits classified as “STD Clinical Visits.” The information collected from PHIS included demographic information such as gender, age, and race/ethnicity. Specific visit information such as clinic visited, date of visit, and visit classification (i.e., “New” or “Return” visit) was also obtained from PHIS. The Quadramed System provided the demographic and visit information for Antelope Valley Health Center.

Additional information on gender of sex partners and HIV testing history was obtained from the H2541 Intake Forms. The H2541 Intake Form is interviewer-administered to patients at new visits (see below for “new visit” definition) and collects more extensive demographic, behavioral risks, std history, and treatment information.

Sexual orientation of patients was based on the self-reported gender of sex partners from the H2541 Intake Form, which recorded the gender of sex partner(s) for the following two time frames: past 12 months and past 3 months. All available H2541 Intake Forms for each particular patient dating back to 2006 was used for information on gender of sex partners. If a patient had previously only reported no sex partners or refused to report the gender of sex partners on the H2541 form(s), then the sexual orientation for that patient was listed as “unknown”.

SUNQUEST, the Public Health Laboratory database system was used to obtain laboratory results for chlamydia, gonorrhea, syphilis, and HIV. The laboratory results were merged with the PHIS/Quadramed data using the medical record number and the date of clinic visit/specimen collection. In addition, a definitive syphilis diagnosis for those with a positive confirmatory syphilis test was obtained from STD*CASEWATCH. “New HIV” status for patients testing HIV positive was based on previous HIV testing information from all available SUNQUEST, STD*CASEWATCH, and H2541 Intake Form data. If there was no record of a previous positive HIV test in any of these data archives, then the HIV positive patient was designated as a “new HIV” case. The final dataset contains one record for each patient, per visit to the STD clinic (ie. individual visit). The “individual visit” is the unit of observation used in all tables and figures and referred to as such.

Data Limitations

Possibility of multiple visits: If a patient sought care for the same event at multiple clinics, each visit was captured as a separate visit.

Definition of new visits: In order to account for multiple return visits for the same event, a separate analysis was conducted examining only visits classified as “New Visits.” The Department of Health Services’ “Procedures for Completing the Clinic Data/Patient Encounter Form” (Policy No. 301) specifies that:

- A “New Visit” is one in which a person is seeking service for the first time, or for a new episode, in an established clinic in the facility. Examples include a first time visit to a STD clinic; or a visit for a new episode, if the patient was previously treated adequately.
- A “Return Visit” is one in which a patient has a follow-up visit to a clinic or ancillary service in the facility for further evaluation, treatment, or results.

Clinic sessions: A clinic session is a half-day when the STD clinic is open. The number of sessions held varies by STD clinic from 2 sessions a week to 9 sessions a week.



STD CLINIC REGISTRATION SCHEDULE

June 2010

Please note, all schedules are subject to change. Patients should be advised to call the STD clinic to confirm that a particular session is being held. Patients are seen on a first come, first served (walk-in) basis. In general, it is advisable for patients to arrive as early as possible to improve the likelihood of being seen.

No fees are charged for services in the Los Angeles County Department of Public Health STD clinics.

| ANTELOPE VALLEY HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|-------------------------------|------|---|------------|-----|-------|-----|
| | A.M. | | 8:00-10:30 | | | |
| | P.M. | 12:00-3:00 Nurse Screening Clinic. STD/HIV testing. Only for patients with NO symptoms. | 12:00-3:30 | | | |

335-B EAST AVENUE K-6, LANCASTER, CA 93535 *Between Division Street & Gingham Avenue*
Business Office (661) 723-4526

| CENTRAL HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|-----------------------|------|------------|------------|------------|---|------------|
| | A.M. | 8:00-10:30 | 8:00-10:30 | 8:00-10:30 | 8:00-10:30 | 8:00-10:30 |
| | P.M. | 12:00-3:30 | 12:00-3:30 | 12:00-3:30 | 12:00-3:30 PM session Closed on 4th Thurs. | |

241 N. FIGUEROA STREET, LOS ANGELES, CA 90012 *Just S of Temple Street* (213) 240-8203/Clinic or (213) 240-8225

| HOLLYWOOD-WILSHIRE HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|----------------------------------|------|------------|------------|------------|------------|------------|
| | A.M. | 7:30-10:30 | 10:00-1:30 | 7:30-10:30 | 7:30-10:30 | 7:30-10:30 |
| | P.M. | 12:00-3:30 | 3:00-5:45 | 12:00-3:30 | | |

5205 MELROSE AVENUE, LOS ANGELES, CA 90038 *At Wilton Place* (323) 769-7932/Clinic

| MONROVIA HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|------------------------|------|------------|------|-------------|------------|-----|
| | A.M. | | | 10:00-12:30 | | |
| | P.M. | 12:30-3:00 | | | 12:30-3:00 | |

330 W. MAPLE AVENUE, MONROVIA, CA 91016 *Between Magnolia Avenue & Mayflower Avenue* (626) 256-1600/Clinic

| NORTH HOLLYWOOD HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|-------------------------------|------|------------|------------|------------|------------|--|
| | A.M. | 8:00-10:30 | 8:00-10:30 | 8:00-10:30 | 8:00-10:30 | |
| | P.M. | 11:30-3:30 | 11:30-3:30 | 11:30-3:30 | 11:30-3:30 | 11:30-3:30 Closed 3 rd Fri |

5300 TUJUNGA AVE., NORTH HOLLYWOOD, CA 91601 *Between Weddington Street & Chandler Boulevard* (818) 766-3982 or (818) 766-3983/Clinic

| POMONA HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|----------------------|------|-----|------------|------------|-------------|-----|
| | A.M. | | | | 10:00-12:30 | |
| | P.M. | | 12:30-3:00 | 12:30-3:00 | | |

750 S. Park Avenue, Pomona, CA 91766 *South of Mission Blvd. at 7th Street* Clinic (909) 868-0235

| SIMMS/MANN/BURKE HEALTH CENTER Note: Public-private partnership clinic; the County operates a free STD clinic on site at times shown. | | MON | TUES | WED | THURS | FRI |
|--|------|-----|------------|-----|-------|-----|
| | A.M. | | 8:00-10:30 | | | |
| | P.M. | | 12:00-2:30 | | | |

2509 PICO BLVD. (3rd Floor), SANTA MONICA, CA 90405 *At 25th Street* (310) 998-3203/Nursing

| SOUTH HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|---------------------|------|------------|------------|-------|------------|------------|
| | A.M. | 8:00-10:30 | 8:00-10:30 | 8:00- | 8:00-10:30 | 8:00-10:30 |
| | P.M. | 12:30-3:00 | 12:30-3:00 | 5:00 | 12:30-3:00 | |

1522 E. 102ND STREET, LOS ANGELES, CA 90002 *Between Success Avenue & Compton Avenue* (323) 563-4112/Clinic

| RUTH TEMPLE HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|---------------------------|------|------------|------------|------------|-------------|------------|
| | A.M. | 7:30-10:30 | 7:30-10:30 | 7:30-10:30 | 10:00-12:30 | 7:30-10:30 |
| | P.M. | 12:00-2:30 | 12:00-2:30 | 12:00-2:30 | 2:00-5:00 | |

3834 S. WESTERN AVE, LOS ANGELES, CA 90062 *Between Exposition and M L King Boulevard* (323) 730-3576/Clinic

| TORRANCE HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|------------------------|------|-----------|------|-----|-------|--|
| | A.M. | | | | | |
| | P.M. | 1:45-5:00 | | | | 11:45 – 3:00 Closed 3 rd Fri |

711 W. DEL AMO BLVD., TORRANCE, CA 90502 *Between Vermont Avenue and 110 Freeway @ Hamilton Avenue*
Business Tel (310) 354-2300

| CURTIS TUCKER HEALTH CENTER | | MON | TUES | WED | THURS | FRI |
|-----------------------------|------|------------|------------|------------|------------|------------|
| | A.M. | 7:45-10:30 | 9:45-12:30 | 7:45-10:30 | | 7:45-10:30 |
| | P.M. | 11:45-3:00 | 1:45-5:00 | 11:45-3:00 | 11:45-3:00 | |

123 W. MANCHESTER BOULEVARD, INGLEWOOD, CA 90301 *West of La Brea at Fir Street* (310) 419-5362/Clinic

| WHITTIER HEALTH CENTER <i>*Please call ahead before referring patients.</i> | | MON | TUES | WED | THURS | FRI |
|--|------|------------|------------|-----|-----------|-----|
| | A.M. | | | | | |
| | P.M. | 12:00-3:00 | 12:00-3:00 | | 4:00-6:00 | |

7643 S. PAINTER AVENUE, WHITTIER, CA 90602 *Between Mar Vista Street & Whittier Boulevard* (562) 464-5350/Clinic

Submit changes to Ana Delia Hernandez at (213) 744-3057; FAX (213) 749-9605 Los Angeles County STD Program
For more information on STDs or a referral to other low- or no-cost clinics in Los Angeles County, please call the STD Hotline at **1-800-758-0880**

Patient's Primary Reason(s) for Visit

Sex partner in clinic today? No / Yes

- Symptoms present
Positive test done elsewhere:
Referral PHI / PHN / Other
Contact to:
Other:
Check-up only (i.e. none of the above).

Gender and Pregnancy Status

- Patient:
Male
Female
Transgender (M to F)
Transgender (F to M)
Other

- Sex partner(s): 12 mths 3 mths
Males
Females
Transgender (M to F)
Transgender (F to M)
Unknown
None

- Pregnant:
No Yes Unk.
Current partner(s) pregnant:
No Yes Unk.

STD History

- Ever Last 12 mths
None
Gonorrhea
Chlamydia
NSU / cervicitis
PID
Trichomonas
Anogenital warts
Genital herpes
Hepatitis A
Hepatitis B
Hepatitis C
Other:
Syphilis ever
Date
Stage
Rx
HIV
Date
In care
On HAART

Test History

- Date last HIV test:
Result:
Last RPR (date/result):
Date last Pap test:
Result:
Pap prior to last Pap:

Vaccination History

- HPV:
Hep A:
Hep B:

Sexual Practices

Last sexual contact Days / Wks / Mths / Yrs Ago Never

No. partners last 3 mths:
New partner last 3 mths: No Yes

- Type of Sex:
Condom / Barrier use 3 mths:
Always Never Some
Vaginal
Anal
Oral
None

If had vaginal sex in the last 5 days:
At risk of unwanted pregnancy? No Yes

Risk Behavior

- LAST 12 MTHS:
No risk identified
Any sex without a condom
Exchanged drugs/money for sex
Incarcerated
Sex with partner recently incarcerated
Sex with IDU
Sex with HIV pos. person
Sex with anonymous partner
Sex at bathhouse/sex club
Met casual partners online
Sex with a man who also has sex with men?
Other HIV risk since last HIV test:

Recreational Drug / Alcohol Use

- In last 12 mths:
None
Marijuana
Methamphetamine
Heroin
Ecstasy
Cocaine
Crack
Viagra/Cialis/Levitra
Binge drinking
Other:

IDU ever: No Yes
IDU last 12 mths? No Yes
Shared equipment 12 mths? No Yes

Tobacco: Current user Ex-user Never used Referred

Other

- Intimate partner violence: at risk? No Yes:
Risk reduction plan: plan developed today? No Yes (page 2:)
Referral made?(other than tobacco) No Yes:
HIV test today: Accepted Declined Not indicated

Intake interview by:
Interviewer Number
PHI/RN/LVN/CW/MD

Contact interview done by:
PHI/PHN/RN

Patient identification

Symptoms Onset / Duration

- Oropharyngeal
Dysuria: external / internal
Urinary: frequency / urgency
Genital rash / lesions
Non-genital rash
Anorectal symptoms / discharge
Testicular pain
Urethral discharge
Vaginal discharge / odor
Vulval itch
Abdominal pain
Dyspareunia: external / internal
Abnormal vaginal bleeding
OTHER:

Females only

- LMP ___/___/___
Pregnant now: # wks
Pregnant in last 3 mths?
Breastfeeding?
Douches?
Current Contraception?
No: Wants FP referral?
Yes: Withdrawal, Condom, Hormonal, IUD, Other

Medications/Drug Allergies

- Current Medication:
Drug Allergies No Yes

Risk Reduction Plan / Counseling / Other

- Condoms given
Explained STD/HIV tests
Difficulty with condoms? No Yes

Interviewer

History of Presenting Illness / Exam

History of Presenting Illness / Exam section with dotted lines for notes.

EXAM FINDINGS

Table with columns for body parts (Oral cavity, Skin, Inguinal LN, etc.) and findings (Nml, Abn, ND).

patient identification

Provider MD/NP

Visit Date (stamp)

Assessment & Plan

Mark all that apply OR STD screening only

STD screening only

STD symptoms, no signs

Syndrome / Diagnosis

- Urethritis
Cervicitis
Vaginitis/balanitis
Proctitis
Pharyngitis
Epididymitis
PID
UTI
GUD (not chancere)
Suspected chancere location:
Signs of secondary Σ:

- Trich
BV
Yeast
Warts
Lice
Scabies
Molluscum

Non-STD condition:

Positive Lab.

- CT # partners 60 days:
GC # partners 60 days:
Herpes
Σ primary
Σ secondary
Σ early latent
Σ late latent/ unknown duration
Σ old treated
BFP
Other:

Contact to

- *interval since last sex with contact:
CT
GC
Trich
Σ Early*
Σ late latent*
Σ unknown duration*
Σ stage unk. / TBD*
Herpes
Other:

Tests

Wet mount: Normal

PMN:EC <1 ≥1
Trich Yeast >20% clue cells

Whiff test: Neg Pos.

Vag. pH: <4.5 ≥4.5

Order Done

Temp BP

Pregnancy: Neg Pos.

Urinalysis:

Leukocyte: Neg Pos.
Nitrates: Neg Pos.

- Vagina CT GC
Urine CT GC
Cervix CT GC
Urethra CT GC
Throat GC
Rectum CT GC
HSV PCR (site)
HSV-2 serology
HIV serology
RPR TPPA
RPR only
PAP Declined Not indicated
Other:

Results entered by:
Date:

Alternate regimen used due to contraindication:

Prescription

Administered / Dispensed

- Azithromycin 1g PO (250mg x 4) admin./disp.
PDPT CT Pack(s) x Azithromycin 1g
PDPT GC/CT Pack(s) x Azithromycin 1g + Cefpodoxime 400mg
Cefpodoxime 400mg PO (200mg x 2) admin./disp.
Ceftriaxone 125mg / 250mg IM R / L buttock / deltoid
Doxycycline 100mg PO BID x days tab.
Acyclovir 400mg PO BID / TID x days tab.
Bicillin LA 2.4 MU IM (give 1.2 MU in each buttock)
Metronidazole 500mg PO BID x 7days 14 tab.
Metronidazole 2g PO (500mg x 4) admin./disp.
Metronidazole vaginal gel 0.75% 70g 1 applicator nightly x5.
Fluconazole 150mg PO admin./disp.
Nystatin-triamcinolone cream 30g, apply to.
Clotrimazole 1% cream 30g Vaginal antifungal: (specify)
Liquid nitrogen TCA
Aldara 5% Cream 25g
Condylox gel 0.5% 3.5g
Twinrix 1ml Engerix B 1ml Havrix 1ml
Levonorgestrel 1.5mg PO x pack(s) admin./disp.
No Rx Other:

Explained: Diagnosis Rx PDPT Abstain 1wk

Referrals: None PMD HIV care FPC/preg.
Mental health Alc./drugs Needle ex. IZ clinic
IPV Gyn. Other:

Follow-up: Phone In-person results MD:

Initials

Provider MD/NP

Broken Phone or Return Appointment

NEEDS: Rx Test Interview Notify of result None
PLAN: Close chart Refer PHI / PHN Call patient B&M

Sign & date MD/RN/NP

CMR completed Health Dept. follow-up initiated

patient identification

Visit Date (stamp)



COUNTY OF LOS ANGELES
Public Health

Sexually Transmitted Disease Program

2615 South Grand Avenue, Room 500
Los Angeles, CA 90007
Tel: (213) 744-3070 Fax: (213) 749-9606

<http://publichealth.lacounty.gov/std>