

Appendix A-1

Demographic Characteristics by Service Planning Area, 1997¹

ADULTS 18 YEARS AND OLDER	ANTELOPE VALLEY			SAN FERNANDO VALLEY			SAN GABRIEL VALLEY			METRO		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
AGE GROUP												
18-29 Years	25%	(±7%)	44,000	25%	(±2%)	331,000	23%	(±2%)	293,000	29%	(±3%)	243,000
30-39 Years	23%	(±7%)	41,000	27%	(±2%)	357,000	25%	(±2%)	310,000	26%	(±3%)	219,000
40-49 Years	27%	(±8%)	49,000	19%	(±2%)	257,000	18%	(±3%)	225,000	18%	(±3%)	154,000
50+ Years	26%	(±7%)	46,000	29%	(±2%)	387,000	34%	(±2%)	433,000	26%	(±3%)	220,000
GENDER												
Males	47%	(± 8%)	84,000	53%	(± 3%)	710,000	50%	(± 3%)	629,000	54%	(± 3%)	452,000
Females	53%	(± 8%)	95,000	47%	(± 3%)	623,000	50%	(± 3%)	632,000	46%	(± 3%)	384,000
RACE AND ETHNICITY												
White	63%	(±8%)	112,000	58%	(±3%)	763,000	38%	(±3%)	479,000	33%	(±3%)	273,000
African American	7%	(±4%)	13,000	4%	(±1%)	50,000	5%	(±1%)	57,000	5%	(±2%)	44,000
Latino	23%	(±7%)	40,000	29%	(±2%)	386,000	38%	(±3%)	481,000	48%	(±3%)	396,000
Asian	5%	(±4%)	9,000	8%	(±1%)	101,000	18%	(±2%)	229,000	13%	(±2%)	111,000
WELFARE STATUS												
On welfare	7%	(±4%)	13,000	6%	(±1%)	79,000	8%	(±1%)	98,000	11%	(±2%)	95,000
Not on welfare	93%	(±4%)	165,000	94%	(±1%)	1,249,000	92%	(±1%)	1,157,000	89%	(±2%)	736,000
EDUCATIONAL STATUS												
Less than high school	18%	(±7%)	33,000	14%	(±2%)	184,000	17%	(±2%)	217,000	25%	(±3%)	208,000
High school	25%	(±7%)	45,000	21%	(±2%)	276,000	23%	(±2%)	290,000	21%	(±3%)	174,000
Some college/trade	32%	(±8%)	57,000	28%	(±2%)	374,000	28%	(±2%)	345,000	22%	(±3%)	184,000
Graduate/post graduate	24%	(±7%)	43,000	37%	(±2%)	491,000	32%	(±3%)	398,000	32%	(±3%)	265,000
POVERTY STATUS ³												
Below 100% FPL	17%	(±6%)	30,000	9%	(±1%)	125,000	14%	(±2%)	175,000	20%	(±3%)	168,000
Between 100-200% FPL	16%	(±6%)	30,000	21%	(±2%)	283,000	26%	(±2%)	330,000	29%	(±3%)	238,000
Above 200% FPL	67%	(±8%)	119,000	69%	(±2%)	925,000	60%	(±3%)	756,000	51%	(±3%)	430,000
HOMELESS STATUS ⁴												
Yes	5%	(±3%)	8,000	5%	(±1%)	66,000	5%	(±1%)	62,000	8%	(±2%)	63,000
No	95%	(±3%)	171,000	95%	(±1%)	1,265,000	95%	(±1%)	1,194,000	92%	(±2%)	770,000

Appendix A-1 Continued

Demographic Characteristics by Service Planning Area, 1997¹

ADULTS 18 YEARS AND OLDER	WEST			SOUTH			EAST			SOUTH BAY		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
AGE GROUP												
18-29 Years	25%	(±4%)	121,000	32%	(±4%)	196,000	29%	(±3%)	252,000	25%	(±2%)	264,000
30-39 Years	23%	(±3%)	110,000	27%	(±4%)	164,000	27%	(±3%)	235,000	28%	(±3%)	289,000
40-49 Years	18%	(±3%)	89,000	17%	(±3%)	103,000	17%	(±2%)	152,000	17%	(±2%)	176,000
50+ Years	34%	(±4%)	164,000	24%	(±3%)	145,000	27%	(±3%)	241,000	30%	(±3%)	318,000
GENDER												
Males	54%	(± 4%)	259,000	40%	(± 4%)	246,000	50%	(± 3%)	445,000	49%	(± 3%)	512,000
Females	46%	(± 4%)	225,000	60%	(± 4%)	365,000	50%	(± 3%)	437,000	51%	(± 3%)	536,000
RACE AND ETHNICITY												
White	69%	(±4%)	328,000	5%	(±2%)	33,000	31%	(±3%)	270,000	46%	(±3%)	478,000
African American	7%	(±2%)	36,000	39%	(±4%)	236,000	5%	(±1%)	42,000	16%	(±2%)	161,000
Latino	14%	(±3%)	65,000	53%	(±4%)	319,000	59%	(±3%)	516,000	27%	(±3%)	280,000
Asian	9%	(±2%)	42,000	2%	(±1%)	13,000	4%	(±1%)	37,000	10%	(±2%)	103,000
WELFARE STATUS												
On welfare	5%	(±2%)	22,000	16%	(±3%)	99,000	9%	(±2%)	82,000	7%	(±1%)	78,000
Not on welfare	95%	(±2%)	461,000	84%	(±3%)	508,000	91%	(±2%)	793,000	93%	(±1%)	966,000
EDUCATIONAL STATUS												
Less than high school	6%	(±2%)	27,000	40%	(±4%)	242,000	26%	(±3%)	225,000	16%	(±2%)	171,000
High school	13%	(±3%)	61,000	26%	(±3%)	155,000	26%	(±3%)	231,000	20%	(±2%)	212,000
Some college/trade	25%	(±4%)	122,000	19%	(±3%)	117,000	29%	(±3%)	259,000	29%	(±3%)	304,000
Graduate/post graduate	56%	(±4%)	268,000	15%	(±3%)	91,000	19%	(±3%)	165,000	34%	(±3%)	356,000
POVERTY STATUS³												
Below 100% FPL	7%	(±2%)	35,000	35%	(±4%)	213,000	19%	(±3%)	164,000	12%	(±2%)	125,000
Between 100-200% FPL	15%	(±3%)	73,000	33%	(±4%)	200,000	29%	(±3%)	257,000	23%	(±2%)	244,000
Above 200% FPL	78%	(±3%)	377,000	32%	(±4%)	198,000	52%	(±3%)	462,000	65%	(±3%)	680,000
HOMELESS STATUS⁴												
Yes	4%	(±2%)	21,000	7%	(±2%)	42,000	6%	(±2%)	53,000	6%	(±1%)	59,000
No	96%	(±2%)	462,000	93%	(±2%)	568,000	94%	(±2%)	827,000	94%	(±1%)	987,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. For a family of four, the 1997 federal poverty level (FPL) was \$16,050 a year and 200% of poverty was \$32,100.

4. Homelessness in the past 5 years.

Appendix A-2

Demographic Characteristics by Service Planning Area, 1997¹

CHILDREN 0 TO 17 YEARS	ANTELOPE VALLEY			SAN FERNANDO VALLEY			SAN GABRIEL VALLEY			METRO		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
AGE GROUP												
0-4 years	22%	(±12%)	23,000	31%	(±5%)	153,000	33%	(±5%)	191,000	34%	(±8%)	112,000
5-12 years	55%	(±15%)	57,000	46%	(±6%)	228,000	47%	(±6%)	272,000	42%	(±8%)	139,000
13-17 years	23%	(±12%)	24,000	23%	(±5%)	111,000	21%	(±5%)	121,000	25%	(±7%)	82,000
GENDER												
Males	50%	(±15%)	52,000	51%	(±6%)	249,000	50%	(±6%)	290,000	46%	(±8%)	152,000
Females	50%	(±15%)	52,000	49%	(±6%)	242,000	50%	(±6%)	294,000	54%	(±8%)	180,000
RACE AND ETHNICITY												
White	54%	(±15%)	56,000	34%	(±5%)	165,000	18%	(±4%)	102,000	6%	(±4%)	21,000
African American	9%	(±8%)	9,000	3%	(±2%)	15,000	4%	(±2%)	23,000	2%	(±2%)	8,000
Latino	33%	(±14%)	34,000	55%	(±6%)	270,000	60%	(±6%)	348,000	83%	(±6%)	272,000
Asian	2%	(±5%)	3,000	8%	(±3%)	38,000	18%	(±4%)	104,000	9%	(±5%)	29,000
PARTICIPATION IN DISABILITY PROGRAMS												
Child health	9%	(±8%)	9,000	22%	(±5%)	106,000	15%	(±4%)	85,000	26%	(±7%)	84,000
Head Start	8%	(±8%)	8,000	11%	(±3%)	53,000	13%	(±4%)	76,000	11%	(±5%)	35,000
WIC	20%	(±12%)	21,000	29%	(±5%)	141,000	30%	(±5%)	171,000	54%	(±8%)	178,000
POVERTY STATUS ³												
Below 100% FPL	31%	(±14%)	32,000	24%	(±5%)	118,000	27%	(±5%)	155,000	54%	(±8%)	180,000
Between 100-200% FPL	16%	(±11%)	17,000	28%	(±5%)	138,000	33%	(±5%)	192,000	29%	(±7%)	96,000
Above 200% FPL	53%	(±15%)	55,000	48%	(±6%)	234,000	41%	(±6%)	237,000	17%	(±6%)	57,000

Appendix A-2 Continued

Demographic Characteristics by Service Planning Area, 1997¹

CHILDREN 0 TO 17 YEARS	WEST			SOUTH			EAST			SOUTH BAY		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
AGE GROUP												
0-4 years	34%	(±12%)	29,000	41%	(±7%)	166,000	28%	(±6%)	127,000	33%	(±6%)	128,000
5-12 years	43%	(±12%)	36,000	36%	(±7%)	147,000	44%	(±6%)	198,000	45%	(±6%)	176,000
13-17 years	23%	(±10%)	19,000	23%	(±6%)	94,000	28%	(±5%)	124,000	21%	(±5%)	83,000
GENDER												
Males	53%	(±12%)	45,000	48%	(± 8%)	194,000	56%	(±6%)	252,000	55%	(±6%)	214,000
Females	47%	(±12%)	40,000	52%	(± 8%)	212,000	44%	(±6%)	197,000	45%	(±6%)	172,000
RACE AND ETHNICITY												
White	51%	(±12%)	43,000	2%	(±2%)	6,000	24%	(±5%)	106,000	23%	(±5%)	90,000
African American	7%	(±6%)	6,000	22%	(±6%)	91,000	4%	(±3%)	20,000	18%	(±5%)	67,000
Latino	34%	(±12%)	28,000	76%	(±7%)	306,000	67%	(±6%)	303,000	49%	(±6%)	188,000
Asian	8%	(±7%)	7,000	0%		0	3%	(±2%)	15,000	10%	(±4%)	38,000
PARTICIPATION IN PROGRAMS												
Child Health & Disability	9%	(±7%)	7,000	23%	(±6%)	93,000	20%	(±5%)	89,000	20%	(±5%)	74,000
Head Start	7%	(±6%)	6,000	13%	(±5%)	51,000	6%	(±3%)	26,000	16%	(±5%)	60,000
WIC	20%	(±10%)	17,000	50%	(±8%)	203,000	35%	(±6%)	157,000	36%	(±6%)	137,000
POVERTY STATUS												
Below 100% FPL	26%	(±11%)	22,000	62%	(±7%)	252,000	32%	(±6%)	144,000	31%	(±6%)	120,000
Between 100-200% FPL	23%	(±10%)	19,000	23%	(±6%)	93,000	28%	(±6%)	126,000	26%	(±6%)	102,000
Above 200% FPL	51%	(±12%)	43,000	15%	(±5%)	61,000	40%	(±6%)	180,000	43%	(±6%)	165,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Parents' poverty status. For a family of four, the 1997 federal poverty level (FPL) was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-1

Health Behaviors, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	CIGARETTE SMOKERS ⁵			HEAVY ALCOHOL USERS ³			HANDGUN OWNERS ⁶			DO NOT USE SEAT BELTS		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER												
Males	22%	(±2%)	740,000	8%	(±1%)	272,000	26%	(±2%)	869,000	14%	(±1%)	450,000
Females	14%	(±1%)	470,000	2%	(±0%)	58,000	14%	(±1%)	463,000	9%	(±1%)	287,000
RACE AND ETHNICITY												
White	21%	(±1%)	564,000	6%	(±1%)	170,000	28%	(±2%)	742,000	12%	(±1%)	325,000
African American	21%	(±3%)	131,000	2%	(±1%)	13,000	22%	(±3%)	135,000	11%	(±2%)	71,000
Latino	16%	(±1%)	387,000	5%	(±1%)	130,000	12%	(±1%)	307,000	11%	(±1%)	262,000
Asian	17%	(±3%)	108,000	2%	(±1%)	11,000	18%	(±3%)	113,000	10%	(±2%)	62,000
POVERTY STATUS⁴												
Below 100% FPL	20%	(±2%)	202,000	4%	(±1%)	45,000	8%	(±2%)	81,000	11%	(±2%)	112,000
Between 100–200% FPL	20%	(±2%)	332,000	6%	(±1%)	92,000	15%	(±2%)	249,000	11%	(±1%)	179,000
Above 200% FPL	17%	(±1%)	675,000	5%	(±1%)	194,000	26%	(±1%)	1,001,000	11%	(±1%)	446,000
AGE GROUP												
18–29 years	18%	(±2%)	323,000	5%	(±2%)	88,000	17%	(±2%)	290,000	14%	(±2%)	245,000
30–39 years	18%	(±2%)	314,000	4%	(±2%)	75,000	18%	(±2%)	313,000	10%	(±1%)	179,000
40–49 years	21%	(±2%)	249,000	4%	(±2%)	50,000	22%	(±2%)	263,000	9%	(±2%)	109,000
50 years and older	17%	(±2%)	322,000	6%	(±2%)	118,000	24%	(±2%)	466,000	11%	(±1%)	203,000
EDUCATION STATUS												
Less than high school	19%	(±2%)	243,000	5%	(±1%)	64,000	9%	(±2%)	121,000	12%	(±2%)	151,000
High school	20%	(±2%)	294,000	6%	(±1%)	80,000	20%	(±2%)	277,000	12%	(±2%)	169,000
Some college or trade school	21%	(±2%)	373,000	6%	(±1%)	95,000	26%	(±2%)	449,000	12%	(±1%)	208,000
Graduate or post graduate	14%	(±1%)	293,000	4%	(±1%)	91,000	24%	(±2%)	482,000	10%	(±1%)	203,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified based on the number of drinks consumed. A drink is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Heavy alcohol users are those who consume 1.0 or more ounces of ethyl alcohol daily.

4. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

5. Adults (18 and older) who currently smoke cigarettes.

6. Adults who own a rifle, shotgun, pistol or revolver.

Appendix B-2

Health Behaviors, Los Angeles County, 1997¹

CHILDREN 0 TO 17 YEARS	DO NOT WEAR SEAT BELTS		
	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER			
Males	5%	(±2%)	72,000
Females	4%	(±1%)	57,000
RACE AND ETHNICITY			
White	3%	(±2%)	19,000
African American	4%	(±3%)	10,000
Latino	5%	(±1%)	84,000
Asian	7%	(±4%)	16,000
POVERTY STATUS³			
Below 100% FPL	5%	(±2%)	55,000
Between 100–200% FPL	4%	(±2%)	35,000
Above 200% FPL	4%	(±1%)	39,000
AGE GROUP			
Less than 4 years	2%	(±3%)	20,000
5–12 years	6%	(±2%)	72,000
13–17 years	6%	(±1%)	37,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-3

Health Status, Los Angeles County, 1997¹

CHILDREN 0 TO 17 YEARS	ASTHMA		
	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER			
Males	8%	(±2%)	114,000
Females	7%	(±2%)	92,000
RACE AND ETHNICITY			
White	8%	(±3%)	46,000
African American	19%	(±6%)	45,000
Latino	6%	(±2%)	97,000
Asian	7%	(±4%)	16,000
POVERTY STATUS³			
Below 100% FPL	8%	(±3%)	77,000
Between 100–200% FPL	7%	(±2%)	53,000
Above 200% FPL	7%	(±2%)	76,000
AGE GROUP			
Less than 4 years	4%	(±2%)	40,000
5–12 years	9%	(±2%)	108,000
13–17 years	9%	(±3%)	57,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-4

Health Status, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	FAIR OR POOR HEALTH ⁵			SEVERELY OVERWEIGHT ⁴			ARTHRITIS		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER									
Males	17%	(±1%)	577,000	13%	(±1%)	437,000	13%	(±1%)	426,000
Females	24%	(±1%)	782,000	15%	(±1%)	457,000	22%	(±1%)	725,000
RACE AND ETHNICITY									
White	12%	(±1%)	315,000	12%	(±1%)	326,000	22%	(±1%)	600,000
African American	18%	(±3%)	114,000	22%	(±3%)	137,000	22%	(±3%)	140,000
Latino	30%	(±2%)	741,000	17%	(±3%)	384,000	12%	(±1%)	287,000
Asian	27%	(±3%)	171,000	4%	(±2%)	25,000	15%	(±3%)	96,000
POVERTY STATUS⁵									
Below 100% FPL	40%	(±3%)	414,000	20%	(±3%)	184,000	18%	(±2%)	178,000
Between 100–200% FPL	27%	(±2%)	447,000	18%	(±2%)	278,000	21%	(±2%)	346,000
Above 200% FPL	13%	(±1%)	498,000	11%	(±1%)	432,000	16%	(±1%)	626,000
AGE GROUP									
18–29 years	17%	(±2%)	293,000	9%	(±1%)	154,000	3%	(±1%)	50,000
30–39 years	18%	(±2%)	315,000	14%	(±2%)	225,000	7%	(±1%)	127,000
40–49 years	21%	(±2%)	253,000	16%	(±2%)	186,000	17%	(±2%)	203,000
50 years and older	26%	(±2%)	497,000	18%	(±2%)	330,000	40%	(±2%)	769,000
EDUCATION STATUS									
Less than high school	42%	(±3%)	546,000	22%	(±2%)	243,000	21%	(±2%)	276,000
High school	23%	(±2%)	337,000	13%	(±2%)	185,000	18%	(±2%)	255,000
Some college or trade school	15%	(±2%)	258,000	15%	(±2%)	249,000	18%	(±2%)	315,000
Graduate or post graduate	10%	(±1%)	201,000	10%	(±1%)	210,000	14%	(±1%)	290,000

Appendix B-4 Continued

Health Status, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	DIABETES			HYPERTENSION		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER						
Males	5%	(±1%)	177,000	14%	(±1%)	460,000
Females	6%	(±1%)	194,000	18%	(±1%)	581,000
RACE AND ETHNICITY						
White	5%	(±1%)	138,000	17%	(±1%)	455,000
African American	9%	(±2%)	57,000	28%	(±3%)	176,000
Latino	6%	(±1%)	136,000	12%	(±1%)	290,000
Asian	5%	(±2%)	30,000	14%	(±3%)	91,000
POVERTY STATUS						
Below 100% FPL	6%	(±1%)	63,000	17%	(±2%)	584,000
Between 100–200% FPL	7%	(±1%)	118,000	18%	(±2%)	290,000
Above 200% FPL	5%	(±1%)	191,000	15%	(±1%)	168,000
AGE GROUP						
18–29 years	1%	(±0%)	15,000	4%	(±1%)	71,000
30–39 years	3%	(±1%)	47,000	7%	(±1%)	114,000
40–49 years	5%	(±1%)	61,000	14%	(±2%)	169,000
50 years and older	13%	(±1%)	249,000	35%	(±2%)	685,000
EDUCATION STATUS						
Less than high school	8%	(±1%)	103,000	19%	(±2%)	243,000
High school	6%	(±1%)	82,000	17%	(±2%)	238,000
Some college or trade school	5%	(±1%)	92,000	17%	(±2%)	290,000
Graduate or post graduate	4%	(±1%)	88,000	13%	(±1%)	261,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Self-perceived health status.

4. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, BMI >= 30 is classified as severely overweight.

5. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-5

Access to Care and Health Service Utilization, Los Angeles County, 1997¹

CHILDREN 0 TO 17 YEARS	UNINSURED			NO REGULAR SOURCE OF CARE		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER						
Males	26%	(±3%)	372,000	15%	(±3%)	216,000
Females	24%	(±3%)	323,000	15%	(±3%)	207,000
RACE AND ETHNICITY						
White	10%	(±3%)	57,000	4%	(±2%)	24,000
African American	14%	(±6%)	33,000	9%	(±5%)	21,000
Latino	33%	(±3%)	556,000	20%	(±3%)	345,000
Asian	20%	(±7%)	46,000	12%	(±6%)	26,000
POVERTY STATUS³						
Below 100% FPL	31%	(±4%)	314,000	20%	(±4%)	200,000
Between 100–200% FPL	32%	(±4%)	246,000	20%	(±4%)	153,000
Above 200% FPL	13%	(±3%)	135,000	7%	(±2%)	70,000
AGE GROUP						
Less than 4 years	24%	(±4%)	213,000	11%	(±3%)	104,000
5–12 years	24%	(±3%)	295,000	15%	(±3%)	190,000
13–17 years	29%	(±5%)	188,000	20%	(±4%)	129,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-6

Access to Care and Health Service Utilization, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	UNINSURED ²			NO REGULAR SOURCE OF CARE			DIFFICULTY GETTING CARE ⁴		
	Percent	(±95%) Confidence Interval ³	Estimated Number	Percent	(±95%) Confidence Interval ³	Estimated Number	Percent	(±95%) Confidence Interval ³	Estimated Number
GENDER									
Males	34%	(±2%)	988,000	30%	(±2%)	988,000	34%	(±2%)	1,095,000
Females	35%	(±2%)	984,000	21%	(±1%)	668,000	35%	(±1%)	1,134,000
RACE AND ETHNICITY									
White	24%	(±2%)	515,000	17%	(±1%)	466,000	23%	(±2%)	620,000
African American	26%	(±4%)	142,000	17%	(±3%)	104,000	32%	(±3%)	199,000
Latino	46%	(±2%)	1,081,000	36%	(±2%)	877,000	47%	(±2%)	1,146,000
Asian	35%	(±4%)	198,000	30%	(±4%)	187,000	36%	(±4%)	219,000
POVERTY STATUS⁵									
Below 100% FPL	56%	(±3%)	545,000	41%	(±3%)	408,000	54%	(±3%)	544,000
Between 100–200% FPL	45%	(±3%)	625,000	32%	(±2%)	512,000	46%	(±2%)	741,000
Above 200% FPL	24%	(±1%)	802,000	19%	(±1%)	736,000	25%	(±1%)	943,000
AGE GROUP									
18–29 years	42%	(±2%)	720,000	39%	(±2%)	665,000	43%	(±2%)	737,000
30–39 years	34%	(±2%)	587,000	28%	(±2%)	478,000	37%	(±2%)	630,000
40–49 years	31%	(±3%)	373,000	24%	(±2%)	280,000	34%	(±3%)	401,000
50 years and older	27%	(±2%)	292,000	12%	(±1%)	231,000	24%	(±2%)	456,000
EDUCATION STATUS									
Less than high school	54%	(±3%)	618,000	40%	(±3%)	509,000	51%	(±3%)	647,000
High school	38%	(±3%)	471,000	27%	(±2%)	381,000	37%	(±2%)	520,000
Some college or trade school	30%	(±2%)	452,000	21%	(±2%)	366,000	32%	(±2%)	552,000
Graduate or post graduate	23%	(±2%)	409,000	19%	(±2%)	385,000	25%	(±2%)	496,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. Adults between the age 18 to 64 years.

3. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

4. Somewhat difficult or very difficult getting care.

5. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-6 Continued

Access to Care and Health Service Utilization, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	PRESCRIPTION MEDICINES			MENTAL HEALTH			DENTAL CARE		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER									
Males	13%	(±1%)	421,000	7%	(±1%)	213,000	24%	(±2%)	788,000
Females	19%	(±1%)	628,000	10%	(±1%)	309,000	29%	(±1%)	965,000
RACE AND ETHNICITY									
White	12%	(±1%)	314,000	7%	(±1%)	194,000	20%	(±1%)	535,000
African American	16%	(±3%)	99,000	7%	(±2%)	44,000	26%	(±3%)	96,000
Latino	22%	(±2%)	542,000	9%	(±1%)	230,000	34%	(±2%)	850,000
Asian	13%	(±3%)	81,000	6%	(±2%)	41,000	27%	(±3%)	174,000
POVERTY STATUS³									
Below 100% FPL	30%	(±3%)	306,000	12%	(±2%)	124,000	43%	(±3%)	449,000
Between 100–200% FPL	22%	(±2%)	360,000	10%	(±1%)	169,000	37%	(±2%)	605,000
Above 200% FPL	10%	(±1%)	383,000	6%	(±1%)	229,000	18%	(±1%)	699,000
AGE GROUP									
18–29 years	17%	(±2%)	301,000	8%	(±1%)	134,000	32%	(±2%)	562,000
30–39 years	18%	(±2%)	305,000	9%	(±1%)	160,000	29%	(±2%)	505,000
40–49 years	17%	(±2%)	205,000	10%	(±2%)	122,000	29%	(±2%)	343,000
50 years and older	12%	(±1%)	237,000	5%	(±1%)	103,000	18%	(±2%)	342,000
EDUCATION STATUS									
Less than high school	24%	(±2%)	316,000	10%	(±2%)	131,000	37%	(±3%)	484,000
High school	17%	(±2%)	238,000	7%	(±1%)	105,000	29%	(±2%)	418,000
Some college or trade school	15%	(±2%)	272,000	7%	(±1%)	130,000	27%	(±2%)	467,000
Graduate or post graduate	10%	(±1%)	213,000	7%	(±1%)	149,000	18%	(±2%)	374,000

Appendix B-6 Continued

Access to Care and Health Service Utilization, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	EYEGLASSES			DIFFICULTY SEEING A DOCTOR		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER						
Males	12%	(±1%)	393,000	15%	(±1%)	507,000
Females	19%	(±1%)	619,000	21%	(±1%)	686,000
RACE AND ETHNICITY						
White	12%	(±1%)	325,000	13%	(±1%)	347,000
African American	15%	(±3%)	96,000	17%	(±3%)	111,000
Latino	19%	(±2%)	469,000	24%	(±2%)	603,000
Asian	15%	(±3%)	98,000	18%	(±3%)	113,000
POVERTY STATUS³						
Below 100% FPL	23%	(±3%)	229,000	33%	(±3%)	343,000
Between 100–200% FPL	21%	(±2%)	345,000	24%	(±2%)	402,000
Above 200% FPL	11%	(±1%)	438,000	11%	(±1%)	448,000
AGE GROUP						
18–29 years	15%	(±2%)	258,000	22%	(±2%)	381,000
30–39 years	14%	(±2%)	239,000	20%	(±2%)	351,000
40–49 years	19%	(±2%)	221,000	19%	(±2%)	234,000
50 years and older	15%	(±1%)	293,000	12%	(±1%)	226,000
EDUCATION STATUS						
Less than high school	21%	(±2%)	274,000	27%	(±2%)	353,000
High school	15%	(±2%)	221,000	19%	(±2%)	275,000
Some college or trade school	16%	(±2%)	283,000	18%	(±2%)	309,000
Graduate or post graduate	11%	(±1%)	226,000	12%	(±1%)	244,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix B-6 Continued

Access to Care and Health Service Utilization, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	HIV/AIDS			PROSTATE EXAM ²			PAP EXAM ³		
	Percent	(±95%) Confidence Interval ⁴	Estimated Number	Percent	(±95%) Confidence Interval ⁴	Estimated Number	Percent	(±95%) Confidence Interval ⁴	Estimated Number
GENDER									
Males	36%	(±2%)	1,190,000	51%	(±3%)	759,000			
Females	36%	(±1%)	1,163,000				73%	(±1%)	2,387,000
RACE AND ETHNICITY									
White	30%	(±2%)	810,000	58%	(±3%)	468,000	75%	(±2%)	985,000
African American	47%	(±4%)	291,000	53%	(±9%)	63,000	79%	(±4%)	286,000
Latino	43%	(±2%)	1,053,000	41%	(±5%)	145,000	70%	(±2%)	890,000
Asian	26%	(±3%)	161,000	36%	(±8%)	58,000	68%	(±5%)	189,000
POVERTY STATUS⁷									
Below 100% FPL	42%	(±3%)	428,000	33%	(±8%)	49,000	65%	(±3%)	408,000
Between 100–200% FPL	37%	(±2%)	610,000	39%	(±6%)	121,000	69%	(±3%)	567,000
Above 200% FPL	34%	(±1%)	1,315,000	57%	(±3%)	589,000	78%	(±2%)	1,413,000
AGE GROUP									
18–29 years	44%	(±2%)	757,000				71%	(±3%)	584,000
30–39 years	40%	(±2%)	683,000				77%	(±2%)	625,000
40–49 years	32%	(±3%)	381,000				77%	(±3%)	466,000
50 years and older	29%	(±2%)	531,000				68%	(±2%)	711,000
EDUCATION STATUS									
Less than high school	42%	(±3%)	542,000	34%	(±7%)	76,000	65%	(±3%)	490,000
High school	36%	(±2%)	515,000	50%	(±6%)	127,000	68%	(±3%)	490,000
Some college or trade school	35%	(±2%)	608,000	52%	(±5%)	201,000	76%	(±2%)	665,000
Graduate or post graduate	33%	(±2%)	672,000	58%	(±4%)	347,000	80%	(±2%)	732,000

Appendix B-6 Continued

Access to Care and Health Service Utilization, Los Angeles County, 1997¹

ADULTS 18 YEARS AND OLDER	CLINICAL BREAST EXAM ³			TESTICULAR EXAM ⁴			MAMMOGRAM ⁵		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
GENDER									
Males				28%	(±2%)	897,000			
Females	70%	(±1%)	2,310,000				76%	(±2%)	796,000
RACE AND ETHNICITY									
White	78%	(±2%)	1,027,000	33%	(±3%)	451,000	77%	(±3%)	466,000
African American	78%	(±4%)	281,000	32%	(±6%)	77,000	80%	(±6%)	99,000
Latino	62%	(±2%)	795,000	25%	(±3%)	289,000	75%	(±5%)	157,000
Asian	62%	(±5%)	173,000	17%	(±4%)	60,000	68%	(±9%)	58,000
POVERTY STATUS⁷									
Below 100% FPL	57%	(±3%)	358,000	20%	(±5%)	78,000	65%	(±8%)	59,000
Between 100–200% FPL	67%	(±3%)	550,000	22%	(±3%)	174,000	72%	(±5%)	208,000
Above 200% FPL	77%	(±2%)	1,401,000	32%	(±2%)	645,000	79%	(±3%)	529,000
AGE GROUP									
18–29 years	61%	(±3%)	502,000	20%	(±3%)	182,000			
30–39 years	66%	(±3%)	537,000	24%	(±3%)	213,000			
40–49 years	75%	(±3%)	457,000	27%	(±4%)	153,000			
50 years and older	78%	(±2%)	813,000	42%	(±3%)	350,000			
EDUCATION STATUS									
Less than high school	60%	(±3%)	449,000	20%	(±4%)	108,000	71%	(±5%)	145,000
High school	66%	(±3%)	473,000	21%	(±3%)	144,000	76%	(±5%)	183,000
Some college or trade school	74%	(±2%)	650,000	31%	(±3%)	261,000	76%	(±4%)	228,000
Graduate or post graduate	80%	(±2%)	727,000	34%	(±3%)	381,000	81%	(±4%)	235,000

1. The percentages and numbers reported reflect the best estimates of the respondents in the Los Angeles County. These estimates should not be considered exact population numbers but should be used as trend data for planning purposes.

2. Men 40 years and older.

3. Women 18 years and older.

4. Men 18 years and older.

5. Women 50 years and older.

6. The actual percentages and numbers may be lower or higher based on the confidence interval reported. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

7. Poverty status is determined based on federal poverty level (FPL). In 1997, FPL for a family of four was \$16,050 a year and 200% of poverty was \$32,100.

Appendix C-1: Antelope Valley

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			ANTELOPE VALLEY SPA		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors						
ALCOHOL USE³-ADULTS⁴						
Non-drinker	43%	(±1%)	2,770,000	39%	(±8%)	69,000
Light	38%	(±1%)	2,440,000	42%	(±8%)	74,000
Moderate	14%	(±1%)	927,000	17%	(±6%)	30,000
Heavy	5%	(±1%)	331,000	1%	(±2%)	1,000
SEAT BELT NOT USED REGULARLY						
Adults	11%	(±1%)	737,000	7%	(±4%)	12,000
Children ⁵	5%	(±1%)	129,000	3%	(±5%)	3,000
CIGARETTE SMOKING						
Adults	18%	(±1%)	1,210,000	21%	(±7%)	38,000
NUMBER OF PACKS SMOKED, ADULTS						
One pack or fewer	94%	(±1%)	1,131,000	97%	(±6%)	37,000
More than one pack	6%	(±1%)	78,000	3%	(±6%)	1,000
HANDGUN OWNERSHIP						
Adults	20%	(±1%)	1,332,000	34%	(±8%)	60,000
Health Status						
WEIGHT STATUS⁶						
Not overweight	51%	(±1%)	3,207,000	51%	(±9%)	89,000
Mild or moderate	34%	(±1%)	2,154,000	36%	(±8%)	63,000
Severely	14%	(±1%)	894,000	13%	(±6%)	22,000
SELF-PERCEIVED HEALTH STATUS, ADULTS						
Very good or excellent	52%	(±1%)	3,448,000	65%	(±8%)	116,000
Good	27%	(±1%)	1,807,000	21%	(±7%)	38,000
Poor or fair	21%	(±1%)	1,359,000	14%	(±6%)	25,000
MEDICAL CONDITION, ADULTS						
Arthritis	18%	(±1%)	1,151,000	21%	(±7%)	38,000
Diabetes	6%	(±1%)	371,000	5%	(±4%)	10,000
BP or hypertension	16%	(±1%)	1,041,000	10%	(±5%)	18,000
MEDICAL CONDITION, CHILDREN						
Asthma	7%	(±1%)	206,000	9%	(±8%)	9,000

Appendix C-1: Antelope Valley Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			ANTELOPE VALLEY SPA		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services						
INSURANCE STATUS, ADULTS						
Medi-Cal	11%	(±1%)	705,000	8%	(±5%)	15,000
Medicare	9%	(±1%)	569,000	7%	(±4%)	13,000
Private	50%	(±1%)	3,279,000	58%	(±8%)	104,000
Uninsured	31%	(±1%)	2,035,000	27%	(±7%)	48,000
INSURANCE STATUS, CHILDREN						
Medi-Cal	31%	(±2%)	851,000	23%	(±13%)	24,000
Private	44%	(±3%)	1,229,000	58%	(±15%)	60,000
Uninsured	25%	(±2%)	696,000	19%	(±12%)	20,000
REGULAR SOURCE OF HEALTH CARE, ADULTS						
Private MD/HMO	56%	(±1%)	3,628,000	69%	(±8%)	123,000
Other regular provider	12%	(±1%)	804,000	11%	(±5%)	19,000
No regular provider	26%	(±1%)	1,656,000	15%	(±6%)	26,000
County or community clinics	6%	(±1%)	415,000	5%	(±4%)	9,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN						
Private MD/HMO	60%	(±3%)	1,678,000	70%	(±14%)	73,000
Other regular provider	10%	(±2%)	289,000	5%	(±6%)	5,000
No regular provider	15%	(±2%)	423,000	18%	(±11%)	19,000
County or community clinics	14%	(±2%)	394,000	7%	(±8%)	8,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS						
Somewhat or very difficult	35%	(±1%)	2,228,000	31%	(±8%)	54,000
Somewhat or very easy	65%	(±1%)	4,238,000	69%	(±8%)	121,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS						
Prescription medicines	16%	(±1%)	1,049,000	16%	(±1%)	29,000
Mental health care	8%	(±1%)	522,000	5%	(±1%)	8,000
Dental care	27%	(±1%)	1,753,000	26%	(±1%)	47,000
Vision care	15%	(±1%)	1,012,000	12%	(±1%)	22,000
Doctor's visit	18%	(±1%)	1,193,000	12%	(±1%)	22,000
PREVENTIVE SERVICES, ADULTS						
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	34%	(±8%)	60,000
Prostate exam ⁷	51%	(±3%)	759,000	49%	(±18%)	22,000
Pap test ⁸	73%	(±1%)	2,387,000	74%	(±9%)	70,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	74%	(±9%)	71,000
Testicular exam ⁹	28%	(±2%)	897,000	29%	(±12%)	24,000
Mammogram ¹⁰	76%	(±2%)	796,000	62%	(±21%)	14,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.
2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.
3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).
4. Adults aged 18 years and older.
5. Children aged 0 to 17 years.
6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.
7. Men aged 40 years and older.
8. Women aged 18 years and older.
9. Men aged 18 years and older.
10. Women aged 50 years and older.

Appendix C-2: San Fernando Valley

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SAN FERNANDO VALLEY SPA			EAST VALLEY HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	43%	(±1%)	2,770,000	39%	(±3%)	501,000	44%	(±6%)	122,000
Light	38%	(±1%)	2,440,000	40%	(±3%)	517,000	37%	(±5%)	103,000
Moderate	14%	(±1%)	927,000	17%	(±2%)	219,000	17%	(±4%)	46,000
Heavy	5%	(±1%)	331,000	5%	(±1%)	59,000	3%	(±2%)	8,000
SEAT BELT NOT USED REGULARLY									
Adults	11%	(±1%)	737,000	12%	(±2%)	162,000	15%	(±4%)	42,000
Children ⁵	5%	(±1%)	129,000	3%	(±2%)	17,000	4%	(±5%)	5,000
CIGARETTE SMOKING									
Adults	18%	(±1%)	1,210,000	19%	(±2%)	250,000	20%	(±4%)	57,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	94%	(±1%)	1,131,000	91%	(±3%)	228,000	89%	(±8%)	51,000
More than one pack	6%	(±1%)	78,000	9%	(±3%)	22,000	11%	(±8%)	6,000
HANDGUN OWNERSHIP									
Adults	20%	(±1%)	1,332,000	24%	(±2%)	317,000	19%	(±4%)	53,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	51%	(±1%)	3,207,000	54%	(±3%)	694,000	57%	(±6%)	155,000
Mild or moderate	34%	(±1%)	2,154,000	34%	(±2%)	436,000	32%	(±5%)	88,000
Severely	14%	(±1%)	894,000	11%	(±2%)	146,000	11%	(±4%)	30,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	52%	(±1%)	3,448,000	57%	(±3%)	753,000	56%	(±5%)	159,000
Good	27%	(±1%)	1,807,000	29%	(±2%)	378,000	29%	(±5%)	83,000
Poor or fair	21%	(±1%)	1,359,000	15%	(±2%)	195,000	15%	(±4%)	43,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±1%)	1,151,000	16%	(±2%)	214,000	11%	(±4%)	32,000
Diabetes	6%	(±1%)	371,000	5%	(±1%)	65,000	5%	(±4%)	13,000
BP or hypertension	16%	(±1%)	1,041,000	15%	(±2%)	192,000	12%	(±2%)	33,000
MEDICAL CONDITION, CHILDREN									
Asthma	7%	(±1%)	206,000	4%	(±2%)	22,000	3%	(±4%)	3,000

Appendix C-2: San Fernando Valley Continued

Summary of Results by Service Planning Area, 1997¹

	GLENDALE HD			SAN FERNANDO HD			WEST VALLEY HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	42%	(±6%)	101,000	35%	(±6%)	85,000	36%	(±4%)	193,000
Light	37%	(±6%)	89,000	43%	(±6%)	105,000	41%	(±4%)	221,000
Moderate	18%	(±5%)	43,000	16%	(±4%)	39,000	17%	(±3%)	91,000
Heavy	3%	(±2%)	6,000	6%	(±3%)	15,000	6%	(±2%)	31,000
SEAT BELT NOT USED REGULARLY									
Adults	13%	(±4%)	32,000	9%	(±3%)	23,000	12%	(±2%)	65,000
Children ⁵	1%	(±3%)	1,000	5%	(±5%)	5,000	3%	(±3%)	6,000
CIGARETTE SMOKING									
Adults	18%	(±5%)	45,000	18%	(±5%)	45,000	19%	(±5%)	102,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	89%	(±10%)	40,000	91%	(±8%)	41,000	94%	(±4%)	96,000
More than one pack	11%	(±10%)	5,000	9%	(±8%)	4,000	6%	(±4%)	6,000
HANDGUN OWNERSHIP									
Adults	23%	(±5%)	54,000	36%	(±6%)	88,000	22%	(±3%)	121,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	55%	(±6%)	130,000	50%	(±6%)	119,000	55%	(±4%)	290,000
Mild or moderate	33%	(±6%)	79,000	40%	(±6%)	93,000	33%	(±4%)	176,000
Severely	12%	(±4%)	29,000	10%	(±4%)	24,000	12%	(±3%)	62,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Poor or fair	53%	(±6%)	128,000	59%	(±6%)	147,000	58%	(±4%)	319,000
Good	32%	(±6%)	78,000	26%	(±5%)	64,000	28%	(±3%)	153,000
Very good or excellent	15%	(±5%)	37,000	15%	(±4%)	38,000	14%	(±3%)	75,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±5%)	45,000	15%	(±4%)	37,000	18%	(±3%)	101,000
Diabetes	4%	(±4%)	10,000	6%	(±4%)	14,000	5%	(±3%)	28,000
BP or hypertension	14%	(±3%)	33,000	12%	(±3%)	30,000	18%	(±2%)	96,000
MEDICAL CONDITION, CHILDREN									
Asthma	3%	(±5%)	2,000	5%	(±5%)	6,000	6%	(±4%)	11,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI < 25 then the person is classified as not overweight, 25 ≤ BMI < 30 as mild or moderately overweight, and BMI ≥ 30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-2: San Fernando Valley Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SAN FERNANDO VALLEY SPA			EAST VALLEY HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	11%	(±1%)	705,000	8%	(±1%)	100,000	7%	(±3%)	21,000
Medicare	9%	(±1%)	569,000	9%	(±1%)	120,000	6%	(±3%)	17,000
Private	50%	(±1%)	3,279,000	55%	(±3%)	726,000	46%	(±5%)	130,000
Uninsured	31%	(±1%)	2,035,000	29%	(±2%)	379,000	41%	(±5%)	117,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	31%	(±2%)	851,000	25%	(±5%)	117,000	29%	(±11%)	33,000
Private	44%	(±3%)	1,229,000	50%	(±6%)	236,000	47%	(±12%)	53,000
Uninsured	25%	(±2%)	696,000	26%	(±5%)	122,000	25%	(±10%)	28,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	56%	(±1%)	3,628,000	60%	(±3%)	782,000	52%	(±6%)	147,000
Other regular provider	12%	(±1%)	804,000	11%	(±2%)	147,000	12%	(±4%)	34,000
No regular provider	26%	(±1%)	1,656,000	24%	(±2%)	311,000	27%	(±5%)	76,000
County or community clinics	6%	(±1%)	415,000	5%	(±1%)	70,000	9%	(±3%)	24,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	60%	(±3%)	1,678,000	68%	(±5%)	323,000	63%	(±12%)	72,000
Other regular provider	10%	(±2%)	289,000	10%	(±3%)	46,000	11%	(±8%)	12,000
No regular provider	15%	(±2%)	423,000	13%	(±4%)	62,000	17%	(±9%)	19,000
County or community clinics	14%	(±2%)	394,000	10%	(±3%)	46,000	9%	(±7%)	10,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	35%	(±1%)	2,228,000	33%	(±2%)	426,000	41%	(±5%)	113,000
Somewhat or very easy	65%	(±1%)	4,238,000	67%	(±2%)	870,000	59%	(±5%)	162,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	16%	(±1%)	1,049,000	15%	(±2%)	197,000	18%	(±7%)	52,000
Mental health care	8%	(±1%)	522,000	8%	(±3%)	108,000	10%	(±3%)	27,000
Dental care	27%	(±1%)	1,753,000	27%	(±2%)	340,000	30%	(±5%)	86,000
Vision care	15%	(±1%)	1,012,000	15%	(±2%)	201,000	16%	(±4%)	45,000
Doctor's visit	18%	(±1%)	1,193,000	16%	(±2%)	216,000	21%	(±4%)	60,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	37%	(±2%)	481,000	38%	(±5%)	105,000
Prostate exam ⁷	51%	(±3%)	759,000	56%	(±5%)	182,000	42%	(±12%)	25,000
Pap test ⁸	73%	(±1%)	2,387,000	74%	(±3%)	459,000	68%	(±6%)	91,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	72%	(±3%)	448,000	67%	(±7%)	89,000
Testicular exam ⁹	28%	(±2%)	897,000	30%	(±4%)	203,000	23%	(±7%)	33,000
Mammogram ¹⁰	76%	(±2%)	796,000	74%	(±5%)	144,000	70%	(±12%)	26,000

Appendix C-2: San Fernando Valley Continued

Summary of Results by Service Planning Area, 1997¹

	GLENDALE HD			SAN FERNANDO HD			WEST VALLEY HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	7%	(±3%)	17,000	5%	(±3%)	13,000	9%	(±2%)	50,000
Medicare	11%	(±4%)	28,000	8%	(±3%)	19,000	10%	(±2%)	56,000
Private	59%	(±6%)	144,000	63%	(±6%)	157,000	54%	(±4%)	296,000
Uninsured	23%	(±5%)	57,000	24%	(±5%)	61,000	27%	(±3%)	146,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	23%	(±13%)	14,000	18%	(±9%)	20,000	27%	(±8%)	16,000
Private	60%	(±15%)	38,000	54%	(±12%)	60,000	45%	(±9%)	86,000
Uninsured	17%	(±12%)	11,000	28%	(±11%)	32,000	28%	(±8%)	15,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	65%	(±6%)	158,000	63%	(±6%)	156,000	59%	(±4%)	321,000
Other regular provider	11%	(±4%)	27,000	10%	(±4%)	25,000	11%	(±2%)	61,000
No regular provider	19%	(±5%)	47,000	22%	(±5%)	54,000	25%	(±3%)	134,000
County or community clinics	5%	(±3%)	11,000	5%	(±3%)	12,000	4%	(±2%)	23,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	76%	(±13%)	49,000	65%	(±11%)	73,000	69%	(±8%)	129,000
Other regular provider	4%	(±6%)	3,000	4%	(±5%)	5,000	14%	(±6%)	26,000
No regular provider	8%	(±8%)	5,000	15%	(±9%)	17,000	11%	(±5%)	21,000
County or community clinics	12%	(±10%)	7,000	15%	(±8%)	17,000	6%	(±4%)	12,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	31%	(±6%)	75,000	30%	(±5%)	73,000	31%	(±4%)	165,000
Somewhat or very easy	69%	(±6%)	166,000	70%	(±5%)	175,000	69%	(±4%)	367,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	11%	(±4%)	28,000	13%	(±4%)	33,000	15%	(±3%)	85,000
Mental health care	9%	(±4%)	20,000	6%	(±3%)	16,000	8%	(±2%)	45,000
Dental care	21%	(±5%)	52,000	23%	(±5%)	57,000	27%	(±3%)	145,000
Vision care	15%	(±5%)	35,000	13%	(±4%)	33,000	16%	(±3%)	87,000
Doctor's visit	14%	(±4%)	34,000	13%	(±4%)	32,000	17%	(±3%)	91,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	33%	(±6%)	78,000	34%	(±6%)	85,000	40%	(±4%)	213,000
Prostate exam ⁷	53%	(±13%)	35,000	40%	(±8%)	27,000	62%	(±8%)	82,000
Pap test ⁸	75%	(±7%)	83,000	80%	(±6%)	94,000	74%	(±4%)	192,000
Clinical breast exam ⁸	72%	(±7%)	81,000	78%	(±6%)	91,000	73%	(±4%)	188,000
Testicular exam ⁹	34%	(±9%)	43,000	30%	(±8%)	40,000	31%	(±5%)	86,000
Mammogram ¹⁰	75%	(±12%)	28,000	78%	(±11%)	28,000	73%	(±8%)	63,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI < 25 then the person is classified as not overweight, 25 = < BMI < 30 as mild or moderately overweight, and BMI > = 30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-3: San Gabriel Valley

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SAN GABRIEL VALLEY SPA			ALHAMBRA HD			EL MONTE HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors												
ALCOHOL USE ³ , ADULTS ⁴												
Non-drinker	43%	(±1%)	2,770,000	44%	(±3%)	544,000	44%	(±6%)	117,000	52%	(±6%)	161,000
Light	38%	(±1%)	2,440,000	38%	(±3%)	471,000	37%	(±6%)	99,000	31%	(±6%)	97,000
Moderate	14%	(±1%)	927,000	12%	(±2%)	153,000	13%	(±4%)	35,000	12%	(±4%)	36,000
Heavy	5%	(±1%)	331,000	5%	(±1%)	67,000	6%	(±3%)	16,000	5%	(±3%)	16,000
SEAT BELT NOT USED REGULARLY												
Adults	11%	(±1%)	737,000	9%	(±2%)	116,000	7%	(±3%)	19,000	9%	(±3%)	28,000
Children ⁵	5%	(±1%)	129,000	4%	(±2%)	25,000	5%	(±5%)	6,000	6%	(±5%)	12,000
CIGARETTE SMOKING												
Adults	18%	(±1%)	1,210,000	19%	(±2%)	235,000	19%	(±5%)	51,000	17%	(±5%)	54,000
NUMBER OF PACKS SMOKED, ADULTS												
One pack or fewer	94%	(±1%)	1,131,000	91%	(±4%)	214,000	95%	(±6%)	48,000	94%	(±7%)	51,000
More than one pack	6%	(±1%)	78,000	9%	(±4%)	21,000	5%	(±6%)	3,000	6%	(±7%)	3,000
HANDGUN OWNERSHIP												
Adults	20%	(±1%)	1,332,000	23%	(±2%)	285,000	20%	(±5%)	53,000	16%	(±4%)	51,000
Health Status												
WEIGHT STATUS ⁶												
Not overweight	51%	(±1%)	3,207,000	51%	(±3%)	611,000	63%	(±6%)	166,000	41%	(±6%)	118,000
Mild or moderate	34%	(±1%)	2,154,000	35%	(±3%)	418,000	29%	(±6%)	75,000	41%	(±6%)	121,000
Severely	14%	(±1%)	894,000	14%	(±2%)	169,000	8%	(±3%)	21,000	18%	(±5%)	53,000
SELF-PERCEIVED HEALTH STATUS, ADULTS												
Very good or excellent	52%	(±1%)	3,448,000	51%	(±3%)	643,000	53%	(±6%)	143,000	40%	(±6%)	125,000
Good	27%	(±1%)	1,807,000	27%	(±2%)	336,000	26%	(±5%)	69,000	31%	(±6%)	96,000
Poor or fair	21%	(±1%)	1,359,000	22%	(±2%)	280,000	22%	(±5%)	60,000	29%	(±5%)	90,000
MEDICAL CONDITION, ADULTS												
Arthritis	18%	(±1%)	1,151,000	20%	(±2%)	246,000	17%	(±5%)	46,000	18%	(±5%)	56,000
Diabetes	6%	(±1%)	371,000	6%	(±1%)	81,000	4%	(±2%)	11,000	8%	(±3%)	24,000
BP or hypertension	16%	(±1%)	1,041,000	18%	(±2%)	221,000	15%	(±4%)	40,000	17%	(±5%)	53,000
MEDICAL CONDITION, CHILDREN												
Asthma	7%	(±1%)	206,000	7%	(±3%)	39,000	7%	(±6%)	8,000	9%	(±6%)	18,000

Appendix C-3: San Gabriel Valley Continued

Summary of Results by Service Planning Area, 1997¹

	FOOTHILL HD			PASADENA HD			POMONA HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	36%	(±6%)	74,000	36%	(±8%)	34,000	44%	(±5%)	157,000
Light	46%	(±6%)	94,000	42%	(±8%)	40,000	40%	(±5%)	141,000
Moderate	12%	(±4%)	25,000	14%	(±6%)	13,000	12%	(±3%)	43,000
Heavy	6%	(±3%)	12,000	8%	(±4%)	8,000	4%	(±2%)	15,000
SEAT BELT NOT USED REGULARLY									
Adults	9%	(±4%)	20,000	15%	(±6%)	15,000	9%	(±3%)	34,000
Children ⁵	1%	(±3%)	1,000	10%	(±11%)	3,000	2%	(±3%)	3,000
CIGARETTE SMOKING									
Adults	20%	(±5%)	43,000	16%	(±6%)	16,000	20%	(±4%)	71,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	87%	(±10%)	37,000	93%	(±10%)	15,000	88%	(±8%)	63,000
More than one pack	13%	(±10%)	6,000	7%	(±10%)	1,000	12%	(±8%)	9,000
HANDGUN OWNERSHIP									
Adults	29%	(±6%)	62,000	22%	(±7%)	22,000	27%	(±5%)	97,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	51%	(±6%)	103,000	51%	(±8%)	49,000	51%	(±5%)	175,000
Mild or moderate	31%	(±6%)	63,000	36%	(±8%)	35,000	36%	(±5%)	123,000
Severely	18%	(±5%)	35,000	13%	(±5%)	12,000	14%	(±4%)	47,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	52%	(±6%)	110,000	67%	(±7%)	66,000	55%	(±5%)	199,000
Good	27%	(±6%)	58,000	17%	(±6%)	17,000	26%	(±5%)	95,000
Poor or fair	21%	(±5%)	44,000	17%	(±6%)	17,000	19%	(±4%)	70,000
MEDICAL CONDITION, ADULTS									
Arthritis	25%	(±5%)	53,000	17%	(±6%)	17,000	21%	(±4%)	75,000
Diabetes	9%	(±4%)	20,000	5%	(±4%)	5,000	6%	(±3%)	21,000
BP or hypertension	20%	(±5%)	43,000	18%	(±6%)	18,000	18%	(±4%)	67,000
MEDICAL CONDITION, CHILDREN									
Asthma	4%	(±5%)	3,000	3%	(±7%)	1,000	6%	(±5%)	10,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.
2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.
3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).
4. Adults aged 18 years and older.
5. Children aged 0 to 17 years.
6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.
7. Men aged 40 years and older.
8. Women aged 18 years and older.
9. Men aged 18 years and older.
10. Women aged 50 years and older.

Appendix C-3: San Gabriel Valley Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SAN GABRIEL VALLEY SPA			ALHAMBRA HD			EL MONTE HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services												
INSURANCE STATUS, ADULTS												
Medi-Cal	11%	(±1%)	705,000	10%	(±2%)	124,000	9%	(±3%)	23,000	14%	(±4%)	44,000
Medicare	9%	(±1%)	569,000	10%	(±2%)	129,000	11%	(±4%)	29,000	6%	(±3%)	19,000
Private	50%	(±1%)	3,279,000	52%	(±3%)	655,000	48%	(±6%)	131,000	46%	(±6%)	142,000
Uninsured	31%	(±1%)	2,035,000	27%	(±2%)	344,000	32%	(±6%)	87,000	34%	(±6%)	105,000
INSURANCE STATUS, CHILDREN												
Medi-Cal	31%	(±2%)	851,000	23%	(±5%)	132,000	19%	(±9%)	21,000	31%	(±10%)	61,000
Private	44%	(±3%)	1,229,000	56%	(±6%)	317,000	63%	(±11%)	70,000	40%	(±11%)	78,000
Uninsured	25%	(±2%)	696,000	21%	(±5%)	121,000	18%	(±9%)	20,000	28%	(±10%)	55,000
REGULAR SOURCE OF HEALTH CARE, ADULTS												
Private MD/HMO	56%	(±1%)	3,628,000	60%	(±3%)	733,000	62%	(±6%)	161,000	49%	(±6%)	148,000
Other regular provider	12%	(±1%)	804,000	12%	(±2%)	144,000	9%	(±4%)	24,000	12%	(±3%)	36,000
No regular provider	26%	(±1%)	1,656,000	24%	(±2%)	290,000	27%	(±5%)	69,000	30%	(±5%)	92,000
County or community clinics	6%	(±1%)	415,000	5%	(±1%)	63,000	2%	(±2%)	6,000	10%	(±3%)	29,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN												
Private MD/HMO	60%	(±3%)	1,678,000	68%	(±5%)	388,000	75%	(±10%)	84,000	61%	(±11%)	115,000
Other regular provider	10%	(±2%)	289,000	9%	(±3%)	51,000	6%	(±6%)	7,000	11%	(±7%)	20,000
No regular provider	15%	(±2%)	423,000	13%	(±4%)	75,000	13%	(±8%)	15,000	15%	(±8%)	27,000
County or community clinics	14%	(±2%)	394,000	10%	(±3%)	56,000	6%	(±6%)	7,000	13%	(±8%)	25,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS												
Somewhat or very difficult	35%	(±1%)	2,228,000	32%	(±3%)	397,000	31%	(±6%)	78,000	38%	(±6%)	117,000
Somewhat or very easy	65%	(±1%)	4,238,000	68%	(±3%)	834,000	69%	(±6%)	177,000	62%	(±6%)	191,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS												
Prescription medicines	16%	(±1%)	1,049,000	16%	(±2%)	195,000	16%	(±4%)	44,000	20%	(±5%)	62,000
Mental health care	8%	(±1%)	522,000	7%	(±3%)	91,000	7%	(±3%)	18,000	9%	(±3%)	28,000
Dental care	27%	(±1%)	1,753,000	27%	(±2%)	333,000	28%	(±5%)	75,000	35%	(±6%)	110,000
Vision care	15%	(±1%)	1,012,000	15%	(±2%)	193,000	13%	(±4%)	36,000	21%	(±5%)	66,000
Doctor's visit	18%	(±1%)	1,193,000	18%	(±2%)	220,000	17%	(±5%)	48,000	21%	(±5%)	66,000
PREVENTIVE SERVICES, ADULTS												
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	29%	(±3%)	357,000	23%	(±5%)	60,000	34%	(±6%)	106,000
Prostate exam ⁷	51%	(±3%)	759,000	50%	(±6%)	157,000	45%	(±12%)	31,000	42%	(±14%)	27,000
Pap test ⁸	73%	(±1%)	2,387,000	73%	(±3%)	459,000	68%	(±7%)	82,000	75%	(±6%)	120,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	71%	(±3%)	448,000	71%	(±7%)	86,000	70%	(±7%)	112,000
Testicular exam ⁹	28%	(±2%)	897,000	26%	(±4%)	158,000	21%	(±7%)	29,000	23%	(±8%)	34,000
Mammogram ¹⁰	76%	(±2%)	796,000	82%	(±4%)	183,000	82%	(±10%)	36,000	88%	(±9%)	39,000

Appendix C-3: San Gabriel Valley Continued

Summary of Results by Service Planning Area, 1997¹

	FOOTHILL HD			PASADENA HD			POMONA HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	10%	(±4%)	20,000	6%	(±4%)	6,000	9%	(±3%)	31,000
Medicare	12%	(±4%)	25,000	14%	(±5%)	14,000	11%	(±3%)	42,000
Private	56%	(±6%)	117,000	49%	(±8%)	48,000	60%	(±5%)	218,000
Uninsured	23%	(±5%)	47,000	32%	(±7%)	31,000	20%	(±4%)	73,000
Medi-Cal	24%	(±12%)	19,000	29%	(±18%)	7,000	15%	(±7%)	24,000
Private	53%	(±14%)	41,000	48%	(±19%)	11,000	71%	(±9%)	117,000
Uninsured	23%	(±11%)	17,000	23%	(±17%)	6,000	15%	(±7%)	24,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	67%	(±6%)	139,000	58%	(±9%)	56,000	63%	(±5%)	228,000
Other regular provider	8%	(±4%)	16,000	14%	(±6%)	14,000	15%	(±4%)	54,000
No regular provider	21%	(±6%)	43,000	25%	(±7%)	24,000	17%	(±4%)	61,000
County or community clinics	4%	(±4%)	8,000	3%	(±3%)	3,000	4%	(±2%)	16,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	82%	(±11%)	62,000	58%	(±19%)	16,000	66%	(±10%)	112,000
Other regular provider	5%	(±6%)	4,000	18%	(±15%)	5,000	10%	(±6%)	16,000
No regular provider	14%	(±9%)	10,000	12%	(±12%)	3,000	12%	(±7%)	20,000
County or community clinics	0%		0	12%	(±12%)	3,000	12%	(±7%)	21,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	28%	(±6%)	58,000	29%	(±7%)	29,000	32%	(±5%)	116,000
Somewhat or very easy	72%	(±6%)	150,000	71%	(±7%)	70,000	68%	(±5%)	247,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	13%	(±4%)	27,000	12%	(±5%)	11,000	14%	(±4%)	51,000
Mental health care	8%	(±4%)	17,000	8%	(±4%)	8,000	5%	(±2%)	20,000
Dental care	23%	(±5%)	49,000	18%	(±6%)	18,000	22%	(±4%)	81,000
Vision care	15%	(±5%)	31,000	10%	(±5%)	10,000	14%	(±4%)	51,000
Doctor's visit	19%	(±5%)	40,000	9%	(±5%)	9,000	16%	(±4%)	58,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	30%	(±6%)	61,000	26%	(±7%)	25,000	30%	(±5%)	105,000
Prostate exam ⁷	62%	(±12%)	36,000	46%	(±17%)	11,000	52%	(±10%)	52,000
Pap test ⁸	73%	(±7%)	83,000	77%	(±9%)	36,000	74%	(±6%)	138,000
Clinical breast exam ⁸	71%	(±7%)	81,000	73%	(±9%)	34,000	73%	(±6%)	135,000
Testicular exam ⁹	34%	(±10%)	31,000	29%	(±11%)	15,000	29%	(±8%)	48,000
Mammogram ¹⁰	79%	(±11%)	32,000	79%	(±13%)	16,000	81%	(±8%)	60,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.
2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.
3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).
4. Adults aged 18 years and older.
5. Children aged 0 to 17 years.
6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.
7. Men aged 40 years and older.
8. Women aged 18 years and older.
9. Men aged 18 years and older.
10. Women aged 50 years and older.

Appendix C-4: Metro

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			METRO SPA			CENTRAL HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	43%	(±1%)	2,770,000	49%	(±3%)	402,000	60%	(±7%)	136,000
Light	38%	(±1%)	2,440,000	33%	(±3%)	270,000	28%	(±6%)	64,000
Moderate	14%	(±1%)	927,000	12%	(±2%)	96,000	5%	(±3%)	11,000
Heavy	5%	(±1%)	331,000	6%	(±2%)	47,000	7%	(±3%)	15,000
SEAT BELT NOT USED REGULARLY									
Adults	11%	(±1%)	737,000	12%	(±2%)	97,000	11%	(±4%)	26,000
Children ⁵	5%	(±1%)	129,000	6%	(±4%)	19,000	2%	(±4%)	2,000
CIGARETTE SMOKING									
Adults	18%	(±1%)	1,210,000	19%	(±3%)	157,000	16%	(±5%)	38,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	94%	(±1%)	1,131,000	98%	(±2%)	153,000	99%	(±4%)	37,000
More than one pack	6%	(±1%)	78,000	2%	(±2%)	4,000	1%	(±4%)	<1,000
HANDGUN OWNERSHIP									
Adults	20%	(±1%)	1,332,000	10%	(±2%)	83,000	9%	(±4%)	20,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	51%	(±1%)	3,207,000	57%	(±4%)	443,000	57%	(±7%)	122,000
Mild or moderate	34%	(±1%)	2,154,000	30%	(±3%)	231,000	29%	(±7%)	62,000
Severely	14%	(±1%)	894,000	13%	(±2%)	100,000	14%	(±5%)	29,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	52%	(±1%)	3,448,000	47%	(±3%)	391,000	39%	(±7%)	89,000
Good	27%	(±1%)	1,807,000	28%	(±3%)	233,000	29%	(±6%)	66,000
Poor or fair	21%	(±1%)	1,359,000	25%	(±3%)	207,000	33%	(±6%)	74,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±1%)	1,151,000	14%	(±2%)	115,000	14%	(±5%)	32,000
Diabetes	6%	(±1%)	371,000	6%	(±2%)	48,000	8%	(±4%)	17,000
BP or hypertension	16%	(±1%)	1,041,000	14%	(±2%)	114,000	15%	(±5%)	33,000
MEDICAL CONDITION, CHILDREN									
Asthma	7%	(±1%)	206,000	5%	(±3%)	16,000	7%	(±7%)	6,000

Appendix C-4: Metro Continued

Summary of Results by Service Planning Area, 1997¹

	HOLLYWOOD HD			NORTHEAST HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors						
ALCOHOL USE ³ , ADULTS ⁴						
Non-drinker	40%	(±5%)	145,000	53%	(±7%)	121,000
Light	38%	(±5%)	138,000	30%	(±6%)	69,000
Moderate	17%	(±4%)	60,000	11%	(±4%)	25,000
Heavy	5%	(±2%)	19,000	6%	(±3%)	13,000
SEAT BELT NOT USED REGULARLY						
Adults	12%	(±3%)	45,000	11%	(±4%)	26,000
Children ⁵	12%	(±8%)	11,000	4%	(±5%)	6,000
CIGARETTE SMOKING						
Adults	19%	(±4%)	71,000	21%	(±6%)	48,000
NUMBER OF PACKS SMOKED, ADULTS						
One pack or fewer	98%	(±4%)	69,000	97%	(±5%)	47,000
More than one pack	2%	(±4%)	2,000	3%	(±5%)	1,000
HANDGUN OWNERSHIP						
Adults	10%	(±3%)	35,000	12%	(±5%)	28,000
Health Status						
WEIGHT STATUS ⁶						
Not overweight	64%	(±5%)	219,000	47%	(±7%)	102,000
Mild or moderate	26%	(±4%)	91,000	36%	(±7%)	78,000
Severely	10%	(±3%)	35,000	17%	(±5%)	36,000
SELF-PERCEIVED HEALTH STATUS, ADULTS						
Very good or excellent	57%	(±5%)	211,000	39%	(±7%)	91,000
Good	24%	(±4%)	90,000	33%	(±6%)	77,000
Poor or fair	18%	(±4%)	68,000	28%	(±6%)	64,000
MEDICAL CONDITION, ADULTS						
Arthritis	13%	(±3%)	48,000	15%	(±5%)	35,000
Diabetes	5%	(±2%)	19,000	5%	(±3%)	12,000
BP or hypertension	14%	(±3%)	52,000	12%	(±5%)	28,000
MEDICAL CONDITION, CHILDREN						
Asthma	5%	(±6%)	5,000	4%	(±5%)	5,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-4: Metro Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			METRO SPA			CENTRAL HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
<i>Access to Health Care and Utilization of Health Care Services</i>									
INSURANCE STATUS, ADULTS									
Medi-Cal	11%	(±1%)	705,000	13%	(±2%)	111,000	15%	(±5%)	34,000
Medicare	9%	(±1%)	569,000	6%	(±2%)	49,000	6%	(±3%)	13,000
Private	50%	(±1%)	3,279,000	37%	(±3%)	304,000	29%	(±6%)	66,000
Uninsured	31%	(±1%)	2,035,000	44%	(±3%)	360,000	50%	(±7%)	114,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	31%	(±2%)	851,000	46%	(±8%)	145,000	46%	(±15%)	42,000
Private	44%	(±3%)	1,229,000	22%	(±7%)	71,000	21%	(±12%)	19,000
Uninsured	25%	(±2%)	696,000	32%	(±7%)	100,000	34%	(±14%)	31,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	56%	(±1%)	3,628,000	42%	(±3%)	340,000	31%	(±6%)	70,000
Other regular provider	12%	(±1%)	804,000	13%	(±2%)	107,000	15%	(±5%)	33,000
No regular provider	26%	(±1%)	1,656,000	37%	(±3%)	302,000	43%	(±7%)	96,000
County or community clinics	6%	(±1%)	415,000	8%	(±2%)	67,000	11%	(±4%)	24,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	60%	(±3%)	1,678,000	45%	(±8%)	145,000	53%	(±15%)	49,000
Other regular provider	10%	(±2%)	289,000	10%	(±5%)	34,000	7%	(±8%)	7,000
No regular provider	15%	(±2%)	423,000	17%	(±6%)	55,000	17%	(±11%)	15,000
County or community clinics	14%	(±2%)	394,000	27%	(±7%)	88,000	22%	(±12%)	20,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	35%	(±1%)	2,228,000	45%	(±3%)	367,000	49%	(±7%)	107,000
Somewhat or very easy	65%	(±1%)	4,238,000	55%	(±3%)	445,000	51%	(±7%)	114,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	16%	(±1%)	1,049,000	18%	(±2%)	145,000	21%	(±6%)	48,000
Mental health care	8%	(±1%)	522,000	10%	(±2%)	80,000	11%	(±4%)	25,000
Dental care	27%	(±1%)	1,753,000	30%	(±2%)	253,000	33%	(±7%)	76,000
Vision care	15%	(±1%)	1,012,000	16%	(±2%)	136,000	21%	(±6%)	48,000
Doctor's visit	18%	(±1%)	1,193,000	23%	(±2%)	193,000	27%	(±6%)	62,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	41%	(±3%)	340,000	37%	(±7%)	85,000
Prostate exam ⁷	51%	(±3%)	759,000	44%	(±7%)	85,000	38%	(±15%)	20,000
Pap test ⁸	73%	(±1%)	2,387,000	67%	(±4%)	256,000	58%	(±8%)	68,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	59%	(±4%)	228,000	44%	(±8%)	52,000
Testicular exam ⁹	28%	(±2%)	897,000	26%	(±5%)	116,000	26%	(±10%)	30,000
Mammogram ¹⁰	76%	(±2%)	796,000	72%	(±7%)	84,000	62%	(±14%)	25,000

Appendix C-4: Metro Continued

Summary of Results by Service Planning Area, 1997¹

	HOLLYWOOD HD			NORTHEAST HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services						
INSURANCE STATUS, ADULTS						
Medi-Cal	11%	(±3%)	42,000	15%	(±5%)	34,000
Medicare	6%	(±2%)	23,000	5%	(±3%)	12,000
Private	43%	(±5%)	156,000	35%	(±7%)	81,000
Uninsured	39%	(±5%)	144,000	44%	(±7%)	102,000
INSURANCE STATUS, CHILDREN						
Medi-Cal	39%	(±12%)	35,000	51%	(±13%)	68,000
Private	24%	(±11%)	22,000	23%	(±11%)	30,000
Uninsured	37%	(±12%)	34,000	26%	(±11%)	35,000
REGULAR SOURCE OF HEALTH CARE, ADULTS						
Private MD/HMO	48%	(±5%)	175,000	42%	(±7%)	95,000
Other regular provider	12%	(±3%)	44,000	13%	(±5%)	29,000
No regular provider	35%	(±5%)	127,000	35%	(±7%)	78,000
County or community clinics	5%	(±2%)	20,000	10%	(±4%)	23,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN						
Private MD/HMO	55%	(±12%)	51,000	33%	(±13%)	45,000
Other regular provider	5%	(±5%)	4,000	17%	(±10%)	23,000
No regular provider	24%	(±11%)	22,000	12%	(±9%)	17,000
County or community clinics	16%	(±9%)	15,000	38%	(±14%)	52,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS						
Somewhat or very difficult	39%	(±5%)	142,000	52%	(±7%)	118,000
Somewhat or very easy	61%	(±5%)	222,000	48%	(±7%)	109,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS						
Prescription medicines	17%	(±4%)	64,000	14%	(±5%)	33,000
Mental health care	9%	(±3%)	34,000	9%	(±4%)	21,000
Dental care	29%	(±4%)	109,000	30%	(±6%)	69,000
Vision care	15%	(±3%)	57,000	14%	(±5%)	32,000
Doctor's visit	23%	(±4%)	85,000	20%	(±6%)	47,000
PREVENTIVE SERVICES, ADULTS						
HIV/AIDS test ⁴	44%	(±5%)	161,000	41%	(±7%)	95,000
Prostate exam ⁷	50%	(±10%)	43,000	41%	(±14%)	23,000
Pap test ⁸	69%	(±6%)	108,000	74%	(±8%)	80,000
Clinical breast exam ⁸	65%	(±6%)	102,000	68%	(±8%)	74,000
Testicular exam ⁹	29%	(±6%)	60,000	21%	(±8%)	25,000
Mammogram ¹⁰	76%	(±10%)	39,000	80%	(±14%)	21,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-5: West

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			WEST SPA		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors						
ALCOHOL USE ³ , ADULTS ⁴						
Non-drinker	43%	(±1%)	2,770,000	29%	(±8%)	137,000
Light	38%	(±1%)	2,440,000	44%	(±8%)	211,000
Moderate	14%	(±1%)	927,000	23%	(±6%)	111,000
Heavy	5%	(±1%)	331,000	3%	(±2%)	16,000
SEAT BELT NOT USED REGULARLY						
Adults	11%	(±1%)	737,000	12%	(±3%)	58,000
Children ⁵	5%	(±1%)	129,000	6%	(±6%)	5,000
CIGARETTE SMOKING						
Adults	18%	(±1%)	1,210,000	13%	(±3%)	64,000
NUMBER OF PACKS SMOKED, ADULTS						
One pack or fewer	94%	(±1%)	1,131,000	90%	(±7%)	58,000
More than one pack	6%	(±1%)	78,000	10%	(±7%)	6,000
HANDGUN OWNERSHIP						
Adults	20%	(±1%)	1,332,000	18%	(±3%)	84,000
Health Status						
WEIGHT STATUS ⁶						
Not overweight	51%	(±1%)	3,207,000	61%	(±4%)	286,000
Mild or Moderate	34%	(±1%)	2,154,000	30%	(±4%)	140,000
Severely	14%	(±1%)	894,000	9%	(±2%)	40,000
SELF-PERCEIVED HEALTH STATUS, ADULTS						
Very good or excellent	52%	(±1%)	3,448,000	66%	(±4%)	318,000
Good	27%	(±1%)	1,807,000	22%	(±3%)	105,000
Poor or Fair	21%	(±1%)	1,359,000	12%	(±3%)	60,000
MEDICAL CONDITION, ADULTS						
Arthritis	18%	(±1%)	1,151,000	15%	(±3%)	70,000
Diabetes	6%	(±1%)	371,000	5%	(±2%)	23,000
BP or hypertension	16%	(±1%)	1,041,000	13%	(±3%)	63,000
MEDICAL CONDITION, CHILDREN						
Asthma	7%	(±1%)	206,000	14%	(±9%)	12,000

Appendix C-5: West Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			WEST SPA		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services						
INSURANCE STATUS, ADULTS						
Medi-Cal	11%	(±1%)	705,000	5%	(±2%)	23,000
Medicare	9%	(±1%)	569,000	12%	(±3%)	58,000
Private	50%	(±1%)	3,279,000	58%	(±4%)	281,000
Uninsured	31%	(±1%)	2,035,000	25%	(±4%)	120,000
INSURANCE STATUS, CHILDREN						
Medi-Cal	31%	(±2%)	851,000	16%	(±9%)	13,000
Private	44%	(±3%)	1,229,000	53%	(±12%)	44,000
Uninsured	25%	(±2%)	696,000	31%	(±12%)	26,000
REGULAR SOURCE OF HEALTH CARE, ADULTS						
Private MD/HMO	56%	(±1%)	3,628,000	60%	(±4%)	288,000
Other regular provider	12%	(±1%)	804,000	15%	(±3%)	72,000
No regular provider	26%	(±1%)	1,656,000	19%	(±3%)	92,000
County or community clinics	6%	(±1%)	415,000	5%	(±2%)	25,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN						
Private MD/HMO	60%	(±3%)	1,678,000	58%	(±12%)	49,000
Other regular provider	10%	(±2%)	289,000	13%	(±8%)	11,000
No regular provider	15%	(±2%)	423,000	12%	(±8%)	10,000
County or community clinics	14%	(±2%)	394,000	18%	(±9%)	15,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS						
Somewhat or very difficult	35%	(±1%)	2,228,000	27%	(±4%)	126,000
Somewhat or very easy	65%	(±1%)	4,238,000	73%	(±4%)	348,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS						
Prescription medicines	16%	(±1%)	1,049,000	14%	(±1%)	68,000
Mental health care	8%	(±1%)	522,000	11%	(±2%)	50,000
Dental care	27%	(±1%)	1,753,000	22%	(±1%)	108,000
Vision care	15%	(±1%)	1,012,000	13%	(±1%)	63,000
Doctor's visit	18%	(±1%)	1,193,000	16%	(±1%)	77,000
PREVENTIVE SERVICES, ADULTS						
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	38%	(±4%)	180,000
Prostate exam ⁷	51%	(±3%)	759,000	54%	(±9%)	61,000
Pap test ⁸	73%	(±1%)	2,387,000	76%	(±4%)	171,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	81%	(±4%)	183,000
Testicular exam ⁹	28%	(±2%)	897,000	34%	(±6%)	86,000
Mammogram ¹⁰	76%	(±2%)	796,000	78%	(±7%)	75,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.
2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.
3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).
4. Adults aged 18 years and older.
5. Children aged 0 to 17 years.
6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.
7. Men aged 40 years and older.
8. Women aged 18 years and older.
9. Men aged 18 years and older.
10. Women aged 50 years and older.

Appendix C-6: South

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SOUTH SPA			COMPTON HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	43%	(±1%)	2,770,000	62%	(±4%)	369,000	59%	(±6%)	100,000
Light	38%	(±1%)	2,440,000	25%	(±3%)	147,000	30%	(±7%)	50,000
Moderate	14%	(±1%)	927,000	9%	(±2%)	54,000	7%	(±4%)	12,000
Heavy	5%	(±1%)	331,000	5%	(±2%)	29,000	5%	(±3%)	8,000
SEAT BELT NOT USED REGULARLY									
Adults	11%	(±1%)	737,000	12%	(±3%)	74,000	10%	(±5%)	18,000
Children ⁵	5%	(±1%)	129,000	4%	(±3%)	16,000	3%	(±5%)	3,000
CIGARETTE SMOKING									
Adults	18%	(±1%)	1,210,000	19%	(±3%)	115,000	19%	(±6%)	33,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	94%	(±1%)	1,131,000	98%	(±3%)	113,000	97%	(±6%)	32,000
More than one pack	6%	(±1%)	78,000	2%	(±3%)	2,000	3%	(±6%)	1,000
HANDGUN OWNERSHIP									
Adults	20%	(±1%)	1,332,000	13%	(±3%)	78,000	15%	(±6%)	25,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	51%	(±1%)	3,207,000	39%	(±3%)	213,000	38%	(±8%)	62,000
Mild or moderate	34%	(±1%)	2,154,000	36%	(±2%)	197,000	36%	(±8%)	59,000
Severely	14%	(±1%)	894,000	25%	(±2%)	139,000	25%	(±7%)	41,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	52%	(±1%)	3,448,000	37%	(±4%)	228,000	45%	(±8%)	77,000
Good	27%	(±1%)	1,807,000	31%	(±4%)	189,000	28%	(±7%)	49,000
Poor or fair	21%	(±1%)	1,359,000	32%	(±4%)	192,000	27%	(±7%)	46,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±1%)	1,151,000	21%	(±3%)	127,000	16%	(±6%)	28,000
Diabetes	6%	(±1%)	371,000	9%	(±2%)	51,000	6%	(±4%)	10,000
BP or hypertension	16%	(±1%)	1,041,000	22%	(±3%)	133,000	18%	(±6%)	30,000
MEDICAL CONDITION, CHILDREN									
Asthma	7%	(±1%)	206,000	8%	(±4%)	32,000	8%	(±9%)	9,000

Appendix C-6: South Continued

Summary of Results by Service Planning Area, 1997¹

	SOUTH HD			SOUTHEAST HD			SOUTHWEST HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	61%	(±8%)	59,000	66%	(±12%)	59,000	63%	(±6%)	152,000
Light	22%	(±7%)	21,000	17%	(±9%)	16,000	25%	(±5%)	60,000
Moderate	14%	(±6%)	13,000	10%	(±7%)	9,000	8%	(±3%)	20,000
Heavy	4%	(±3%)	4,000	7%	(±6%)	7,000	4%	(±2%)	10,000
SEAT BELT NOT USED REGULARLY									
Adults	14%	(±6%)	14,000	16%	(±9%)	14,000	11%	(±4%)	27,000
Children ⁵	13%	(±9%)	10,000	2%	(±5%)	2,000	1%	(±3%)	1,000
CIGARETTE SMOKING									
Adults	22%	(±7%)	22,000	14%	(±8%)	13,000	19%	(±5%)	47,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	98%	(±5%)	22,000	100%	(±0%)	13,000	98%	(±3%)	46,000
More than one pack	2%	(±5%)	<1,000	0%		0	2%	(±3%)	1,000
HANDGUN OWNERSHIP									
Adults	13%	(±6%)	12,000	7%	(±6%)	7,000	14%	(±4%)	34,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	31%	(±8%)	28,000	41%	(±13%)	30,000	41%	(±6%)	93,000
Mild or moderate	37%	(±8%)	33,000	41%	(±13%)	30,000	34%	(±6%)	75,000
Severely	32%	(±8%)	29,000	18%	(±10%)	13,000	25%	(±5%)	56,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	26%	(±7%)	26,000	26%	(±10%)	24,000	41%	(±6%)	101,000
Good	33%	(±8%)	33,000	37%	(±12%)	35,000	30%	(±5%)	72,000
Poor or fair	41%	(±8%)	40,000	37%	(±12%)	34,000	29%	(±5%)	71,000
MEDICAL CONDITION, ADULTS									
Arthritis	21%	(±5%)	21,000	20%	(±10%)	18,000	25%	(±7%)	61,000
Diabetes	11%	(±4%)	10,000	4%	(±4%)	3,000	11%	(±4%)	27,000
BP or hypertension	26%	(±7%)	26,000	17%	(±9%)	15,000	26%	(±5%)	62,000
MEDICAL CONDITION, CHILDREN									
Asthma	9%	(±8%)	7,000	5%	(±8%)	5,000	9%	(±7%)	11,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI < 25 then the person is classified as not overweight, 25 ≤ BMI < 30 as mild or moderately overweight, and BMI ≥ 30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-6: South Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SOUTH SPA			COMPTON HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
<i>Access to Health Care and Utilization of Health Care Services</i>									
INSURANCE STATUS, ADULTS									
Medi-Cal	11%	(±1%)	705,000	18%	(±3%)	109,000	16%	(±6%)	28,000
Medicare	9%	(±1%)	569,000	7%	(±2%)	40,000	5%	(±4%)	8,000
Private	50%	(±1%)	3,279,000	35%	(±4%)	214,000	43%	(±7%)	73,000
Uninsured	31%	(±1%)	2,035,000	40%	(±4%)	244,000	36%	(±8%)	60,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	31%	(±2%)	851,000	45%	(±8%)	180,000	43%	(±16%)	47,000
Private	44%	(±3%)	1,229,000	22%	(±6%)	89,000	32%	(±15%)	36,000
Uninsured	25%	(±2%)	696,000	33%	(±7%)	131,000	25%	(±14%)	28,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	56%	(±1%)	3,628,000	44%	(±4%)	258,000	52%	(±8%)	85,000
Other regular provider	12%	(±1%)	804,000	15%	(±3%)	87,000	16%	(±6%)	26,000
No regular provider	26%	(±1%)	1,656,000	29%	(±4%)	169,000	21%	(±7%)	34,000
County or community clinics	6%	(±1%)	415,000	13%	(±3%)	75,000	11%	(±5%)	18,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	60%	(±3%)	1,678,000	47%	(±8%)	187,000	60%	(±16%)	66,000
Other regular provider	10%	(±2%)	289,000	14%	(±5%)	57,000	19%	(±13%)	21,000
No regular provider	15%	(±2%)	423,000	18%	(±6%)	71,000	6%	(±8%)	70,000
County or community clinics	14%	(±2%)	394,000	21%	(±6%)	84,000	15%	(±11%)	16,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	35%	(±1%)	2,228,000	44%	(±4%)	258,000	37%	(±8%)	62,000
Somewhat or very easy	65%	(±1%)	4,238,000	56%	(±4%)	333,000	63%	(±8%)	105,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	16%	(±1%)	1,049,000	21%	(±2%)	126,000	19%	(±6%)	32,000
Mental health care	8%	(±1%)	522,000	9%	(±3%)	55,000	7%	(±4%)	12,000
Dental care	27%	(±1%)	1,753,000	33%	(±1%)	199,000	35%	(±8%)	60,000
Vision care	15%	(±1%)	1,012,000	18%	(±2%)	105,000	16%	(±6%)	27,000
Doctor's visit	18%	(±1%)	1,193,000	23%	(±2%)	138,000	21%	(±7%)	36,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	45%	(±4%)	270,000	42%	(±8%)	71,000
Prostate exam ⁷	51%	(±3%)	759,000	42%	(±12%)	33,000	31%	(±25%)	6,000
Pap test ⁸	73%	(±1%)	2,387,000	71%	(±4%)	257,000	78%	(±8%)	78,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	67%	(±4%)	242,000	72%	(±8%)	72,000
Testicular exam ⁹	28%	(±2%)	897,000	23%	(±6%)	53,000	13%	(±10%)	9,000
Mammogram ¹⁰	76%	(±2%)	796,000	79%	(±7%)	80,000	80%	(±15%)	20,000

Appendix C-6: South Continued

Summary of Results by Service Planning Area, 1997¹

	SOUTH HD			SOUTHEAST HD			SOUTHWEST HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	19%	(±3%)	19,000	22%	(±10%)	21,000	17%	(±4%)	41,000
Medicare	7%	(±4%)	7,000	7%	(±6%)	7,000	8%	(±3%)	18,000
Private	26%	(±6%)	26,000	23%	(±10%)	21,000	38%	(±6%)	94,000
Uninsured	48%	(±5%)	48,000	48%	(±12%)	44,000	37%	(±6%)	91,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	45%	(±14%)	35,000	56%	(±19%)	52,000	38%	(±12%)	45,000
Private	9%	(±8%)	7,000	9%	(±11%)	8,000	32%	(±11%)	38,000
Uninsured	46%	(±14%)	36,000	34%	(±18%)	32,000	30%	(±11%)	35,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	37%	(±8%)	35,000	24%	(±10%)	22,000	49%	(±6%)	116,000
Other regular provider	16%	(±6%)	15,000	11%	(±8%)	11,000	15%	(±4%)	35,000
No regular provider	32%	(±8%)	30,000	40%	(±12%)	36,000	29%	(±5%)	68,000
County or community clinics	16%	(±6%)	15,000	25%	(±10%)	23,000	8%	(±3%)	18,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	34%	(±13%)	26,000	33%	(±18%)	32,000	54%	(±12%)	63,000
Other regular provider	8%	(±8%)	6,000	13%	(±13%)	13,000	15%	(±9%)	17,000
No regular provider	25%	(±12%)	19,000	25%	(±16%)	24,000	17%	(±9%)	20,000
County or community clinics	33%	(±13%)	26,000	28%	(±17%)	27,000	13%	(±8%)	15,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	51%	(±8%)	49,000	50%	(±12%)	46,000	43%	(±6%)	102,000
Somewhat or very easy	49%	(±8%)	47,000	50%	(±12%)	46,000	57%	(±6%)	136,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	26%	(±7%)	26,000	31%	(±11%)	29,000	16%	(±4%)	38,000
Mental health care	12%	(±5%)	12,000	11%	(±7%)	10,000	9%	(±3%)	21,000
Dental care	34%	(±8%)	34,000	39%	(±12%)	36,000	28%	(±5%)	68,000
Vision care	17%	(±6%)	17,000	19%	(±9%)	17,000	18%	(±4%)	44,000
Doctor's visit	30%	(±8%)	29,000	31%	(±11%)	29,000	18%	(±4%)	44,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	51%	(±8%)	49,000	45%	(±12%)	42,000	46%	(±6%)	108,000
Prostate exam ⁷	38%	(±23%)	5,000	18%	(±34%)	2,000	54%	(±16%)	20,000
Pap test ⁸	69%	(±9%)	40,000	64%	(±13%)	37,000	70%	(±6%)	103,000
Clinical breast exam ⁸	62%	(±9%)	35,000	61%	(±13%)	36,000	68%	(±6%)	99,000
Testicular exam ⁹	23%	(±12%)	9,000	29%	(±22%)	10,000	29%	(±10%)	25,000
Mammogram ¹⁰	78%	(±18%)	9,000	85%	(±27%)	6,000	78%	(±9%)	45,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-7: East

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			EAST SPA			BELLFLOWER HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	43%	(±1%)	2,770,000	42%	(±3%)	364,000	39%	(±5%)	95,000
Light	38%	(±1%)	2,440,000	39%	(±3%)	337,000	44%	(±5%)	105,000
Moderate	14%	(±1%)	927,000	13%	(±2%)	110,000	13%	(±3%)	31,000
Heavy	5%	(±1%)	331,000	6%	(±1%)	47,000	4%	(±2%)	10,000
SEAT BELT NOT USED REGULARLY									
Adults	11%	(±1%)	737,000	11%	(±2%)	100,000	11%	(±3%)	26,000
Children ⁵	5%	(±1%)	129,000	4%	(±2%)	17,000	5%	(±4%)	6,000
CIGARETTE SMOKING									
Adults	18%	(±1%)	1,210,000	19%	(±3%)	167,000	23%	(±4%)	56,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	94%	(±1%)	1,131,000	95%	(±3%)	159,000	93%	(±5%)	52,000
More than one pack	6%	(±1%)	78,000	5%	(±3%)	8,000	7%	(±5%)	4,000
HANDGUN OWNERSHIP									
Adults	20%	(±1%)	1,332,000	23%	(±3%)	198,000	29%	(±5%)	71,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	51%	(±1%)	3,207,000	43%	(±3%)	356,000	48%	(±5%)	110,000
Mild or moderate	34%	(±1%)	2,154,000	39%	(±3%)	322,000	32%	(±5%)	73,000
Severely	14%	(±1%)	894,000	18%	(±3%)	147,000	20%	(±4%)	47,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	52%	(±1%)	3,448,000	47%	(±3%)	415,000	53%	(±5%)	130,000
Good	27%	(±1%)	1,807,000	29%	(±3%)	253,000	30%	(±5%)	73,000
Poor or fair	21%	(±1%)	1,357,000	24%	(±3%)	213,000	18%	(±4%)	43,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±1%)	1,151,000	18%	(±3%)	161,000	21%	(±4%)	51,000
Diabetes	6%	(±1%)	371,000	5%	(±1%)	42,000	5%	(±2%)	13,000
BP or hypertension	16%	(±1%)	1,041,000	16%	(±2%)	139,000	16%	(±4%)	39,000
MEDICAL CONDITION, CHILDREN									
Asthma	7%	(±1%)	206,000	8%	(±3%)	34,000	9%	(±6%)	11,000

Appendix C-7: East Continued

Summary of Results by Service Planning Area, 1997¹

	EAST LOS ANGELES HD			SAN ANTONIO HD			WHITTIER HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	44%	(±9%)	62,000	47%	(±6%)	123,000	39%	(±7%)	83,000
Light	33%	(±8%)	47,000	37%	(±6%)	99,000	41%	(±7%)	87,000
Moderate	16%	(±6%)	22,000	12%	(±4%)	31,000	13%	(±5%)	27,000
Heavy	7%	(±5%)	10,000	4%	(±2%)	11,000	8%	(±4%)	16,000
SEAT BELT NOT USED REGULARLY									
Adults	14%	(±6%)	20,000	12%	(±4%)	33,000	10%	(±4%)	21,000
Children ⁵	5%	(±8%)	3,000	4%	(±4%)	5,000	2%	(±4%)	3,000
CIGARETTE SMOKING									
Adults	17%	(±7%)	25,000	18%	(±5%)	50,000	17%	(±5%)	37,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	100%	(±0%)	25,000	93%	(±8%)	46,000	97%	(±6%)	36,000
More than one pack	0%		0	7%	(±8%)	4,000	3%	(±6%)	1,000
HANDGUN OWNERSHIP									
Adults	17%	(±7%)	25,000	18%	(±5%)	48,000	25%	(±6%)	54,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	46%	(±9%)	62,000	42%	(±6%)	104,000	38%	(±7%)	80,000
Mild or moderate	37%	(±9%)	50,000	43%	(±6%)	109,000	43%	(±7%)	90,000
Severely	17%	(±7%)	23,000	15%	(±5%)	38,000	19%	(±6%)	40,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	34%	(±8%)	50,000	43%	(±6%)	118,000	54%	(±7%)	118,000
Good	33%	(±8%)	49,000	25%	(±5%)	68,000	29%	(±6%)	64,000
Poor or fair	33%	(±8%)	48,000	32%	(±6%)	86,000	17%	(±5%)	36,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±7%)	27,000	14%	(±4%)	38,000	21%	(±6%)	46,000
Diabetes	3%	(±3%)	4,000	5%	(±3%)	13,000	5%	(±3%)	11,000
BP or hypertension	12%	(±6%)	17,000	16%	(±4%)	43,000	19%	(±5%)	41,000
MEDICAL CONDITION, CHILDREN									
Asthma	9%	(±10%)	6,000	9%	(±6%)	14,000	3%	(±5%)	3,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.

Appendix C-7: East Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			EAST SPA			BELLFLOWER HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	11%	(±1%)	705,000	13%	(±2%)	112,000	12%	(±3%)	29,000
Medicare	9%	(±1%)	569,000	8%	(±2%)	75,000	9%	(±3%)	21,000
Private	50%	(±1%)	3,279,000	50%	(±3%)	439,000	56%	(±5%)	136,000
Uninsured	31%	(±1%)	2,035,000	29%	(±3%)	254,000	24%	(±4%)	59,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	31%	(±2%)	851,000	31%	(±6%)	137,000	28%	(±9%)	34,000
Private	44%	(±3%)	1,229,000	49%	(±6%)	220,000	59%	(±9%)	72,000
Uninsured	25%	(±2%)	696,000	20%	(±5%)	91,000	14%	(±7%)	17,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	56%	(±1%)	3,628,000	55%	(±3%)	476,000	62%	(±5%)	150,000
Other regular provider	12%	(±1%)	804,000	11%	(±2%)	100,000	14%	(±4%)	35,000
No regular provider	26%	(±1%)	1,656,000	26%	(±3%)	228,000	21%	(±4%)	50,000
County or community clinics	6%	(±1%)	415,000	8%	(±2%)	66,000	3%	(±2%)	7,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	60%	(±3%)	1,656,000	63%	(±5%)	281,000	75%	(±8%)	91,000
Other regular provider	10%	(±2%)	289,000	9%	(±4%)	41,000	7%	(±5%)	8,000
No regular provider	15%	(±2%)	423,000	15%	(±4%)	65,000	15%	(±7%)	18,000
County or community clinics	14%	(±2%)	394,000	13%	(±4%)	58,000	3%	(±3%)	4,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	35%	(±1%)	2,228,000	36%	(±3%)	313,000	33%	(±5%)	80,000
Somewhat or very easy	65%	(±1%)	4,238,000	64%	(±3%)	556,000	67%	(±5%)	165,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	16%	(±1%)	1,049,000	17%	(±2%)	152,000	18%	(±4%)	43,000
Mental health care	8%	(±1%)	522,000	8%	(±3%)	67,000	8%	(±3%)	19,000
Dental care	27%	(±1%)	1,753,000	28%	(±2%)	250,000	30%	(±5%)	73,000
Vision care	15%	(±1%)	1,012,000	16%	(±2%)	136,000	16%	(±4%)	38,000
Doctor's visit	18%	(±1%)	1,193,000	18%	(±2%)	162,000	19%	(±4%)	47,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	33%	(±3%)	287,000	31%	(±5%)	74,000
Prostate exam ⁷	51%	(±3%)	759,000	50%	(±7%)	94,000	51%	(±11%)	30,000
Pap test ⁸	73%	(±1%)	2,387,000	70%	(±4%)	307,000	72%	(±6%)	93,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	69%	(±4%)	303,000	75%	(±5%)	96,000
Testicular exam ⁹	28%	(±2%)	897,000	26%	(±5%)	111,000	27%	(±8%)	30,000
Mammogram ¹⁰	76%	(±2%)	796,000	71%	(±7%)	89,000	78%	(±10%)	28,000

Appendix C-7: East Continued

Summary of Results by Service Planning Area, 1997¹

	EAST LOS ANGELES HD			SAN ANTONIO HD			WHITTIER HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	18%	(±7%)	26,000	16%	(±4%)	42,000	7%	(±3%)	15,000
Medicare	5%	(±4%)	8,000	9%	(±4%)	26,000	9%	(±4%)	21,000
Private	42%	(±9%)	62,000	39%	(±6%)	107,000	62%	(±7%)	135,000
Uninsured	35%	(±8%)	51,000	36%	(±6%)	96,000	22%	(±6%)	48,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	44%	(±17%)	28,000	41%	(±11%)	60,000	13%	(±9%)	16,000
Private	29%	(±15%)	19,000	30%	(±10%)	43,000	74%	(±12%)	86,000
Uninsured	27%	(±15%)	17,000	29%	(±10%)	42,000	13%	(±9%)	15,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	47%	(±9%)	68,000	44%	(±6%)	118,000	65%	(±7%)	140,000
Other regular provider	10%	(±5%)	15,000	10%	(±4%)	26,000	11%	(±4%)	24,000
No regular provider	29%	(±8%)	42,000	35%	(±6%)	95,000	19%	(±5%)	41,000
County or community clinics	14%	(±6%)	20,000	11%	(±4%)	30,000	4%	(±3%)	9,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	45%	(±17%)	29,000	49%	(±11%)	71,000	78%	(±12%)	90,000
Other regular provider	16%	(±12%)	10,000	10%	(±6%)	14,000	7%	(±7%)	9,000
No regular provider	13%	(±12%)	9,000	18%	(±8%)	26,000	11%	(±9%)	12,000
County or community clinics	26%	(±15%)	17,000	23%	(±9%)	34,000	4%	(±5%)	4,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	47%	(±9%)	66,000	39%	(±6%)	104,000	29%	(±6%)	63,000
Somewhat or very easy	53%	(±9%)	75,000	61%	(±6%)	164,000	71%	(±6%)	153,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	23%	(±7%)	34,000	18%	(±4%)	49,000	12%	(±4%)	26,000
Mental health care	8%	(±5%)	12,000	7%	(±3%)	18,000	8%	(±4%)	18,000
Dental care	26%	(±8%)	38,000	31%	(±6%)	85,000	24%	(±6%)	53,000
Vision care	16%	(±6%)	23,000	14%	(±4%)	37,000	17%	(±5%)	37,000
Doctor's visit	22%	(±7%)	31,000	19%	(±5%)	52,000	15%	(±5%)	32,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	32%	(±8%)	46,000	38%	(±6%)	103,000	30%	(±6%)	64,000
Prostate exam ⁷	46%	(±20%)	13,000	63%	(±15%)	28,000	61%	(±14%)	34,000
Pap test ⁸	67%	(±10%)	45,000	70%	(±7%)	95,000	70%	(±8%)	75,000
Clinical breast exam ⁸	60%	(±11%)	40,000	65%	(±7%)	88,000	74%	(±8%)	78,000
Testicular exam ⁹	16%	(±10%)	12,000	26%	(±8%)	34,000	33%	(±10%)	35,000
Mammogram ¹⁰	64%	(±21%)	11,000	72%	(±14%)	21,000	67%	(±13%)	28,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.
2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.
3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).
4. Adults aged 18 years and older.
5. Children aged 0 to 17 years.
6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI < 25 then the person is classified as not overweight, 25 = < BMI < 30 as mild or moderately overweight, and BMI > = 30 as severely overweight persons.
7. Men aged 40 years and older.
8. Women aged 18 years and older.
9. Men aged 18 years and older.
10. Women aged 50 years and older.

Appendix C-8: South Bay

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SOUTH BAY SPA			HARBOR HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	43%	(±1%)	2,770,000	38%	(±3%)	385,000	43%	(±7%)	58,000
Light	38%	(±1%)	2,440,000	41%	(±3%)	413,000	36%	(±7%)	48,000
Moderate	14%	(±1%)	927,000	15%	(±2%)	153,000	13%	(±5%)	18,000
Heavy	5%	(±1%)	331,000	6%	(±1%)	64,000	8%	(±4%)	11,000
SEAT BELT NOT USED REGULARLY									
Adults	11%	(±1%)	737,000	11%	(±2%)	118,000	12%	(±5%)	16,000
Children ⁵	5%	(±1%)	129,000	7%	(±3%)	27,000	17%	(±12%)	12,000
CIGARETTE SMOKING									
Adults	18%	(±1%)	1,210,000	17%	(±2%)	183,000	18%	(±6%)	24,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	94%	(±1%)	1,131,000	92%	(±4%)	168,000	97%	(±6%)	24,000
More than one pack	6%	(±1%)	78,000	8%	(±4%)	14,000	3%	(±6%)	1,000
HANDGUN OWNERSHIP									
Adults	20%	(±1%)	1,332,000	22%	(±2%)	226,000	22%	(±6%)	30,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	51%	(±1%)	3,207,000	52%	(±3%)	514,000	45%	(±8%)	57,000
Mild or moderate	34%	(±1%)	2,154,000	35%	(±3%)	347,000	39%	(±8%)	50,000
Severely	14%	(±1%)	894,000	13%	(±2%)	132,000	16%	(±6%)	20,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	52%	(±1%)	3,448,000	56%	(±3%)	584,000	55%	(±8%)	77,000
Good	27%	(±1%)	1,807,000	26%	(±2%)	274,000	27%	(±7%)	37,000
Poor or Fair	21%	(±1%)	1,359,000	18%	(±2%)	187,000	18%	(±6%)	25,000
MEDICAL CONDITION, ADULTS									
Arthritis	18%	(±1%)	1,151,000	17%	(±2%)	178,000	17%	(±6%)	23,000
Diabetes	6%	(±1%)	371,000	5%	(±1%)	52,000	4%	(±3%)	6,000
BP or hypertension	16%	(±1%)	1,041,000	16%	(±2%)	161,000	12%	(±5%)	17,000
MEDICAL CONDITION, CHILDREN									
Asthma	7%	(±1%)	206,000	11%	(±4%)	42,000	11%	(±10%)	8,000

Appendix C-8: South Bay Continued

Summary of Results by Service Planning Area, 1997¹

	INGLEWOOD HD			LONG BEACH HD			TORRANCE HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Health Risk Behaviors									
ALCOHOL USE ³ , ADULTS ⁴									
Non-drinker	46%	(±6%)	122,000	35%	(±5%)	104,000	32%	(±5%)	101,000
Light	38%	(±6%)	100,000	41%	(±5%)	121,000	45%	(±5%)	143,000
Moderate	10%	(±3%)	25,000	18%	(±4%)	53,000	18%	(±4%)	56,000
Heavy	6%	(±3%)	16,000	6%	(±3%)	18,000	6%	(±2%)	19,000
SEAT BELT NOT USED REGULARLY									
Adults	13%	(±4%)	34,000	11%	(±3%)	34,000	10%	(±3%)	34,000
Children ⁵	5%	(±5%)	6,000	5%	(±5%)	5,000	5%	(±5%)	4,000
CIGARETTE SMOKING									
Adults	17%	(±4%)	47,000	18%	(±4%)	55,000	17%	(±4%)	56,000
NUMBER OF PACKS SMOKED, ADULTS									
One pack or fewer	90%	(±8%)	42,000	92%	(±7%)	51,000	92%	(±6%)	52,000
More than one pack	10%	(±8%)	5,000	8%	(±7%)	5,000	8%	(±6%)	4,000
HANDGUN OWNERSHIP									
Adults	22%	(±5%)	58,000	19%	(±4%)	57,000	24%	(±4%)	81,000
Health Status									
WEIGHT STATUS ⁶									
Not overweight	48%	(±5%)	122,000	55%	(±6%)	159,000	55%	(±5%)	176,000
Mild or moderate	37%	(±5%)	95,000	32%	(±6%)	94,000	34%	(±5%)	108,000
Severely	14%	(±4%)	36,000	13%	(±4%)	38,000	12%	(±3%)	38,000
SELF-PERCEIVED HEALTH STATUS, ADULTS									
Very good or excellent	51%	(±6%)	138,000	53%	(±5%)	161,000	63%	(±5%)	208,000
Good	28%	(±5%)	77,000	28%	(±5%)	84,000	23%	(±4%)	76,000
Poor or fair	21%	(±5%)	57,000	19%	(±4%)	59,000	14%	(±3%)	46,000
MEDICAL CONDITION, ADULTS									
Arthritis	16%	(±4%)	43,000	18%	(±4%)	53,000	18%	(±4%)	59,000
Diabetes	6%	(±3%)	15,000	5%	(±2%)	15,000	5%	(±2%)	15,000
BP or hypertension	18%	(±4%)	49,000	15%	(±4%)	46,000	15%	(±3%)	49,000
MEDICAL CONDITION, CHILDREN									
Asthma	14%	(±8%)	17,000	8%	(±6%)	8,000	11%	(±7%)	9,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.
2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.
3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).
4. Adults aged 18 years and older.
5. Children aged 0 to 17 years.
6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI<25 then the person is classified as not overweight, 25=<BMI<30 as mild or moderately overweight, and BMI>=30 as severely overweight persons.
7. Men aged 40 years and older.
8. Women aged 18 years and older.
9. Men aged 18 years and older.
10. Women aged 50 years and older.

Appendix C-8: South Bay Continued

Summary of Results by Service Planning Area, 1997¹

	LOS ANGELES COUNTY			SOUTH BAY SPA			HARBOR HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	11%	(±1%)	705,000	11%	(±2%)	111,000	13%	(±5%)	18,000
Medicare	9%	(±1%)	569,000	8%	(±2%)	86,000	12%	(±5%)	17,000
Private	50%	(±1%)	3,279,000	54%	(±3%)	556,000	51%	(±8%)	69,000
Uninsured	31%	(±1%)	2,035,000	28%	(±3%)	286,000	24%	(±7%)	32,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	31%	(±2%)	851,000	27%	(±6%)	102,000	30%	(±15%)	21,000
Private	44%	(±3%)	1,229,000	51%	(±6%)	192,000	40%	(±15%)	28,000
Uninsured	25%	(±2%)	696,000	22%	(±5%)	84,000	30%	(±14%)	21,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	56%	(±1%)	3,628,000	61%	(±3%)	627,000	61%	(±7%)	84,000
Other regular provider	12%	(±1%)	804,000	12%	(±2%)	129,000	12%	(±5%)	17,000
No regular provider	26%	(±1%)	1,656,000	23%	(±2%)	238,000	22%	(±6%)	31,000
County or Community clinics	6%	(±1%)	415,000	4%	(±1%)	40,000	5%	(±3%)	6,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	60%	(±3%)	1,678,000	61%	(±6%)	233,000	39%	(±15%)	28,000
Other regular provider	10%	(±2%)	289,000	11%	(±4%)	44,000	14%	(±11%)	10,000
No regular provider	15%	(±2%)	423,000	17%	(±5%)	66,000	31%	(±15%)	22,000
County or Community clinics	14%	(±2%)	394,000	10%	(±4%)	40,000	15%	(±11%)	11,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	35%	(±1%)	2,228,000	28%	(±3%)	288,000	24%	(±7%)	32,000
Somewhat or very easy	65%	(±1%)	4,238,000	72%	(±3%)	731,000	76%	(±7%)	105,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	16%	(±1%)	1,049,000	13%	(±2%)	136,000	8%	(±4%)	11,000
Mental health care	8%	(±1%)	522,000	6%	(±3%)	62,000	6%	(±4%)	8,000
Dental care	27%	(±1%)	1,753,000	21%	(±2%)	224,000	15%	(±5%)	21,000
Vision care	15%	(±1%)	1,012,000	15%	(±2%)	155,000	12%	(±5%)	17,000
Doctor's visit	18%	(±1%)	1,193,000	16%	(±2%)	165,000	11%	(±5%)	16,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	36%	(±1%)	2,353,000	37%	(±3%)	377,000	33%	(±7%)	45,000
Prostate exam ⁷	51%	(±3%)	759,000	55%	(±6%)	123,000	33%	(±15%)	11,000
Pap test ⁸	73%	(±1%)	2,387,000	77%	(±3%)	407,000	75%	(±8%)	52,000
Clinical breast exam ⁸	70%	(±1%)	2,310,000	73%	(±3%)	389,000	72%	(±8%)	50,000
Testicular exam ⁹	28%	(±2%)	897,000	30%	(±4%)	147,000	30%	(±12%)	19,000
Mammogram ¹⁰	76%	(±2%)	796,000	77%	(±5%)	128,000	76%	(±15%)	16,000

Appendix C-8: South Bay Continued

Summary of Results by Service Planning Area, 1997¹

	INGLEWOOD HD			LONG BEACH HD			TORRANCE HD		
	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number	Percent	(±95%) Confidence Interval ²	Estimated Number
Access to Health Care and Utilization of Health Care Services									
INSURANCE STATUS, ADULTS									
Medi-Cal	13%	(±4%)	36,000	10%	(±3%)	31,000	8%	(±3%)	26,000
Medicare	7%	(±3%)	19,000	6%	(±3%)	20,000	9%	(±3%)	30,000
Private	50%	(±6%)	136,000	51%	(±5%)	155,000	60%	(±5%)	195,000
Uninsured	30%	(±5%)	81,000	32%	(±5%)	97,000	23%	(±4%)	76,000
INSURANCE STATUS, CHILDREN									
Medi-Cal	25%	(±11%)	31,000	37%	(±11%)	38,000	15%	(±8%)	12,000
Private	53%	(±12%)	65,000	43%	(±11%)	44,000	66%	(±11%)	55,000
Uninsured	22%	(±10%)	27,000	20%	(±9%)	20,000	19%	(±9%)	16,000
REGULAR SOURCE OF HEALTH CARE, ADULTS									
Private MD/HMO	56%	(±6%)	151,000	55%	(±5%)	165,000	69%	(±4%)	228,000
Other regular provider	13%	(±4%)	35,000	16%	(±4%)	47,000	9%	(±3%)	29,000
No regular provider	26%	(±5%)	71,000	24%	(±5%)	72,000	19%	(±4%)	63,000
County or community clinics	5%	(±2%)	13,000	4%	(±2%)	13,000	3%	(±2%)	9,000
REGULAR SOURCE OF HEALTH CARE, CHILDREN									
Private MD/HMO	61%	(±12%)	75,000	61%	(±11%)	62,000	79%	(±9%)	68,000
Other regular provider	9%	(±7%)	11,000	14%	(±8%)	14,000	10%	(±7%)	8,000
No regular provider	22%	(±10%)	27,000	11%	(±7%)	12,000	6%	(±5%)	5,000
County or community clinics	9%	(±7%)	11,000	13%	(±7%)	14,000	5%	(±5%)	5,000
DIFFICULTY GETTING MEDICAL CARE, ADULTS									
Somewhat or very difficult	35%	(±6%)	94,000	32%	(±5%)	94,000	21%	(±4%)	67,000
Somewhat or very easy	65%	(±6%)	172,000	68%	(±5%)	199,000	79%	(±4%)	254,000
EXPERIENCED DIFFICULTY OBTAINING MEDICAL SERVICES, ADULTS									
Prescription medicines	18%	(±4%)	49,000	14%	(±4%)	43,000	10%	(±3%)	33,000
Mental health care	7%	(±3%)	20,000	5%	(±2%)	16,000	6%	(±2%)	18,000
Dental care	26%	(±5%)	70,000	23%	(±4%)	70,000	19%	(±4%)	63,000
Vision care	19%	(±5%)	53,000	16%	(±4%)	49,000	11%	(±3%)	37,000
Doctor's visit	20%	(±5%)	55,000	19%	(±4%)	59,000	11%	(±3%)	35,000
PREVENTIVE SERVICES, ADULTS									
HIV/AIDS test ⁴	45%	(±6%)	119,000	42%	(±5%)	125,000	28%	(±4%)	89,000
Prostate exam ⁷	47%	(±13%)	26,000	57%	(±13%)	34,000	54%	(±11%)	40,000
Pap test ⁸	74%	(±6%)	98,000	78%	(±5%)	125,000	78%	(±5%)	132,000
Clinical breast exam ⁸	67%	(±7%)	88,000	74%	(±6%)	119,000	77%	(±5%)	131,000
Testicular exam ⁹	30%	(±8%)	39,000	33%	(±8%)	46,000	28%	(±7%)	43,000
Mammogram ¹⁰	82%	(±11%)	31,000	70%	(±12%)	28,000	79%	(±8%)	52,000

1. The percentages and numbers reported reflect the best estimates in each area. These estimates should not be considered exact population numbers but, should be used as trend data for planning purposes.

2. The actual percentages and numbers may be lower or higher based on the confidence interval reported for each area. We are 95% confident that the actual percentage in each area is within the range of the confidence interval presented for that area.

3. Alcohol users were classified into four groups based on the number of the number of drinks consumed. A "drink" is defined as a can or bottle of beer, one glass of wine or cocktail or shot of liquor. Each drink is assumed to contain 0.5 oz of ethyl alcohol. Accordingly, a non-drinker is one who consumes 0.0 oz of ethyl alcohol daily, light drinker (0.001 to 0.210 oz), moderate drinker (0.211 to 0.990 oz), and heavy drinker (1.0 oz or more).

4. Adults aged 18 years and older.

5. Children aged 0 to 17 years.

6. Weight status is determined based on the body mass index (BMI). BMI is calculated by taking into account weight and height of the respondent. Accordingly, when was BMI < 25 then the person is classified as not overweight, 25 = < BMI < 30 as mild or moderately overweight, and BMI > = 30 as severely overweight persons.

7. Men aged 40 years and older.

8. Women aged 18 years and older.

9. Men aged 18 years and older.

10. Women aged 50 years and older.