

# ZIKA SCREENING TOOL FOR PREGNANT WOMEN



*(To be administered by nurse, check-in receptionist, or other healthcare provider)*

All pregnant women should be assessed for possible Zika virus exposure<sup>1</sup> at each prenatal care visit. Use this tool to evaluate pregnant women for exposure to Zika virus and for signs and symptoms of Zika virus disease to determine whether testing is indicated.

**NOTE:** If your pregnant patient has questions about Zika testing, educational factsheets are available on CDC's website: <http://www.cdc.gov/zika/hc-providers/pregnant-woman.html>

## Assess for Possible Exposure<sup>1</sup> to Zika Virus Infection

Circle response:

Do you live in or do you frequently travel (daily or weekly) to an area with active Zika virus transmission<sup>2</sup>?

YES | NO

Have you traveled to an area with Zika<sup>2</sup> during pregnancy or just before you became pregnant [8 weeks before conception or 6 weeks before your last menstrual period]?

YES | NO

Have you had sex (vaginal, anal, or oral sex) without a condom or shared sex toys with a partner(s) who lives in or has traveled to an area with Zika<sup>2</sup>?

YES | NO

➔ If your pregnant patient answered **"NO"** to ALL questions, she is at low risk for exposure to Zika.

## If Pregnant Patient Answered "Yes" to Any Question, Assess for Signs and Symptoms of Zika Virus Disease

Circle response:

Do you currently have or have you had (in the last 12 weeks) fever, rash, joint pain, or conjunctivitis (red eyes)?

YES | NO

- ➔ If your pregnant patient answered **"YES"** to having any of these signs or symptoms, she might have symptomatic Zika virus infection. Test in accordance with CDC guidance for symptomatic persons<sup>3</sup>.
- ➔ If your pregnant patient answered **"NO"** to having any signs or symptoms, she has been exposed and might have an asymptomatic Zika virus infection. Test in accordance with CDC guidance for asymptomatic pregnant women<sup>3</sup>.

**References:**

1. Possible exposure to Zika virus that warrants testing includes one or more of the following:
  - a. Living in an area with active transmission
  - b. Travel to an area with active transmission
  - c. Sex (vaginal, anal, and oral sex) without a condom or the sharing of sex toys with a person who traveled to or lives in an area with Zika.
2. Visit CDC's website to see areas with active Zika transmission: <http://www.cdc.gov/zika/geo/index.html>
3. Please see the algorithm on the back from CDC's Updated Interim Guidance for Health Care Providers Caring for Pregnant Women with Possible Zika Virus Exposure to guide testing and interpretation of results. ([http://www.cdc.gov/mmwr/volumes/65/wr/mm6529e1.htm?s\\_cid=mm6529e1\\_e](http://www.cdc.gov/mmwr/volumes/65/wr/mm6529e1.htm?s_cid=mm6529e1_e))



U.S. Department of Health and Human Services  
Centers for Disease Control and Prevention