2010-2012 LA HOPE Surveys

Domain/Subdomain			Tobacco, Alcohol & Substance Use / Smoking						Tobacco, Alcohol & Substance Use / Alcohol Consumption			Tobacco, Alcohol & Substance Use / Secondhand Smoke Exposure		
Indicator Description			Percentage of women having an infant/fetal death who smoked during pregnancy.			smoked cigarettes during						Percentage of women having an infant/fetal death who reported that during their pregnancy they have been exposed to secondhand smoke.		
			%	% 95% CI		% 95% CI			% 95% CI			% 95% CI		
	LA County		4.4	2.9	5.9	10.6	8.5	12.6	9.8	7.7	11.9	8.3	6.4	10.3
Race/Ethnicity	White		5.8*	1.6	10.0	17.8	11.2	24.5	21.1	14.2	27.9	8.3*	3.3	13.2
	Latina		1.8*	0.6	3.0	7.4	5.1	9.7	7.8	5.4	10.2	5.8	3.6	7.9
	African-American		13.1	7.1	19.1	21.2	14.0	28.4	11.3*	4.8	17.9	20.5	13.4	27.5
	Asian/Pacific Islander		5.0*	0.0	11.9	2.1*	0.0	6.1	3.5*	0.0	9.3	4.1*	0.0	9.6
Service Planning Areas	SPA 1	Antelope Valley	7.2*	0.9	13.5	17.7*	8.7	26.7	13.5*	5.5	21.5	14.6*	5.9	23.2
	SPA 2	San Fernando	3.9*	0.0	7.9	9.1*	4.5	13.6	9.6*	4.4	14.8	5.8*	2.0	9.5
	SPA 3	San Gabriel	1.6*	0.0	4.6	6.7*	1.7	11.6	4.3*	0.7	8.0	8.2*	2.6	13.8
	SPA 4	Metro	1.9*	0.0	4.6	6.0*	1.5	10.5	10.8*	5.0	16.7	2.7*	0.0	5.7
	SPA 5	West	-	-	-	7.8*	0.2	15.4	27.8*	13.0	42.6	5.7*	0.0	13.9
	SPA 6	South	8.3*	4.2	12.4	15.3	9.9	20.7	6.3*	2.5	10.0	14.2	8.9	19.6
	SPA 7	East	2.5*	0.0	5.3	5.8*	1.6	10.0	10.4*	4.2	16.5	7.4*	2.4	12.4
	SPA 8	South Bay	6.3*	1.5	11.2	15.6	8.7	22.5	12.7*	6.0	19.3	7.3*	2.0	12.5

^{(-) =} null value

For more information about LA HOPE project please click here

^{*}The estimate is statistically unstable (relative standard error > 25%) and therefore may not be appropriate to use for planning or policy purposes.