



COUNTY OF LOS ANGELES
Public Health

Sexually Transmitted Disease Program

Annual Adolescent STD Morbidity Report 2005



2615 South Grand Avenue, Room 500
Los Angeles, CA 90007
Tel: (213) 744-3070 Fax: (213) 749-9606

www.lapublichealth.org/std



COUNTY OF LOS ANGELES
Public Health

JONATHAN E. FIELDING, M.D., M.P.H.
Acting Director and Health Officer

JOHN SCHUNHOFF, Ph.D.
Acting Chief Deputy

Sexually Transmitted Disease Program

PETER R. KERNDT, M.D. M.P.H., Director

2615 South Grand Avenue, Room 500
Los Angeles, California 90007
TEL (213) 744-3070 • FAX (213) 749-9606

www.lapublichealth.org



BOARD OF SUPERVISORS

Gloria Molina
First District

Yvonne Brathwaite Burke
Second District

Zev Yaroslavsky
Third District

Don Knabe
Fourth District

Michael D. Antonovich
Fifth District

September 1, 2006

Dear Colleague:

I am pleased to release the *Annual Adolescent STD Morbidity Report, 2005*. This report provides a special, detailed assessment of reportable STDs among youth from 2001 to 2005 for Los Angeles County. The report encompasses cases and rates of chlamydia, gonorrhea, and syphilis infection in youth (ages 10 to 24). Caution should be taken when interpreting data for less than 20 observations, as directed by the CDC, 2002 (see Technical Notes).

This report is intended to help policy makers, community organizations, schools, and health providers recognize the STD burden and risk for youth, and to take appropriate action. It is important that sexually active youth be able to access routine STD screening. New tools make screening and treatment for chlamydia and gonorrhea easy and effective. Both diseases can now be diagnosed through a simple but highly accurate urine test, which can be administered in non-clinical settings. Complete cures for both gonorrhea and chlamydia can now be achieved with single doses of medication. In addition, state law now enables health providers to offer chlamydia patients oral medication to give directly to their sex partners, thereby decreasing the likelihood of reinfection.

Please consult the contact section in the back of this report to obtain more information, or learn how you or your agency can partner with the Los Angeles County STD Program to ensure effective prevention, screening, and treatment of STDs among youth.

Sincerely,

Peter R. Kerndt, MD, MPH
Director, STD Program

The *Sexually Transmitted Disease Adolescent Morbidity Report* is published on a yearly basis by the Sexually Transmitted Disease Program of the Los Angeles County Department of Health Services. This report is also available in PDF format, online at <http://www.lapublichealth.org/std>.

If you would like to receive surveillance reports and other information from the STD Program, please fill out the form in Appendix C and mail or fax it in to the STD Program. You may also register for STDInfo online at <http://ladhs.org/listserv> to receive surveillance reports and other information from the STD Program via e-mail.

SUGGESTED CITATION

Sexually Transmitted Disease Program, Los Angeles County Department of Health Services. *Annual Sexually Transmitted Disease Adolescent Morbidity Report, 2005*: pp. 1-110

Department of Public Health

Jonathan E. Fielding, MD, MPH
Director, Public Health & Health Officer

John Schunhoff, PhD
Acting Chief Deputy

Robert Kim-Farley, MD, MPH
Director, Disease Control Programs & Deputy Health Officer

~

Sexually Transmitted Disease Program

Peter R. Kerndt, MD, MPH
Director, Sexually Transmitted Disease Program

Sarah Guerry, MD
Medical Director

Mary Hayes
Field Services

Harlan Rotblatt, BA, Co-Director
Melina Boudov, MA, Co-Director
Adolescent Surveillance Team

Marian Eldahaby, BA, Linda Kim, Stephanie Koai,
Ryan Murphy, Danté Tolbert

Lisa V. Smith, MS, DrPH
Epidemiology

Getahun Aynalem, Patricia Barnett, Guillermo Campos, Cynthia Davison, Marian Eldahaby, Clarice Gillis, Kellie Hawkins, Chandra Higgins, Marjan Javanbakht, Stephanie Koai, Troy Meisner,

Jacqueline Moore, Ryan Murphy, Yifei Shi, Danté Tolbert, Joyce Wu

Mary Hayes
Field Services

Anissa Brown, Tony Bustamante, Deborah Carr, Sandra Chacon, David DeLaRiva, Oscar Dimas, Jim Fountain, Armando Garcia, Kelly Ho, Tameeka Johnson, Natalie Kelley, Lori Laubacher, Vince Lewis, Gentamu Mckinney, Judith Melendez, Margarita Melgoza, Gabriel Montano, Kenneth Myers, Leanne Olea, Jane Pannor, Ruth Pioquinto, Frank Ramirez, Marco Reyes, Roberto Rocha, Bob Simcik, Tracey Smith, Dave Tesfaye, Florence Thomas, Eleanor Tupas, Elizabeth Valencia, Kim Young

Kai-Jen Cheng, MS

Information Systems & Data Management

Sonia Alvarado, Leon Ayzazian, Diane Battaglia, Yetimwork Campbell, Debra Campos, Jacklyn Centeno, Teo Cruz, Chona Felix, Chamagne Footman, Yolanda Galvan, Mildred Gonzales, Alice Gonzalez, Nawal Habashi, Yoshiko Hayakawa, Peter He, Beatrice Johnson, JaMeisha Kennedy, Linda Koai, Lan Lam, Julius Lim, Kenn Liu, Shakeh Novagian, Janet Paul, Tien Pham, Harry Rollins, Terrie Ross, Joselene Samson, Michael Saucedo, Bertha Solis, Lillian Tom, Debra Valverde, David Yu, Yan Xu

Karen Menacker, RN, PHN

Nursing Unit

Jeanette Brown, Keisha Davis, Shirlesa Edwards, Fayga Greiner, Lenita Johnson, Alicia Magbanna, Terry Miller, Audrey Miyahira, Monica Munoz, Regina Umukoro, Franklin Villacres,

Betsy Swanson-Hollinger, MPH

Health Education

Emma Fredua, Ana Hernandez, Phoebe Macon, Susan Perez, Philip Phan, Sally Villanueva

Melina Boudov, MA

Infertility Prevention Projects

Kim Coffee, Gloria Escoto, Kim Harrison, Marisol Mejia, Kirsten Wilson

Jorge A. Montoya, PhD

Communications & Behavioral Research

Chi-Wai Au, Michael Granado, Carol Magee, Aaron Plant

Harlan Rotblatt, BA
Prevention Policy & Youth Services

The STD burden on youth is alarmingly high, both in Los Angeles County and nationally.

- Los Angeles County STD cases reported among youth ages 10-24 in 2005 totaled 30,063, including 24,592 cases of chlamydia, 5,320 cases of gonorrhea, and 151 cases of early syphilis in this age group. These cases represent approximately 63% of all chlamydia cases and 51% of all gonorrhea cases reported in the county in 2005.
- The number of reported STD cases among youth ages 10 to 24 in 2005 was greater than the total number of all reportable communicable diseases in Los Angeles County in 2004 (most recent data available) for all other age groups combined.
- Older youth, ages 15-19 and 20-24, have the most cases and the highest rates of reportable STDs of any five-year age group in the County.
- Chlamydia rates from 2001 to 2005 increased about 16% among 15-19 year-olds (to 1,543 cases per 100,000 population), and 20% among 20-24 year-olds (to 2,020 cases per 100,000). National chlamydia rates in 2004 (most recent data available) were higher for 15-19 year-olds and lower for 20-24 year-olds than rates in L.A. County.
- Gonorrhea rates increased about 38% among 15-19 year-olds (to 325 cases per 100,000), and 39% among 20-24 year-olds (to 427 cases per 100,000). National gonorrhea rates in 2004 (most recent data available) for youth were much higher than in Los Angeles County.
- Early syphilis rates increased about 176% among 10-24 year-olds to (6.9 cases per 100,000).

There are serious disparities in the STD burden on Los Angeles County youth.

- For chlamydia in 2005, African American female youth ages 15 to 19 had a rate over three and a half times higher than Hispanic females, who in turn had a rate more than two and a half times higher than white females in the same age group. African American males ages 15-19 had a chlamydia rate more than five times higher than Hispanic males, who in turn had a rate almost four times higher than that of White males in the same age group.
- For gonorrhea in 2005, African American female youth ages 15 to 19 had a rate 13.5 times higher than Hispanic females, who in turn had a rate nearly 1.5 times that of white females in the same age group. African American males ages 15-19 had a gonorrhea rate over 17 times higher than Hispanic males, who in turn had a rate about one and a half that of White males in the same age group.
- The 2005 rate for chlamydia among 15-19 year-olds in Service Planning Area (SPA) 6 (South), of approximately 3,416 cases per 100,000, was more than double that in the SPA with the second highest rate in the County (SPA 8, South Bay), and five times that in the SPA with the lowest rate (SPA 5, West).
- The 2005 rate for gonorrhea among 15-19 year-olds in SPA 6 (South), of 1068 cases per 100,000, was more than double that in the SPA with the second highest rate in the County (SPA 8, South Bay), and almost ten times higher than that in the SPA with the lowest rate (SPA 5, West).

CONTENTS

Table 1-1:	Sexually Transmitted Diseases Reported for 10-24 Year Olds in Los Angeles County, 2001-2005.	1
Table 1-2:	Sexually Transmitted Diseases Reported for 10-24 Year Olds in Los Angeles County by Age Group, 2001-2005.	1a

Chlamydia

Figure 1:	Reported Chlamydia Rates for 10-14 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	2
Table 2:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	3
Table 3:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005.	4
Table 4-1:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	5
Table 4-2:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	6
Table 4-3:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	7
Table 4-4:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	8
Table 4-5:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	9
Table 4-6:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	10
Table 4-7:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	11
Table 4-8:	Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	12
Figure 2:	Reported Chlamydia Rates for 15-19 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	13
Table 5:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	14
Table 6:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005.	15
Table 7-1:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	16
Table 7-2:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	17

Table 7-3:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	18
Table 7-4:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	19
Table 7-5:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	20
Table 7-6:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	21
Table 7-7:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	22
Table 7-8:	Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	23
Figure 3:	Reported Chlamydia Rates for 20-24 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	24
Table 8:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	25
Table 9:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005.	26
Table 10-1:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	27
Table 10-2:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	28
Table 10-3:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	29
Table 10-4:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	30
Table 10-5:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	31
Table 10-6:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	32
Table 10-7:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	33
Table 10-8:	Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	34
Figure 4:	2005 Reported Chlamydia Rates Per 100,000 10-24 Year Olds, by Service Planning Area (SPA)	35

Gonorrhea

Figure 5:	Reported Gonorrhea Rates for 10-14 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	36
Table 11:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	37
Table 12:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005. . .	38
Table 13-1:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	39
Table 13-2:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	40
Table 13-3:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	41
Table 13-4:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	42
Table 13-5:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	43
Table 13-6:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	44
Table 13-7:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	45
Table 13-8:	Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	46
Figure 6:	Reported Gonorrhea Rates for 15-19 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	47
Table 14:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	48
Table 15:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005. . .	49
Table 16-1:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	50
Table 16-2:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	51
Table 16-3:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	52
Table 16-4:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	53
Table 16-5:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	54

Table 16-6:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	55
Table 16-7:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	56
Table 16-8:	Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	57
Figure 7:	Reported Gonorrhea Rates for 20-24 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	58
Table 17:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	59
Table 18:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005.	60
Table 19-1:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	61
Table 19-2:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	62
Table 19-3:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	63
Table 19-4:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	64
Table 19-5:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	65
Table 19-6:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	66
Table 19-7:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	67
Table 19-8:	Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	68
Figure 8:	2005 Reported Gonorrhea Rates Per 100,000 10-24 Year Olds, by Service Planning Area (SPA)	69

Early Syphilis

Figure 9:	Reported Early Syphilis Rates for 10-14 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	70
Table 20:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	71
Table 21:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005.	72
Table 22-1:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	73

Table 22-2:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	74
Table 22-3:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	75
Table 22-4:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	76
Table 22-5:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	77
Table 22-6:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	78
Table 22-7:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	79
Table 22-8:	Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	80
Figure 10:	Reported Early Syphilis Rates for 15-19 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	81
Table 23:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	82
Table 24:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005.	83
Table 25-1:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	84
Table 25-2:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	85
Table 25-3:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	86
Table 25-4:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	87
Table 25-5:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	88
Table 25-6:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	89
Table 25-7:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	90
Table 25-8:	Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	91
Figure 11:	Reported Early Syphilis Rates for 20-24 Year Olds by Year, United States, California, and Los Angeles County (LAC), 2001-2005.	92
Table 26:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by	

Table 27:	Gender and Race/Ethnicity, Los Angeles County, 2001-2005.	93
Table 28-1:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Service Planning Area (SPA) and Gender, Los Angeles County, 2001-2005. . .	94
Table 28-2:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005.	95
Table 28-3:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005.	96
Table 28-4:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005.	97
Table 28-5:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005.	98
Table 28-6:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005.	99
Table 28-7:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005.	100
Table 28-8:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005.	101
Figure 12:	Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005.	102
Figure 12:	2005 Reported Early Syphilis Rates Per 100,000 10-24 Year Olds, by Service Planning Area (SPA)	103

Special Projects

Special Project Summary.	104
----------------------------------	-----

Appendices

Appendix A. Case Definitions for Reportable STDs.	111
Appendix B. Technical Notes.	114
Appendix C. STD Program Surveillance Report Request.	115
Appendix D. Where to Report Communicable Diseases in Los Angeles County.	116

Table 1-1. Sexually Transmitted Diseases Reported for 10-24 Year Olds in Los Angeles County, 2001-2005

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Chlamydia ¹	20,386	84.5	991.0	21,859	85.3	1,034.6	22,525	85.1	1,035.3	24,097	83.2	1,087.2	24,592	81.8	1,154.7
Gonorrhea ¹	3,685	15.3	179.1	3,675	14.3	173.9	3,849	14.5	176.9	4,756	16.4	214.6	5,320	17.7	249.8
Early Syphilis ^{1,2}	51	0.2	2.5	95	0.4	4.5	84	0.3	3.9	118	0.4	5.3	151	0.5	7.1
Total	24,122			25,629			26,458			28,971			30,063		

1. Rates per 100,000 population.

2. Primary, secondary, and early latent syphilis cases.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.


**Table 1-2. Sexually Transmitted Diseases Reported for 10-24 Year Olds in Los Angeles County
by Age Group, 2001-2005**

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
<u>Chlamydia</u>															
Age Group															
10-14	397	1.9	54.4	421	1.9	55.4	346	1.5	44.1	391	1.6	49.6	381	1.5	49.7
15-19	8,984	44.1	1,330.6	9,729	44.5	1,397.3	9,725	43.2	1,348.6	10,532	43.7	1,410.5	10,823	44.0	1,543.8
20-24	<u>11,005</u>	54.0	1,688.8	<u>11,709</u>	53.6	1,784.8	<u>12,454</u>	55.3	1,857.0	<u>13,174</u>	54.7	1,933.6	<u>13,388</u>	54.4	2,020.0
Total	<u>20,386</u>		991.0	<u>21,859</u>		1,034.6	<u>22,525</u>		1,035.3	<u>24,097</u>		1,087.2	<u>24,592</u>		1,154.7
<u>Gonorrhea</u>															
Age Group															
10-14	87	2.4	11.9	84	2.3	11.0	84	2.2	10.7	106	2.2	13.5	109	2.0	13.9
15-19	1,589	43.1	235.3	1,620	44.1	232.7	1,702	44.2	236.0	2,035	42.8	282.2	2,346	44.1	325.3
20-24	<u>2,009</u>	54.5	308.3	<u>1,971</u>	53.6	300.4	<u>2,063</u>	53.6	307.6	<u>2,615</u>	55.0	389.9	<u>2,865</u>	53.9	427.2
Total	<u>3,685</u>		179.1	<u>3,675</u>		173.9	<u>3,849</u>		176.9	<u>4,756</u>		218.6	<u>5,320</u>		244.5
<u>Early Syphilis¹</u>															
Age Group															
10-14	3	5.9	0.4	1	1.1	0.1	0	0.0	0.0	2	1.7	0.3	0	0.0	0.0
15-19	14	27.5	2.1	21	22.1	3.0	17	20.2	2.4	21	17.8	2.9	29	19.2	4.0
20-24	<u>34</u>	66.7	5.2	<u>73</u>	76.8	11.1	<u>67</u>	79.8	10.0	<u>95</u>	80.5	14.2	<u>122</u>	80.8	18.2
Total	<u>51</u>		2.5	<u>95</u>		4.5	<u>84</u>		3.9	<u>118</u>		5.4	<u>151</u>		6.9
Total	24,122		1,172.6	25,629		1,213.0	26,458		1,321.9	28,971		1,408.3	30,063		1,422.9

1. Primary, secondary, and early latent syphilis cases.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Figure 1. Reported Chlamydia Rates for 10-14 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 2. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	2	3.3	2.8	33	54.1	99.9	16	26.2	8.9	1	1.6	3.0	0	0.0	----	61	15.4	16.4
2002	2	3.2	2.9	19	30.6	59.1	29	46.8	15.8	1	1.6	3.0	0	0.0	----	62	14.8	16.0
2003	6	11.3	8.5	23	43.4	69.4	16	30.2	8.0	0	0.0	0.0	0	0.0	----	53	100.0	13.2
2004	0	0.0	0.0	36	54.5	109.1	21	31.8	10.1	0	0.0	0.0	0	0.0	----	66	100.0	16.4
2005	1	1.9	1.4	23	44.2	72.0	20	38.5	10.1	0	0.0	0.0	0	0.0	----	52	100.0	13.3
FEMALE																		
2001	12	3.6	19.8	116	34.5	398.8	124	36.9	80.7	3	0.9	10.7	1	0.3	----	336	84.6	94.0
2002	12	3.4	19.0	125	34.9	409.6	142	39.7	84.2	2	0.6	6.7	1	0.3	----	358	85.2	96.1
2003	15	5.1	21.9	106	36.3	316.8	125	42.8	64.6	5	1.7	15.0	0	0.0	----	292	84.6	76.1
2004	17	5.2	25.7	130	40.0	393.0	130	40.0	65.1	4	1.2	11.9	0	0.0	----	325	83.1	84.3
2005	23	7.0	33.8	129	39.3	400.5	135	41.2	68.7	0	0.0	0.0	1	0.3	----	328	86.3	87.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2002; 1 in 2003; 1 in 2004; and 1 in 2005.

3. Includes 89 cases reported as unknown race/ethnicity in 2001; 87 in 2002; 49 in 2003; 53 in 2004; and 48 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 3. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

SPA ² /Gender	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Antelope Valley (1)	21	5.3	64.2	21	5.0	62.7	35	10.1	103.1	30	7.7	88.2	33	8.7	100.4
Male	2	0.5	12.1	3	0.7	17.7	8	0.0	46.9	7	1.8	41.0	2	0.5	12.1
Female	19	4.8	118.0	18	4.3	108.6	27	7.8	160.1	23	5.9	135.8	31	8.1	190.5
East (7)	49	12.3	41.8	45	10.7	36.9	32	9.2	25.5	34	8.7	26.8	29	7.6	23.5
Male	5	1.3	8.3	5	1.2	8.0	5	1.4	7.8	3	0.8	4.6	2	0.5	3.2
Female	44	11.1	76.8	40	9.5	67.1	27	7.8	43.9	31	7.9	50.0	27	7.1	44.7
Metro (4)	27	6.8	34.2	41	9.7	48.7	41	11.8	46.4	39	10.0	43.4	42	11.0	48.0
Male	2	0.5	5.0	6	1.4	13.9	4	1.2	8.8	3	0.8	6.5	6	1.6	13.4
Female	25	6.3	64.8	35	8.3	85.3	37	10.7	86.0	36	9.2	82.5	36	9.4	83.9
San Fernando (2)	39	9.8	25.2	43	10.2	26.6	37	10.7	22.2	32	8.2	19.0	45	11.8	27.6
Male	3	0.8	3.8	6	1.4	7.3	7	2.0	8.2	8	2.0	9.3	6	1.6	7.2
Female	36	9.1	47.6	37	8.8	46.9	30	8.7	36.9	24	6.1	29.3	39	10.2	49.1
San Gabriel (3)	36	9.1	27.0	37	8.8	27.1	21	6.1	15.2	31	7.9	22.6	30	7.9	22.3
Male	3	0.8	4.4	8	1.9	11.5	1	0.3	1.4	4	1.0	5.7	1	0.3	1.5
Female	33	8.3	50.4	29	6.9	43.4	20	5.8	29.6	27	6.9	40.3	29	7.6	44.1
South (6)³	135	34.0	138.0	138	32.8	135.7	118	34.1	112.5	157	40.2	148.8	144	37.8	137.8
Male	30	7.6	60.3	21	5.0	40.5	15	4.3	28.0	27	6.9	50.1	22	5.8	41.2
Female	105	26.4	218.5	116	27.6	232.6	103	29.8	200.4	130	33.2	251.6	121	31.8	236.8
South Bay (8)	45	11.3	53.6	52	12.4	59.2	34	9.8	37.5	32	8.2	35.1	30	7.9	34.0
Male	6	1.5	14.0	6	1.4	13.4	5	1.4	10.8	10	2.6	21.6	4	1.0	8.9
Female	39	9.8	95.0	46	10.9	106.8	29	8.4	65.0	22	5.6	49.0	26	6.8	60.1
West (5)	6	1.5	19.2	6	1.4	17.9	5	1.4	14.2	6	1.5	16.9	2	0.5	6.4
Male	2	0.5	12.6	0	0.0	0.0	1	0.3	5.6	0	0.0	0.0	0	0.0	0.0
Female	4	1.0	25.9	6	1.4	36.1	4	1.2	23.0	6	1.5	34.1	2	0.5	13.1
Unknown	39	9.8	-----	38	9.0	-----	23	6.6	-----	30	7.7	-----	26	6.8	-----
Total	397		54.4	421		55.4	346		44.1	391		49.6	381		49.7

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 1 case reported with "unknown/other" gender in 2002; 1 in 2003; 1 in 2004; and 1 in 2005.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-1. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	50.0	33.9	0	0.0	0.0	0	0.0	----	2	9.5	12.1
2002	1	33.3	14.3	1	33.3	32.0	1	33.3	15.9	0	0.0	0.0	0	0.0	----	3	14.3	17.7
2003	2	25.0	34.2	4	50.0	139.2	1	12.5	17.2	0	0.0	0.0	0	0.0	----	8	22.9	46.9
2004	0	0.0	0.0	7	100.0	202.5	0	0.0	0.0	0	0.0	0.0	0	0.0	----	7	23.3	41.0
2005	0	0.0	0.0	0	0.0	0.0	1	50.0	31.4	0	0.0	0.0	0	0.0	----	2	6.1	12.1
FEMALE																		
2001	2	10.5	39.1	8	42.1	384.1	4	21.1	93.7	0	0.0	0.0	0	0.0	----	19	90.5	118.0
2002	2	11.1	37.7	10	55.6	424.7	1	5.6	20.8	1	5.6	282.6	0	0.0	----	18	85.7	108.6
2003	5	18.5	97.8	11	40.7	437.6	4	14.8	78.7	1	3.7	293.5	0	0.0	----	27	77.1	160.1
2004	3	13.0	59.1	10	43.5	353.0	6	26.1	105.7	0	0.0	0.0	0	0.0	----	23	76.7	135.8
2005	4	12.9	71.6	19	61.3	907.6	2	6.5	40.0	0	0.0	0.0	0	0.0	----	31	93.9	190.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 6 cases reported as unknown race/ethnicity in 2001; 4 in 2002; 7 in 2003; 4 in 2004; and 7 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-2. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	33.3	3.2	1	33.3	30.6	1	33.3	2.7	0	0.0	0.0	0	0.0	----	3	7.7	3.8
2002	1	16.7	3.2	0	0.0	0.0	4	66.7	10.1	1	16.7	12.9	0	0.0	----	6	14.0	7.3
2003	1	14.3	3.7	1	14.3	33.3	4	57.1	11.1	0	0.0	0.0	0	0.0	----	7	18.9	8.2
2004	0	0.0	0.0	4	50.0	112.8	4	50.0	9.2	0	0.0	0.0	0	0.0	----	8	25.0	9.3
2005	1	16.7	3.8	1	16.7	37.6	3	50.0	8.8	0	0.0	0.0	0	0.0	----	6	13.3	7.2
FEMALE																		
2001	3	8.3	10.9	6	16.7	210.9	23	63.9	71.1	1	2.8	15.7	0	0.0	----	36	92.3	47.6
2002	0	0.0	0.0	7	18.9	252.0	25	67.6	76.2	0	0.0	0.0	0	0.0	----	37	86.0	46.9
2003	6	20.0	21.2	4	13.3	130.1	18	60.0	47.9	0	0.0	0.0	0	0.0	----	30	81.1	36.9
2004	2	8.3	8.6	2	8.3	75.5	14	58.3	42.5	1	4.2	17.3	0	0.0	----	24	75.0	29.3
2005	7	17.9	24.5	8	20.5	285.5	22	56.4	58.9	0	0.0	0.0	0	0.0	----	39	86.7	49.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 3 cases reported as unknown race/ethnicity in 2001; 5 in 2002; 3 in 2003; 5 in 2004; and 3 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-3. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	33.3	8.2	1	33.3	31.6	1	33.3	2.7	0	0.0	0.0	0	0.0	----	3	8.3	4.4
2002	0	0.0	0.0	2	25.0	74.5	5	62.5	14.6	0	0.0	0.0	0	0.0	----	8	21.6	11.5
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	4.8	1.4
2004	0	0.0	0.0	3	75.0	103.8	1	25.0	2.4	0	0.0	0.0	0	0.0	----	4	12.9	5.7
2005	0	0.0	0.0	0	0.0	0.0	1	100.0	2.5	0	0.0	0.0	0	0.0	----	1	3.3	1.5
FEMALE																		
2001	3	9.1	30.9	5	15.2	203.4	18	54.5	60.6	1	3.0	8.7	0	0.0	----	33	91.7	50.4
2002	2	6.9	21.3	7	24.1	285.9	14	48.3	44.8	1	3.4	8.3	0	0.0	----	29	78.4	43.4
2003	1	5.0	10.3	1	5.0	38.9	14	70.0	39.9	2	10.0	15.1	0	0.0	----	20	95.2	29.6
2004	3	11.1	38.5	4	14.8	188.2	14	51.9	45.8	0	0.0	0.0	0	0.0	----	27	87.1	40.3
2005	3	10.3	31.3	8	27.6	316.2	14	48.3	42.0	0	0.0	0.0	0	0.0	----	29	96.7	44.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 6 cases reported as unknown race/ethnicity in 2001; 6 in 2002; 3 in 2003; 6 in 2004; and 4 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 4-4. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	50.0	7.1	0	0.0	0.0	0	0.0	----	2	7.4	5.0
2002	0	0.0	0.0	0	0.0	0.0	5	83.3	19.8	0	0.0	0.0	0	0.0	----	6	14.6	13.9
2003	1	25.0	17.7	1	25.0	49.5	2	50.0	6.2	0	0.0	0.0	0	0.0	----	4	9.8	8.8
2004	0	0.0	0.0	2	66.7	101.0	1	33.3	3.0	0	0.0	0.0	0	0.0	----	3	7.7	6.5
2005	0	0.0	0.0	2	33.3	99.9	4	66.7	12.1	0	0.0	0.0	0	0.0	----	6	14.6	13.4
FEMALE																		
2001	1	4.0	35.1	3	12.0	269.0	11	44.0	67.7	0	0.0	0.0	0	0.0	----	25	92.6	64.8
2002	1	2.9	25.2	7	20.0	473.3	19	54.3	85.5	0	0.0	0.0	0	0.0	----	35	85.4	85.3
2003	0	0.0	0.0	8	21.6	517.1	21	56.8	88.2	0	0.0	0.0	0	0.0	----	37	90.2	86.0
2004	1	2.8	22.0	5	13.9	291.9	25	69.4	93.8	0	0.0	0.0	0	0.0	----	36	92.3	82.5
2005	3	8.6	82.9	4	11.4	233.2	24	68.6	85.3	0	0.0	0.0	0	0.0	----	35	85.4	81.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 11 cases reported as unknown race/ethnicity in 2001; 9 in 2002; 8 in 2003; 5 in 2004; and 4 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-5. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 1998-2002

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	2	100.0	137.5	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	33.3	12.6
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	1	0.0	0.0	0	0.0	0.0	0	0.0	----	1	20.0	5.6
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	1	25.0	96.1	1	25.0	36.5	1	25.0	96.1	0	0.0	----	4	66.7	25.9
2002	2	33.3	31.3	0	0.0	0.0	2	33.3	75.5	0	0.0	0.0	0	0.0	----	6	100.0	36.1
2003	0	0.0	0.0	2	50.0	134.4	2	50.0	47.1	0	0.0	0.0	0	0.0	----	4	80.0	23.0
2004	0	0.0	0.0	5	83.3	330.7	1	16.7	22.9	0	0.0	0.0	0	0.0	----	6	100.0	34.1
2005	0	0.0	0.0	1	50.0	76.3	1	50.0	23.2	0	0.0	0.0	0	0.0	----	2	100.0	13.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2001; and 2 in 2002.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-6. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	21	70.0	147.9	6	20.0	19.2	1	3.3	270.6	0	0.0	----	30	22.2	60.3
2002	0	0.0	0.0	12	57.1	93.5	6	28.6	19.6	0	0.0	0.0	0	0.0	----	21	15.3	40.5
2003	1	6.7	144.5	11	73.3	75.4	3	20.0	8.0	0	0.0	0.0	0	0.0	----	15	12.7	28.0
2004	0	0.0	0.0	14	51.9	114.5	10	37.0	28.9	0	0.0	0.0	0	0.0	----	27	17.2	50.1
2005	0	0.0	0.0	16	72.7	105.0	6	27.3	16.1	0	0.0	0.0	0	0.0	----	22	15.4	41.2
FEMALE																		
2001	0	0.0	0.0	63	60.0	506.6	22	21.0	84.6	0	0.0	0.0	1	1.0	----	105	77.8	218.5
2002	0	0.0	0.0	62	53.4	515.1	31	26.7	114.4	0	0.0	0.0	0	0.0	----	116	84.7	232.6
2003	0	0.0	0.0	59	57.3	440.4	34	33.0	104.6	1	1.0	286.1	0	0.0	----	103	87.3	200.4
2004	1	0.8	161.9	86	66.2	670.7	33	25.4	97.6	0	0.0	0.0	0	0.0	----	130	82.8	251.6
2005	4	3.3	787.3	66	54.5	493.2	39	32.2	122.9	0	0.0	0.0	0	0.0	----	121	84.6	236.8

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2002; 1 in 2004; and 1 in 2005.

3. Includes 21 cases reported as unknown race/ethnicity in 2001; 26 in 2002; 9 in 2003; 13 in 2004; and 12 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-7. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	1	20.0	46.8	4	80.0	8.7	0	0.0	0.0	0	0.0	----	5	10.2	8.3
2002	0	0.0	0.0	1	20.0	45.4	4	80.0	8.3	0	0.0	0.0	0	0.0	----	5	11.1	8.0
2003	0	0.0	0.0	1	20.0	56.0	3	60.0	7.4	0	0.0	0.0	0	0.0	----	5	15.6	7.8
2004	0	0.0	0.0	1	33.3	44.0	2	66.7	3.9	0	0.0	0.0	0	0.0	----	3	8.8	4.6
2005	0	0.0	0.0	0	0.0	0.0	2	100.0	4.0	0	0.0	0.0	0	0.0	----	2	6.9	3.2
FEMALE																		
2001	2	4.5	33.4	3	6.8	167.8	32	72.7	86.6	0	0.0	0.0	0	0.0	----	44	89.8	76.8
2002	0	0.0	0.0	6	15.0	289.3	32	80.0	72.6	0	0.0	0.0	0	0.0	----	40	88.9	67.1
2003	2	7.4	36.6	3	11.1	157.9	18	66.7	43.4	0	0.0	0.0	0	0.0	----	27	84.4	43.9
2004	2	6.5	38.8	2	6.5	101.5	23	74.2	53.1	0	0.0	0.0	0	0.0	----	31	91.2	50.0
2005	2	7.4	30.8	1	3.7	54.9	23	85.2	49.9	0	0.0	0.0	0	0.0	----	27	93.1	44.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 7 cases reported as unknown race/ethnicity in 2001; 2 in 2002; 5 in 2003; 4 in 2004; and 1 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 4-8. Reported Chlamydia Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	4	66.7	46.6	2	33.3	10.5	0	0.0	0.0	0	0.0	----	6	13.3	14.0
2002	0	0.0	0.0	3	50.0	34.1	3	50.0	14.7	0	0.0	0.0	0	0.0	----	6	11.5	13.4
2003	0	0.0	0.0	4	80.0	45.2	1	20.0	4.6	0	0.0	0.0	0	0.0	----	5	14.7	10.8
2004	0	0.0	0.0	4	40.0	65.5	3	30.0	19.3	0	0.0	0.0	0	0.0	----	10	31.3	21.6
2005	0	0.0	0.0	3	75.0	36.4	1	25.0	4.7	0	0.0	0.0	0	0.0	----	4	13.3	8.9
FEMALE																		
2001	1	2.6	12.9	22	56.4	308.5	9	23.1	61.0	0	0.0	0.0	0	0.0	----	39	86.7	95.0
2002	4	8.7	48.0	20	43.5	255.8	15	32.6	89.6	0	0.0	0.0	1	2.2	----	46	88.5	106.8
2003	1	3.4	12.1	15	51.7	189.7	9	31.0	50.7	0	0.0	0.0	0	0.0	----	29	85.3	65.0
2004	3	13.6	32.3	11	50.0	121.0	8	36.4	37.5	0	0.0	0.0	0	0.0	----	22	68.8	49.0
2005	0	0.0	0.0	17	65.4	233.7	6	23.1	33.0	0	0.0	0.0	0	0.0	----	26	86.7	60.1


1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 7 cases reported as unknown race/ethnicity in 2001; 6 in 2002; 4 in 2003; 3 in 2004; and 3 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.

Figure 2. Reported Chlamydia Rates for 15-19 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 5. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	69	4.2	108.7	509	30.8	1954.1	686	41.5	465.1	31	1.9	91.1	4	0.2	----	1,655	18.4	477.6
2002	75	3.8	113.7	640	32.2	2368.4	822	41.3	537.7	32	1.6	90.9	8	0.4	----	1,989	20.5	559.0
2003	86	4.3	127.1	671	33.7	2414.0	791	39.8	494.2	33	1.7	91.0	5	0.3	----	1,989	100.0	541.7
2004	117	5.4	169.3	776	36.1	2721.5	798	37.1	475.6	25	1.2	67.2	5	0.2	----	2,149	100.0	566.8
2005	87	3.8	130.7	877	38.7	2955.7	820	36.2	520.8	33	1.5	101.9	4	0.2	----	2,268	100.0	634.3
FEMALE																		
2001	426	5.8	775.0	1,768	24.1	7519.4	2,831	38.7	2255.4	172	2.3	589.2	19	0.3	----	7,323	81.6	2228.0
2002	462	6.0	762.4	1,959	25.3	7537.6	3,210	41.5	2320.3	196	2.5	615.7	25	0.3	----	7,730	79.5	2270.5
2003	543	7.0	805.8	2,247	29.1	7682.6	3,336	43.1	2130.8	181	2.3	513.6	15	0.2	----	7,734	79.5	2185.0
2004	564	6.7	830.4	2,509	30.0	8432.7	3,457	41.3	2142.9	167	2.0	469.7	43	0.5	----	8,377	79.6	2279.3
2005	596	7.0	892.3	2,683	31.4	8937.1	3,603	42.2	2312.4	167	2.0	531.7	43	0.5	----	8,545	79.0	2487.3

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 6 cases reported with "unknown/other" gender in 2001; 10 in 2002; 9 in 2003; 10 in 2004; and 12 in 2005.

3. Includes 2,463 cases reported as unknown race/ethnicity in 2001; 2,290 in 2002; 1,815 in 2003; 2,065 in 2004; and 1,900 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 6. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

SPA ² /Gender	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
Antelope Valley (1)³	330	3.7	1148.5	360	3.7	1202.1	477	4.9	1547.3	503	4.8	1585.7	553	5.1	1646.7
Male	51	0.6	341.1	64	0.7	413.6	105	1.1	664.2	96	0.9	595.1	109	1.0	626.8
Female	279	3.1	2024.2	296	3.0	2045.0	372	3.8	2476.9	407	3.9	2610.5	444	4.1	2742.1
East (7)³	1,217	13.5	1146.0	1,294	13.3	1197.1	1,212	12.5	1094.3	1,249	11.9	1098.3	1,289	11.9	1162.3
Male	204	2.3	372.7	229	2.4	412.5	206	2.1	363.0	194	1.8	333.7	232	2.1	406.5
Female	1,013	11.3	1968.7	1,064	10.9	2023.7	1006	10.3	1862.8	1,055	10.0	1898.0	1,056	9.8	1962.0
Metro (4)³	973	10.8	1314.3	1,166	12.0	1515.0	1,076	11.1	1324.3	1,146	10.9	1327.3	1,060	9.8	1461.7
Male	175	1.9	453.7	250	2.6	626.5	195	2.0	464.9	278	2.6	624.4	223	2.1	600.3
Female	798	8.9	2250.7	914	9.4	2466.1	881	9.1	2241.4	868	8.2	2075.9	835	7.7	2360.8
San Fernando (2)³	1,097	12.2	774.0	1,259	12.9	852.6	1,247	12.8	812.6	1,276	12.1	800.2	1,315	12.2	890.9
Male	172	1.9	235.5	230	2.4	304.2	234	2.4	299.1	225	2.1	277.5	244	2.3	322.7
Female	924	10.3	1345.2	1,029	10.6	1427.7	1,013	10.4	1346.9	1,050	10.0	1339.7	1,071	9.9	1487.7
San Gabriel (3)³	1,015	11.3	792.0	1,055	10.8	815.8	1,107	11.4	846.2	1,133	10.8	858.7	1,188	11.0	906.9
Male	198	2.2	299.6	198	2.0	297.6	210	2.2	313.0	233	2.2	345.6	206	1.9	309.2
Female	816	9.1	1314.6	857	8.8	1364.9	897	9.2	1407.3	899	8.5	1393.6	982	9.1	1525.5
South (6)³	2,352	26.2	2697.8	2,853	29.3	3201.5	2,955	30.4	3210.2	3,075	29.2	3234.8	3,186	29.4	3415.7
Male	471	5.2	1073.4	661	6.8	1476.0	649	6.7	1403.3	658	6.2	1376.5	717	6.6	1522.0
Female	1,881	20.9	4343.7	2,190	22.5	4940.0	2,306	23.7	5034.9	2,417	22.9	5114.6	2,467	22.8	5343.6
South Bay (8)³	1,023	11.4	1406.4	1,123	11.5	1480.8	1,055	10.8	1326.7	1,188	11.3	1433.1	1,295	12.0	1664.1
Male	205	2.3	550.4	229	2.4	592.1	202	2.1	500.5	252	2.4	601.0	277	2.6	697.0
Female	817	9.1	2301.8	893	9.2	2403.1	853	8.8	2178.4	935	8.9	2282.3	1,017	9.4	2670.9
West (5)	175	1.9	480.3	206	2.1	524.2	225	2.3	530.2	217	2.1	476.6	235	2.2	683.3
Male	36	0.4	199.9	33	0.3	171.0	36	0.4	173.9	38	0.4	171.9	43	0.4	255.5
Female	139	1.5	754.3	173	1.8	865.3	189	1.9	869.8	179	1.7	764.3	192	1.8	1093.6
Unknown	802	8.9	-----	413	4.2	-----	371	3.8	-----	745	7.1	-----	702	6.5	-----
Total	8,984		1330.6	9,729		1397.3	9,725		1348.6	10,532		1410.5	10,823		1543.8

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 3 cases reported with "unknown/other" gender in 2001; 6 in 2002; 9 in 2003; 10 in 2004; and 12 in 2005.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-1. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	8	15.7	153.3	25	49.0	1,336.5	7	13.7	168.6	0	0.0	0.0	0	0.0	----	51	15.5	341.1
2002	6	9.4	123.3	31	48.4	1,602.0	10	15.6	242.9	0	0.0	0.0	0	0.0	----	64	17.8	413.6
2003	10	9.5	188.4	44	41.9	1,882.4	30	28.6	625.3	1	1.0	242.3	1	1.0	----	105	22.0	664.2
2004	8	8.3	162.0	52	54.2	2,167.4	15	15.6	310.3	0	0.0	0.0	0	0.0	----	96	19.1	595.1
2005	5	4.6	81.8	65	59.6	2,854.6	14	12.8	289.1	2	1.8	500.0	0	0.0	----	109	19.7	626.8
FEMALE																		
2001	38	13.6	975.2	89	31.9	6,419.3	43	15.4	1,488.0	1	0.4	396.5	1	0.4	----	279	84.5	2,024.2
2002	45	15.2	1,163.0	89	30.1	5,805.5	42	14.2	1,354.5	4	1.4	1,628.2	0	0.0	----	296	82.2	2,045.0
2003	68	18.3	1,563.8	122	32.8	6,402.9	67	18.0	1,774.6	0	0.0	0.0	0	0.0	----	372	78.0	2,476.9
2004	66	16.2	1,359.0	162	39.8	6,936.2	81	19.9	1,753.4	7	1.7	2,231.8	3	0.7	----	407	80.9	2,610.5
2005	83	18.7	1,371.7	179	40.3	8,332.4	97	21.8	2,122.9	3	0.7	746.0	2	0.5	----	444	80.3	2,742.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2003; and 1 in 2004.

3. Includes 118 cases reported as unknown race/ethnicity in 2001; 133 in 2002; 134 in 2003; 109 in 2004; and 103 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-2. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	14	8.1	60.1	20	11.6	794.9	100	58.1	378.8	4	2.3	64.0	0	0.0	----	172	15.7	235.5
2002	24	10.4	100.7	30	13.0	1,153.2	121	52.6	447.1	6	2.6	94.4	2	0.9	----	230	18.3	304.2
2003	28	12.0	115.8	34	14.5	1,260.4	114	48.7	402.3	8	3.4	125.0	1	0.4	----	234	18.8	299.1
2004	31	13.8	121.4	41	18.2	1,389.6	109	48.4	350.2	3	1.3	44.0	1	0.4	----	225	17.6	277.5
2005	23	9.4	90.1	37	15.2	1,619.4	133	54.5	483.5	5	2.0	81.7	1	0.4	----	244	18.6	322.7
FEMALE																		
2001	96	10.4	441.4	81	8.8	3,332.9	525	56.8	2,139.4	31	3.4	540.1	2	0.2	----	924	84.3	1,345.2
2002	136	13.2	618.1	109	10.6	4,342.3	518	50.3	2,069.6	25	2.4	433.2	5	0.5	----	1,029	81.7	1,427.7
2003	170	16.8	692.8	94	9.3	3,269.1	551	54.4	1,912.5	31	3.1	482.1	0	0.0	----	1,013	81.2	1,346.9
2004	165	15.7	681.3	110	10.5	3,755.8	538	51.2	1,825.7	23	2.2	359.2	13	1.2	----	1,050	82.4	1,339.7
2005	154	14.4	615.5	114	10.6	5,015.0	593	55.4	2,184.2	35	3.3	599.4	4	0.4	----	1,071	81.4	1,487.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001; and 2 in 2004.

3. Includes 223 cases reported as unknown race/ethnicity in 2001; 283 in 2002; 216 in 2003; 241 in 2004; and 216 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-3. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	19	9.6	162.9	26	13.1	896.7	124	62.6	407.9	10	5.1	67.1	1	0.5	----	198	19.5	299.6
2002	10	5.1	93.0	31	15.7	1,146.7	124	62.6	418.3	7	3.5	48.2	0	0.0	----	198	18.8	297.6
2003	14	6.7	150.8	26	12.4	1,124.8	114	54.3	413.8	11	5.2	82.0	0	0.0	----	210	19.0	313.0
2004	19	8.2	199.3	37	15.9	1,568.5	135	57.9	442.1	6	2.6	41.0	1	0.4	----	233	20.6	345.6
2005	13	6.3	116.9	31	15.0	1,207.3	126	61.2	409.1	9	4.4	68.0	0	0.0	----	206	17.3	309.2
FEMALE																		
2001	98	12.0	984.2	98	12.0	4,318.1	436	53.4	1,646.2	41	5.0	329.0	2	0.2	----	816	80.5	1,314.6
2002	76	8.9	807.4	73	8.5	3,358.7	485	56.6	1,834.8	54	6.3	436.9	2	0.2	----	857	81.2	1,364.9
2003	75	8.4	780.1	103	11.5	4,617.3	523	58.3	1,818.7	55	6.1	416.1	4	0.4	----	897	81.0	1,407.3
2004	82	9.1	880.3	108	12.0	4,951.4	528	58.7	1,780.5	43	4.8	319.9	2	0.2	----	899	79.4	1,393.6
2005	97	9.9	874.8	107	10.9	4,439.1	583	59.4	1,963.6	52	5.3	433.3	6	0.6	----	982	82.7	1,525.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001; 1 in 2003; and 1 in 2005.

3. Includes 159 cases reported as unknown race/ethnicity in 2001; 193 in 2002; 182 in 2003; 171 in 2004; and 164 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 7-4. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 1998-2002

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	10	5.7	229.8	20	11.4	1,247.3	115	65.7	520.5	3	1.7	66.9	0	0.0	----	175	18.0	453.7
2002	3	1.2	66.4	29	11.6	1,965.5	151	60.4	750.5	6	2.4	142.6	2	0.8	----	250	21.5	626.5
2003	2	1.0	35.8	27	13.8	1,628.7	128	65.6	568.6	3	1.5	63.4	1	0.5	----	195	18.1	464.9
2004	12	4.3	198.7	39	14.0	2,315.9	161	57.9	712.1	8	2.9	167.0	0	0.0	----	278	24.3	624.4
2005	8	3.6	250.7	38	17.0	2,355.1	127	57.0	593.1	8	3.6	202.2	1	0.4	----	223	21.1	600.3
FEMALE																		
2001	33	4.1	934.9	114	14.3	8,817.2	399	50.0	2,388.0	23	2.9	619.4	1	0.1	----	798	82.0	2,250.7
2002	28	3.1	644.1	127	13.9	8,719.1	510	55.8	2,761.7	36	3.9	870.2	2	0.2	----	914	78.5	2,466.1
2003	23	2.6	457.3	141	16.0	9,082.7	504	57.2	2,556.9	19	2.2	434.8	3	0.3	----	881	81.9	2,241.4
2004	31	3.6	540.7	127	14.6	7,701.0	511	58.9	2,406.5	21	2.4	450.2	4	0.5	----	868	75.7	2,075.9
2005	34	4.1	1,040.0	128	15.3	8,842.2	520	62.3	2,413.2	26	3.1	677.3	6	0.7	----	835	78.9	2,360.8

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2002; 1 in 2003; 3 in 2004; and 1 in 2005.

3. Includes 255 cases reported as unknown race/ethnicity in 2001; 270 in 2002; 225 in 2003; 232 in 2004; and 162 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-5. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	3	8.3	37.4	15	41.7	1,324.3	10	27.8	291.2	1	2.8	43.5	1	2.8	----	36	20.6	199.9
2002	1	3.0	14.0	10	30.3	1,050.4	9	27.3	319.0	2	6.1	102.8	0	0.0	----	33	16.0	171.0
2003	5	13.9	53.4	13	36.1	1,080.2	10	27.8	282.7	1	2.8	39.5	0	0.0	----	36	16.0	173.9
2004	12	31.6	104.4	14	36.8	957.8	8	21.1	188.2	1	2.6	32.0	0	0.0	----	38	17.5	171.9
2005	3	7.0	43.2	16	37.2	1,625.1	11	25.6	369.6	2	4.7	126.2	0	0.0	----	43	18.3	255.5
FEMALE																		
2001	21	15.1	280.1	40	28.8	3,810.0	37	26.6	1,278.6	6	4.3	237.5	2	1.4	----	139	79.4	754.3
2002	33	19.1	404.0	41	23.7	3,682.7	44	25.4	1,484.4	10	5.8	382.7	1	0.6	----	173	84.0	865.3
2003	47	24.9	438.5	47	24.9	3,282.1	59	31.2	1,563.6	15	7.9	444.9	0	0.0	----	189	84.0	869.8
2004	39	21.8	355.6	52	29.1	3,635.4	45	25.1	1,172.5	12	6.7	349.1	3	1.7	----	179	82.5	764.3
2005	35	18.2	443.6	67	34.9	5,996.1	43	22.4	1,351.0	8	4.2	384.9	4	2.1	----	192	81.7	1,093.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 39 cases reported as unknown race/ethnicity in 2001; 55 in 2002; 28 in 2003; 31 in 2004; and 46 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-6. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	4	0.8	322.5	267	56.7	2,581.7	120	25.5	489.1	4	0.8	635.1	0	0.0	----	471	20.0	1,073.4
2002	9	1.4	739.8	355	53.7	3,447.8	177	26.8	706.4	7	1.1	1,151.6	2	0.3	----	661	23.2	1,476.0
2003	9	1.4	677.5	383	59.0	3,508.4	179	27.6	643.0	1	0.2	150.4	1	0.2	----	649	22.0	1,403.3
2004	9	1.4	657.7	394	59.9	3,623.5	177	26.9	606.8	0	0.0	0.0	1	0.2	----	658	21.4	1,376.5
2005	7	1.0	809.2	460	64.2	3,342.3	184	25.7	665.0	1	0.1	194.7	1	0.1	----	717	22.5	1,522.0
FEMALE																		
2001	26	1.4	2,102.4	886	47.1	8,809.5	523	27.8	2,449.3	12	0.6	1,830.1	3	0.2	----	1,881	80.0	4,343.7
2002	33	1.5	2,601.8	1,012	46.2	9,576.0	701	32.0	3,032.1	12	0.5	1,771.8	7	0.3	----	2,190	76.8	4,940.0
2003	18	0.8	1,290.4	1,219	52.9	10,577.2	763	33.1	2,916.3	7	0.3	972.1	1	0.0	----	2,306	78.0	5,034.9
2004	31	1.3	2,178.5	1,301	53.8	11,153.1	774	32.0	2,818.7	4	0.2	542.2	6	0.2	----	2,417	78.6	5,114.6
2005	32	1.3	3,777.9	1,371	55.6	10,159.8	759	30.8	2,946.0	4	0.2	839.7	7	0.3	----	2,467	77.5	5,343.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2002; 1 in 2003; 1 in 2004; and 2 in 2005.

3. Includes 507 cases reported as unknown race/ethnicity in 2001; and 536 in 2002.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-7. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	5	2.5	88.8	14	6.9	873.7	134	65.7	435.8	2	1.0	57.6	0	0.0	----	204	16.8	372.7
2002	8	3.5	148.3	20	8.7	1,201.0	146	63.8	461.2	0	0.0	0.0	1	0.4	----	229	17.7	412.5
2003	7	3.4	141.3	22	10.7	1,356.8	121	58.7	384.0	1	0.5	29.5	0	0.0	----	206	17.0	363.0
2004	16	8.2	309.2	20	10.3	1,109.4	116	59.8	327.6	2	1.0	55.3	0	0.0	----	194	15.5	333.7
2005	11	4.7	174.6	30	12.9	1,730.4	149	64.2	417.0	3	1.3	84.0	0	0.0	----	232	18.0	406.5
FEMALE																		
2001	58	5.7	1,146.5	62	6.1	5,168.8	576	56.9	2,117.2	22	2.2	744.5	2	0.2	----	1,013	83.2	1,968.7
2002	47	4.4	952.7	78	7.3	5,894.6	606	57.0	2,142.7	28	2.6	935.3	4	0.4	----	1,064	82.3	2,023.7
2003	54	5.4	1,095.0	91	9.0	6,078.9	580	57.7	1,895.0	22	2.2	707.2	4	0.4	----	1,006	83.0	1,862.8
2004	61	5.8	1,237.0	96	9.1	5,867.9	631	59.8	1,916.1	20	1.9	624.8	6	0.6	----	1,055	84.5	1,898.0
2005	65	6.2	1,083.0	116	11.0	7,745.3	668	63.3	1,975.4	25	2.4	752.5	5	0.5	----	1,056	82.0	1,962.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2002; 1 in 2003; and 2 in 2005.

3. Includes 342 cases reported as unknown race/ethnicity in 2001; 355 in 2002; 310 in 2003; 281 in 2004; and 216 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 7-8. Reported Chlamydia Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	5	2.4	63.5	109	53.2	1,780.0	62	30.2	419.9	7	3.4	148.3	2	1.0	----	205	20.1	550.4
2002	12	5.2	145.7	122	53.3	1,875.2	69	30.1	448.8	4	1.7	81.5	1	0.4	----	229	20.4	592.1
2003	5	2.5	62.1	97	48.0	1,478.1	65	32.2	421.5	7	3.5	145.6	1	0.5	----	202	19.1	500.5
2004	6	2.4	71.3	160	63.5	2,296.6	50	19.8	301.5	5	2.0	100.1	2	0.8	----	252	21.2	601.0
2005	13	4.7	159.1	167	60.3	2,423.7	62	22.4	408.4	1	0.4	21.5	1	0.4	----	277	21.4	697.0
FEMALE																		
2001	48	5.9	717.7	363	44.4	6,135.3	241	29.5	1,844.5	30	3.7	747.5	4	0.5	----	817	79.9	2,301.8
2002	54	6.0	775.5	395	44.2	6,318.8	264	29.6	1,955.5	26	2.9	636.6	4	0.4	----	893	79.6	2,403.1
2003	65	7.6	843.7	405	47.5	5,658.7	252	29.5	1,668.2	28	3.3	630.9	2	0.2	----	853	80.9	2,178.4
2004	57	6.1	702.2	475	50.8	6,138.8	275	29.4	1,695.4	29	3.1	629.4	4	0.4	----	935	78.8	2,282.3
2005	70	6.9	881.9	542	53.3	7,608.9	288	28.3	1,941.0	13	1.3	294.0	7	0.7	----	1,017	78.6	2,670.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".


2. Excludes 1 case reported with "unknown/other" gender in 2001; 1 in 2002; and 1 in 2003.

3. Includes 151 cases reported as unknown race/ethnicity in 2001; 171 in 2002; 128 in 2003; 124 in 2004; and 130 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.

Figure 3. Reported Chlamydia Rates for 20-24 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 8. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	154	5.9	311.8	602	23.0	3,393.2	977	37.4	741.6	58	2.2	196.9	7	0.3	----	2,612	23.8	784.3
2002	177	5.8	335.4	694	22.6	3,527.2	1,264	41.2	900.5	107	3.5	330.7	12	0.4	----	3,070	26.2	915.5
2003	258	8.1	461.9	707	22.2	3,339.9	1,331	41.8	917.7	89	2.8	256.7	7	0.2	----	3,183	100.0	928.5
2004	272	7.8	471.4	874	25.1	3,998.7	1,346	38.7	932.2	102	2.9	284.9	18	0.5	----	3,481	100.0	1,000.9
2005	287	8.0	479.2	958	26.8	4,184.2	1,442	40.4	962.4	88	2.5	261.1	18	0.5	----	3,573	100.0	1,044.4
FEMALE																		
2001	469	5.6	1,028.0	1,470	17.5	7,802.1	3,261	38.9	2,847.8	251	3.0	889.2	25	0.3	----	8,383	76.2	2,630.9
2002	557	6.5	1,122.3	1,565	18.1	7,369.4	3,735	43.3	2,927.6	330	3.8	1,032.6	45	0.5	----	8,631	73.8	2,691.1
2003	743	8.0	1,369.9	1,780	19.2	7,605.6	4,238	45.7	3,037.3	396	4.3	1,114.7	11	0.1	----	9,268	74.4	2,827.1
2004	819	8.5	1,492.6	1,893	19.5	8,034.7	4,213	43.5	3,036.2	399	4.1	1,112.3	60	0.6	----	9,684	73.6	2,903.5
2005	887	9.0	1,502.8	2,040	20.8	8,504.2	4,513	46.0	3,188.4	426	4.3	1,284.3	47	0.5	----	9,804	73.3	3,057.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 10 cases reported with "unknown/other" gender in 2001; 8 in 2002; 4 in 2003; 2 in 2004; and 8 in 2005.

3. Includes 3,721 cases reported as unknown race/ethnicity in 2001; 3,215 in 2002; 2,891 in 2003; 3,169 in 2004; and 2,671 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 9. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 1998-2002

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)	249	2.3	1,299.1	295	2.5	1,389.7	367	2.9	1,557.7	424	3.2	1,642.1	454	3.4	1,558.9
Male	55	0.5	557.2	67	0.6	617.0	94	0.8	782.7	106	0.8	808.2	122	0.9	803.5
Female	194	1.8	2,086.7	228	1.9	2,198.9	273	2.2	2,363.4	318	2.4	2,503.1	332	2.5	2,381.6
East (7)³	1,520	13.8	1,524.7	1,682	14.4	1,670.3	1,659	13.3	1,612.6	1,712	13.0	1,644.4	1,738	13.0	1,671.7
Male	333	3.0	655.3	373	3.2	727.7	366	2.9	698.4	387	2.9	729.9	417	3.1	782.3
Female	1,186	10.8	2,426.7	1,306	11.2	2,641.5	1,293	10.4	2,561.8	1,324	10.1	2,591.3	1,318	9.8	2,601.4
Metro (4)³	1,479	13.4	1,675.3	1,647	14.1	1,950.4	1,640	13.2	1,989.5	1,626	12.3	2,007.5	1,633	12.2	2,210.9
Male	370	3.4	791.3	497	4.2	1,108.9	429	3.4	982.5	445	3.4	1,043.0	464	3.5	39,008.0
Female	1,108	10.1	2,668.4	1,150	9.8	2,902.2	1,210	9.7	3,121.1	1,180	9.0	3,078.4	1,169	8.7	34,852.0
San Fernando (2)³	1,467	13.3	1,115.1	1,752	15.0	1,308.4	1,895	15.2	1,365.0	1,927	14.6	1,345.4	1,973	14.7	1,421.9
Male	329	3.0	484.7	435	3.7	628.1	492	4.0	685.4	485	3.7	656.8	517	3.9	715.2
Female	1,138	10.3	1,787.3	1,316	11.2	2,035.5	1,403	11.3	2,092.8	1,442	10.9	2,078.2	1,455	10.9	2,188.9
San Gabriel (3)³	1,274	11.6	1,085.7	1,489	12.7	1,237.9	1,656	13.3	1,332.2	1,620	12.3	1,286.0	1,667	12.5	1,292.1
Male	270	2.5	450.0	371	3.2	604.7	375	3.0	591.8	391	3.0	608.4	388	2.9	66,316.0
Female	1,003	9.1	1,749.4	1,117	9.5	1,895.2	1,281	10.3	2,102.1	1,228	9.3	1,990.3	1,278	9.5	62,696.0
South (6)³	2,491	22.6	2,931.0	2,709	23.1	3,155.0	2,935	23.6	3,374.8	3,057	23.2	3,518.5	2,924	21.8	3,428.1
Male	608	5.5	1,408.3	734	6.3	1,689.1	733	5.9	1,668.6	863	6.6	1,972.0	829	6.2	1,912.0
Female	1,881	17.1	4,498.2	1,975	16.9	4,657.0	2,200	17.7	5,111.6	2,192	16.6	5,083.4	2,095	15.6	4,995.5
South Bay (8)³	1,242	11.3	1,927.9	1,271	10.9	1,943.6	1,420	11.4	2,089.4	1,401	10.6	1,992.8	1,574	11.8	2,239.1
Male	336	3.1	1,043.1	338	2.9	1,027.4	408	3.3	1,188.5	403	3.1	1,135.7	459	3.4	1,268.9
Female	904	8.2	2,806.6	932	8.0	2,868.0	1,012	8.1	3,008.9	996	7.6	2,860.8	1,114	8.3	3,264.6
West (5)	358	3.3	774.7	369	3.2	834.2	444	3.6	1,015.4	454	3.4	1,032.0	460	3.4	1,417.2
Male	99	0.9	444.0	110	0.9	513.1	130	1.0	609.2	125	0.9	578.4	134	1.0	812.4
Female	259	2.4	1,083.0	259	2.2	1,136.2	314	2.5	1,402.6	329	2.5	1,470.0	326	2.4	2,042.1
Unknown	925	8.4	-----	495	4.2	-----	438	3.5	-----	953	7.2	-----	965	7.2	-----
Total	11,005		1,688.8	11,709		1,784.8	12,454		1,857.0	13,174		1,933.6	13,388		2,020.0

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 7 cases reported with "unknown/other" gender in 2001; 6 in 2002; 5 in 2003; and 2 in 2004.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-1. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	6	10.9	206.4	16	29.1	1,567.9	16	29.1	625.9	0	0.0	0.0	0	0.0	----	55	22.1	557.2
2002	9	13.4	235.0	31	46.3	2,166.4	16	23.9	473.0	0	0.0	0.0	0	0.0	----	67	22.7	617.0
2003	15	16.0	402.1	29	30.9	1,961.9	26	27.7	788.0	0	0.0	0.0	0	0.0	----	94	25.6	782.7
2004	19	17.9	430.2	49	46.2	2,608.2	19	17.9	480.2	0	0.0	0.0	0	0.0	----	106	25.0	808.2
2005	13	10.7	234.4	52	42.6	3,015.7	33	27.0	751.5	0	0.0	0.0	1	0.8	----	122	26.9	803.5
FEMALE																		
2001	18	9.3	712.2	52	26.8	6,100.6	48	24.7	2,222.9	2	1.0	1,130.5	0	0.0	----	194	77.9	2,086.7
2002	27	11.8	950.7	58	25.4	5,594.5	55	24.1	2,185.5	6	2.6	2,856.7	0	0.0	----	228	77.3	2,198.9
2003	54	19.8	1,556.3	69	25.3	5,095.8	69	25.3	2,189.2	4	1.5	1,481.8	0	0.0	----	273	74.4	2,363.4
2004	61	19.2	1,605.5	87	27.4	5,349.9	85	26.7	2,362.5	1	0.3	335.3	1	0.3	----	318	75.0	2,503.1
2005	76	22.9	1,457.1	89	26.8	5,696.5	95	28.6	2,447.8	6	1.8	1,386.8	0	0.0	----	332	73.1	2,381.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 91 cases reported as unknown race/ethnicity in 2001; 93 in 2002; 101 in 2002; 102 in 2003; 102 in 2004; and 89 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-2. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	42	12.8	255.1	39	11.9	1,863.4	140	42.6	566.2	11	3.3	231.3	2	0.6	----	329	22.4	484.7
2002	39	9.0	242.8	36	8.3	1,709.6	202	46.4	850.4	16	3.7	332.6	2	0.5	----	435	24.8	628.1
2003	72	14.6	401.9	42	8.5	1,793.0	222	45.1	869.4	15	3.0	272.7	2	0.4	----	492	26.0	685.4
2004	84	17.3	443.8	50	10.3	2,027.9	197	40.6	766.7	13	2.7	219.8	7	1.4	----	485	25.2	656.8
2005	86	16.6	381.1	65	12.6	3,406.3	237	45.8	863.7	25	4.8	393.2	6	1.2	----	517	26.2	715.2
FEMALE																		
2001	118	10.4	742.1	76	6.7	3,770.1	574	50.4	2,604.5	32	2.8	669.8	3	0.3	----	1,138	77.6	1,787.3
2002	173	13.1	1,060.0	88	6.7	4,049.9	627	47.6	2,727.5	55	4.2	1,081.9	11	0.8	----	1,316	75.2	2,035.5
2003	216	15.4	1,233.7	101	7.2	4,288.4	666	47.5	2,720.2	64	4.6	1,156.3	0	0.0	----	1,403	74.0	2,092.8
2004	223	15.5	1,213.7	94	6.5	3,799.0	688	47.7	2,748.7	63	4.4	1,086.8	12	0.8	----	1,442	74.8	2,078.2
2005	233	16.0	1,109.5	120	8.2	6,670.8	753	51.8	3,081.0	51	3.5	858.4	13	0.9	----	1,455	73.8	2,188.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2002; 1 in 2003; and 2 in 2004.

3. Includes 430 cases reported as unknown race/ethnicity in 2001; 502 in 2002; 495 in 2003; 496 in 2004; and 383 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-3. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	15	5.6	166.7	29	10.7	1,714.1	157	58.1	622.8	15	5.6	123.4	1	0.4	----	270	21.2	450.0
2002	28	7.5	301.1	46	12.4	2,459.7	206	55.5	790.5	24	6.5	182.2	2	0.5	----	371	24.9	604.7
2003	25	6.7	280.1	40	10.7	2,073.4	197	52.5	782.8	30	8.0	225.3	1	0.3	----	375	22.6	591.8
2004	31	7.8	329.2	45	11.3	2,088.3	227	57.2	836.3	28	7.1	188.6	1	0.3	----	397	24.4	617.7
2005	38	9.8	335.7	38	9.8	1,720.9	237	61.1	803.3	18	4.6	127.5	5	1.3	----	388	23.3	585.1
FEMALE																		
2001	86	8.6	1,064.6	78	7.8	4,881.0	511	50.9	2,289.8	72	7.2	667.6	5	0.5	----	1,003	78.8	1,749.4
2002	109	9.8	1,241.7	86	7.7	4,753.2	611	54.7	2,483.1	95	8.5	766.6	5	0.4	----	1,117	75.1	1,895.2
2003	123	9.6	1,332.4	93	7.3	4,710.5	762	59.5	2,860.9	114	8.9	874.4	0	0.0	----	1,281	77.4	2,138.9
2004	135	10.9	1,550.5	99	8.0	5,048.1	680	55.2	2,597.5	103	8.4	813.2	5	0.4	----	1,233	75.6	2,055.2
2005	147	11.5	1,328.7	87	6.8	4,008.2	759	59.4	2,692.4	119	9.3	891.0	9	0.7	----	1,278	76.7	2,038.4

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001; 1 in 2002; 1 in 2003; and 2 in 2004.

3. Includes 304 cases reported as unknown race/ethnicity in 2001; 276 in 2002; 271 in 2003; 276 in 2004; and 209 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 10-4. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	26	7.0	494.9	46	12.4	2,823.7	190	51.4	827.3	10	2.7	221.5	1	0.3	----	370	25.0	791.3
2002	30	6.0	590.8	44	8.9	2,683.7	278	55.9	1,216.4	25	5.0	543.3	4	0.8	----	497	30.2	1,108.9
2003	41	9.6	768.9	48	11.2	2,826.1	245	57.2	1,051.0	12	2.8	247.5	0	0.0	----	428	26.1	980.2
2004	37	8.3	665.5	45	10.1	2,756.1	256	57.5	1,151.8	20	4.5	410.9	1	0.2	----	445	27.4	1,043.0
2005	43	9.3	1,308.0	50	10.8	3,262.6	276	59.5	1,197.0	16	3.4	362.7	0	0.0	----	464	28.4	1,189.5
FEMALE																		
2001	71	6.4	1,550.5	82	7.4	5,909.0	544	49.1	3,122.1	42	3.8	976.4	3	0.3	----	1,108	75.0	2,668.4
2002	51	4.4	1,103.8	91	7.9	6,114.5	661	57.5	3,512.6	53	4.6	1,143.2	6	0.5	----	1,150	69.8	2,902.2
2003	73	6.0	1,633.5	107	8.8	7,395.7	653	54.0	3,539.4	69	5.7	1,523.9	3	0.2	----	1,210	73.9	3,121.1
2004	78	6.6	1,585.5	120	10.2	7,928.8	661	56.0	3,450.2	75	6.4	1,565.2	5	0.4	----	1,180	72.6	3,078.4
2005	93	8.0	2,847.2	123	10.5	8,674.5	675	57.7	3,349.3	88	7.5	1,979.3	7	0.6	----	1,169	71.6	3,354.2

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001; 4 in 2004; and 1 in 2005.

3. Includes 463 cases reported as unknown race/ethnicity in 2001; 404 in 2002; 387 in 2003; 327 in 2004; 262 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-5. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	22	22.2	284.1	23	23.2	2,199.6	20	20.2	548.2	6	6.1	170.1	0	0.0	----	99	27.7	444.0
2002	34	30.9	412.1	16	14.5	1,366.5	26	23.6	670.1	10	9.1	279.0	1	0.9	----	110	29.8	513.1
2003	37	28.5	439.0	25	19.2	2,102.1	29	22.3	776.0	10	7.7	300.9	1	0.8	----	130	29.3	609.2
2004	27	21.6	292.7	37	29.6	2,966.6	28	22.4	748.8	7	5.6	212.6	3	2.4	----	125	27.5	578.4
2005	38	28.4	613.5	31	23.1	3,483.1	29	21.6	789.7	3	2.2	172.3	1	0.7	----	134	29.1	812.4
FEMALE																		
2001	42	16.2	544.7	49	18.9	4,390.1	64	24.7	1,808.7	24	9.3	582.5	0	0.0	----	259	72.3	1,083.0
2002	44	17.0	556.7	47	18.1	3,906.6	62	23.9	1,656.4	41	15.8	965.2	1	0.4	----	259	70.2	1,136.2
2003	85	27.1	1,010.6	51	16.2	4,037.6	71	22.6	1,851.5	41	13.1	952.3	3	1.0	----	314	70.7	1,402.6
2004	83	25.2	894.0	63	19.1	4,760.7	93	28.3	2,372.5	35	10.6	792.5	6	1.8	----	329	72.5	1,470.0
2005	77	23.6	1,167.6	64	19.6	6,879.0	82	25.2	2,386.5	42	12.9	1,990.4	3	0.9	----	326	70.9	2,042.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2003.

3. Includes 108 cases reported as unknown race/ethnicity in 2001; 87 in 2002; 91 in 2003; 72 in 2004; and 90 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-6. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	14	2.3	703.5	264	43.4	3,623.9	198	32.6	840.7	2	0.3	177.9	1	0.2	----	608	24.4	1,408.3
2002	7	1.0	351.4	329	44.8	4,345.6	248	33.8	1,025.8	5	0.7	425.0	2	0.3	----	734	27.1	1,689.1
2003	11	1.5	534.9	331	45.2	4,017.1	286	39.0	1,094.2	1	0.1	79.7	1	0.1	----	733	25.0	1,668.6
2004	17	2.0	883.9	429	49.7	5,272.4	283	32.8	1,100.2	1	0.1	81.9	1	0.1	----	863	28.2	1,972.0
2005	15	1.8	1,219.2	429	51.7	4,166.7	266	32.1	1,059.2	3	0.4	442.1	2	0.2	----	829	28.4	1,912.0
FEMALE																		
2001	21	1.1	1,303.6	752	40.0	8,993.9	636	33.8	3,088.3	13	0.7	1,164.7	7	0.4	----	1,881	75.6	4,498.2
2002	24	1.2	1,392.9	793	40.2	9,019.5	740	37.5	3,374.5	10	0.5	788.9	7	0.4	----	1,975	72.9	4,657.0
2003	29	1.3	1,569.4	915	41.6	9,551.3	947	43.0	3,906.2	6	0.3	431.7	1	0.0	----	2,200	75.0	5,111.6
2004	38	1.7	2,121.9	971	44.3	10,213.4	860	39.2	3,524.4	14	0.6	1,033.1	5	0.2	----	2,192	71.8	5,083.4
2005	36	1.7	2,955.4	985	47.0	8,687.7	830	39.6	3,456.0	15	0.7	1,918.8	4	0.2	----	2,095	71.6	4,995.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2001; 1 in 2003; 2 in 2004; and 1 in 2005.

3. Includes 581 cases reported as unknown race/ethnicity in 2001; 544 in 2002; 405 in 2003; 436 in 2004; and 339 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-7. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	14	4.2	369.0	19	5.7	2,449.8	157	47.1	677.8	7	2.1	312.9	0	0.0	----	333	21.9	655.3
2002	16	4.3	406.4	27	7.2	2,957.0	180	48.3	736.4	9	2.4	361.9	0	0.0	----	373	22.2	727.7
2003	25	6.8	656.1	12	3.3	1,186.4	177	48.4	727.8	8	2.2	304.3	1	0.3	----	366	22.1	698.4
2004	17	4.4	399.9	33	8.5	2,626.0	200	51.7	719.0	15	3.9	477.6	3	0.8	----	387	22.6	729.9
2005	23	5.5	415.4	41	9.8	3,856.8	239	57.3	794.3	11	2.6	313.1	2	0.5	----	417	24.0	782.3
FEMALE																		
2001	42	3.5	1,178.5	46	3.9	5,977.0	562	47.4	2,650.3	22	1.9	1,043.7	2	0.2	----	1,186	78.1	2,426.7
2002	68	5.2	1,799.7	52	4.0	5,746.5	651	49.8	2,813.2	26	2.0	1,090.8	6	0.5	----	1,306	77.8	2,641.5
2003	64	4.9	1,614.9	53	4.1	5,083.1	692	53.5	2,745.2	36	2.8	1,332.1	2	0.2	----	1,293	77.9	2,562.4
2004	86	6.5	2,090.1	54	4.1	4,527.4	750	56.6	2,750.5	29	2.2	973.5	8	0.6	----	1,324	77.4	2,591.3
2005	82	6.2	1,416.5	83	6.3	7,704.7	860	65.3	2,850.5	29	2.2	837.5	4	0.3	----	1,318	76.0	2,601.4

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001; 3 in 2002; and 1 in 2004.

3. Includes 648 cases reported as unknown race/ethnicity in 2001; 644 in 2002; 589 in 2003; 516 in 2004; and 361 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 10-8. Reported Chlamydia Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	13	3.9	255.1	161	47.9	3,980.3	87	25.9	662.3	7	2.1	200.5	2	0.6	----	336	27.1	1,043.1
2002	10	3.0	186.3	157	46.4	3,607.5	97	28.7	706.4	16	4.7	421.1	1	0.3	----	338	26.6	1,027.4
2003	28	7.2	475.2	163	41.9	3,417.4	130	33.4	906.0	10	2.6	238.7	1	0.3	----	389	27.1	1,133.1
2004	29	7.2	499.6	166	41.3	3,584.7	108	26.9	813.1	12	3.0	294.4	1	0.2	----	402	28.8	1,132.8
2005	24	5.2	365.3	232	50.5	4,773.6	108	23.5	777.9	12	2.6	278.9	1	0.2	----	459	29.1	1,268.9
FEMALE																		
2001	55	6.1	1,185.3	310	34.3	6,327.8	284	31.4	2,439.3	36	4.0	1,121.2	2	0.2	----	904	72.9	2,806.6
2002	49	5.3	1,078.7	321	34.4	6,525.4	285	30.6	2,428.1	43	4.6	1,301.5	9	1.0	----	932	73.4	2,868.0
2003	66	6.3	1,333.6	366	35.0	6,966.8	336	32.1	2,690.2	52	5.0	1,434.7	2	0.2	----	1,047	72.9	3,113.0
2004	76	7.6	1,356.3	368	37.0	6,446.8	320	32.2	2,370.7	58	5.8	1,423.6	7	0.7	----	995	71.2	2,857.9
2005	88	7.9	1,400.0	426	38.2	8,070.3	362	32.4	2,758.5	65	5.8	1,558.3	7	0.6	----	1,116	70.9	3,270.4

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2001; 1 in 2002; 3 in 2004; and 1 in 2005.

3. Includes 283 cases reported as unknown race/ethnicity in 2001; 282 in 2002; 282 in 2003; 262 in 2004; and 250 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.





Figure 4. 2005 Reported Chlamydia Rates Per 100,000 10-24 Year Olds, by Service Planning Area (SPA).

Figure 5. Reported Gonorrhea Rates for 10-14 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 11. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	5	71.4	18.1	0	0.0	0.0	0	0.0	0.0	0	0.0	----	7	8.0	1.9
2002	0	0.0	0.0	4	44.4	13.2	3	33.3	1.7	0	0.0	0.0	0	0.0	----	9	10.7	2.3
2003	1	5.6	1.6	9	50.0	29.6	3	16.7	1.6	1	5.6	3.1	0	0.0	----	18	21.4	4.5
2004	0	0.0	0.0	13	81.3	36.3	2	12.5	0.9	0	0.0	0.0	0	0.0	----	16	15.1	4.0
2005	0	0.0	0.0	13	56.5	41.7	6	26.1	3.1	0	0.0	0.0	0	0.0	----	23	21.1	5.9
FEMALE																		
2001	0	0.0	0.0	49	61.3	162.8	13	16.3	8.2	1	1.3	3.4	0	0.0	----	80	92.0	22.4
2002	2	2.7	2.9	49	65.3	148.4	12	16.0	6.6	0	0.0	0.0	1	1.3	----	75	89.3	20.1
2003	2	3.0	3.0	35	53.0	107.9	17	25.8	9.1	1	1.5	3.1	0	0.0	----	66	78.6	17.2
2004	9	10.0	13.9	57	63.3	176.4	9	10.0	4.6	1	1.1	3.0	0	0.0	----	90	84.9	23.3
2005	7	8.1	10.2	55	64.0	169.8	13	15.1	6.6	0	0.0	0.0	1	1.2	----	86	78.9	23.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 19 cases reported as unknown race/ethnicity in 2001; 13 in 2002; 15 in 2003; 15 in 2004; and 14 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 12. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)	3	3.4	9.2	3	3.6	9.0	12	14.3	35.4	11	10.4	32.3	9	8.3	27.4
Male	0	0.0	0.0	0	0.0	0.0	3	3.6	17.6	2	1.9	11.7	3	2.8	18.1
Female	3	3.4	18.6	3	3.6	18.1	9	10.7	53.4	9	8.5	53.1	6	5.5	36.9
East (7)	4	4.6	3.4	3	3.6	2.5	4	4.8	3.2	5	4.7	3.9	7	6.4	5.7
Male	0	0.0	0.0	1	1.2	1.6	0	0.0	0.0	0	0.0	0.0	2	1.8	3.2
Female	4	4.6	7.0	2	2.4	3.4	4	4.8	6.5	5	4.7	8.1	5	4.6	8.3
Metro (4)	4	4.6	5.1	2	2.4	2.4	4	4.8	4.5	7	6.6	7.8	6	5.5	6.9
Male	1	1.1	2.5	0	0.0	0.0	1	1.2	2.2	2	1.9	4.3	1	0.9	2.2
Female	3	3.4	7.8	2	2.4	4.9	3	3.6	7.0	5	4.7	11.5	5	4.6	11.7
San Fernando (2)	6	6.9	3.9	4	4.8	2.5	7	8.3	4.2	3	2.8	1.8	14	12.8	8.6
Male	1	1.1	1.3	1	1.2	1.2	1	1.2	1.2	0	0.0	0.0	2	1.8	2.4
Female	5	5.7	6.6	3	3.6	3.8	6	7.1	7.4	3	2.8	3.7	12	11.0	15.1
San Gabriel (3)	2	2.3	1.5	7	8.3	5.1	4	4.8	2.9	7	6.6	5.1	8	7.3	5.9
Male	0	0.0	0.0	0	0.0	0.0	1	1.2	1.4	1	0.9	1.4	0	0.0	0.0
Female	2	2.3	3.1	7	8.3	10.5	3	3.6	4.4	6	5.7	9.0	8	7.3	12.2
South (6)	46	52.9	47.0	40	47.6	39.3	35	41.7	33.4	52	49.1	49.3	47	43.1	45.0
Male	4	4.6	8.0	6	7.1	11.6	8	9.5	15.0	9	8.5	16.7	9	8.3	16.8
Female	42	48.3	87.4	34	40.5	68.2	27	32.1	52.5	43	40.6	83.2	38	34.9	74.4
South Bay (8)	11	12.6	13.1	23	27.4	26.2	7	8.3	7.7	9	8.5	9.9	8	7.3	9.1
Male	1	1.1	2.3	1	1.2	2.2	2	2.4	4.3	0	0.0	0.0	4	3.7	8.9
Female	10	11.5	24.4	22	26.2	51.1	5	6.0	11.2	9	8.5	20.0	4	3.7	9.2
West (5)	2	2.3	6.4	0	0.0	0.0	2	2.4	5.7	5	4.7	14.0	2	1.8	6.4
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	2	1.9	11.1	0	0.0	0.0
Female	2	2.3	12.9	0	0.0	0.0	2	2.4	11.5	3	2.8	17.1	2	1.8	13.1
Unknown	9	10.3	----	2	2.4	----	9	10.7	----	7	6.6	----	8	7.3	----
Total	87	11.9	84	11.0	84	10.7	106	13.4	109	14.2					

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 13-1. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	1	33.3	22.5	1	0.0	45.7	0	0.0	0.0	0	0.0	0.0	0	0.0	----	3	25.0	17.6
2004	0	0.0	0.0	2	0.0	57.9	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	18.2	11.7
2005	0	0.0	0.0	2	0.0	75.5	1	0.0	15.7	0	0.0	0.0	0	0.0	----	3	33.3	18.1
FEMALE																		
2001	0	0.0	0.0	2	66.7	70.7	1	33.3	17.3	0	0.0	0.0	0	0.0	----	3	100.0	18.6
2002	0	0.0	0.0	2	66.7	99.1	0	0.0	0.0	0	0.0	0.0	0	0.0	----	3	100.0	18.1
2003	1	11.1	22.8	4	44.4	185.6	1	11.1	22.9	0	0.0	0.0	0	0.0	----	9	75.0	53.4
2004	1	11.1	24.4	4	44.4	175.0	1	11.1	21.8	0	0.0	0.0	0	0.0	----	9	81.8	53.1
2005	2	33.3	34.7	3	50.0	138.7	0	0.0	0.0	0	0.0	0.0	0	0.0	----	6	66.7	36.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2002; 4 in 2003; 3 in 2004; and 1 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 13-2. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	1	100.0	30.6	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	16.7	1.3
2002	0	0.0	0.0	0	0.0	0.0	1	100.0	2.5	0	0.0	0.0	0	0.0	----	1	25.0	1.2
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	2.4	0	0.0	0.0	0	0.0	----	1	14.3	1.2
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	1	50.0	4.9	0	0.0	0.0	0	0.0	----	2	14.3	2.4
FEMALE																		
2001	0	0.0	0.0	2	40.0	80.6	1	20.0	3.5	1	20.0	18.0	0	0.0	----	5	83.3	6.6
2002	0	0.0	0.0	1	33.3	31.1	2	66.7	5.3	0	0.0	0.0	0	0.0	----	3	75.0	3.8
2003	1	16.7	4.0	1	16.7	36.4	3	50.0	8.9	0	0.0	0.0	0	0.0	----	6	85.7	7.4
2004	2	66.7	6.8	1	33.3	29.9	0	0.0	0.0	0	0.0	0.0	0	0.0	----	3	100.0	3.7
2005	2	16.7	8.0	6	50.0	243.7	2	16.7	6.1	0	0.0	0.0	0	0.0	----	12	85.7	15.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2000; 1 in 2003; and 2 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 13-3. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	25.0	1.4
2004	0	0.0	0.0	0	0.0	0.0	1	0.0	2.4	0	0.0	0.0	0	0.0	----	1	14.3	1.4
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	1	50.0	33.3	1	50.0	2.8	0	0.0	0.0	0	0.0	----	2	100.0	3.1
2002	0	0.0	0.0	5	71.4	169.0	2	28.6	5.3	0	0.0	0.0	0	0.0	----	7	100.0	10.5
2003	0	0.0	0.0	0	0.0	0.0	2	66.7	7.7	0	0.0	0.0	0	0.0	----	3	75.0	4.4
2004	2	33.3	24.0	1	16.7	43.9	2	33.3	6.1	0	0.0	0.0	0	0.0	----	6	85.7	9.0
2005	2	25.0	20.6	4	50.0	155.8	1	12.5	3.0	0	0.0	0.0	0	0.0	----	8	100.0	12.2

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 2 case reported as unknown race/ethnicity in 2003; 1 in 2004; and 1 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 13-4. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	25.0	2.5
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	440.6	0	0.0	0.0	0	0.0	----	1	25.0	2.2
2004	0	0.0	0.0	2	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	28.6	4.3
2005	0	0.0	0.0	0	0.0	0.0	1	100.0	410.1	0	0.0	0.0	0	0.0	----	1	16.7	2.2
FEMALE																		
2001	0	0.0	0.0	1	33.3	53.8	2	66.7	7.4	0	0.0	0.0	0	0.0	----	3	75.0	7.8
2002	1	50.0	19.4	1	50.0	52.2	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	100.0	4.9
2003	0	0.0	0.0	3	100.0	152.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	3	75.0	7.0
2004	2	40.0	38.0	2	40.0	100.6	1	20.0	3.2	0	0.0	0.0	0	0.0	----	5	71.4	11.5
2005	0	0.0	0.0	4	80.0	206.5	1	20.0	3.1	0	0.0	0.0	0	0.0	----	5	83.3	11.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2001.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 13-5. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	2	0.0	125.7	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	40.0	11.1
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	1	50.0	72.0	1	50.0	27.4	0	0.0	0.0	0	0.0	----	2	100.0	12.9
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	2	100.0	134.4	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	100.0	11.5
2004	1	33.3	10.1	2	66.7	132.3	0	0.0	0.0	0	0.0	0.0	0	0.0	----	3	60.0	17.1
2005	0	0.0	0.0	1	50.0	76.3	0	0.0	0.0	0	0.0	0.0	1	50.0	----	2	100.0	13.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 13-6. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	3	75.0	26.3	0	0.0	0.0	0	0.0	0.0	0	0.0	----	4	8.7	8.0
2002	0	0.0	0.0	2	33.3	20.0	2	33.3	8.4	0	0.0	0.0	0	0.0	----	6	15.0	11.6
2003	0	0.0	0.0	7	87.5	54.9	0	0.0	0.0	0	0.0	0.0	0	0.0	----	8	22.9	15.0
2004	0	0.0	0.0	7	77.8	57.3	1	11.1	2.9	0	0.0	0.0	0	0.0	----	9	17.3	16.7
2005	0	0.0	0.0	7	77.8	45.9	2	22.2	5.4	0	0.0	0.0	0	0.0	----	9	19.1	16.8
FEMALE																		
2001	0	0.0	0.0	33	78.6	246.2	4	9.5	14.3	0	0.0	0.0	0	0.0	----	42	91.3	87.4
2002	0	0.0	0.0	27	79.4	210.9	2	5.9	6.9	0	0.0	0.0	0	0.0	----	34	85.0	68.2
2003	0	0.0	0.0	18	66.7	137.9	6	22.2	19.0	0	0.0	0.0	0	0.0	----	27	77.1	52.5
2004	0	0.0	0.0	36	83.7	309.6	0	0.0	0.0	0	0.0	0.0	0	0.0	----	43	82.7	83.2
2005	1	2.6	192.5	31	81.6	226.6	3	7.9	9.2	0	0.0	0.0	0	0.0	----	38	80.9	74.4

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 6 cases reported as unknown race/ethnicity in 2001; 7 in 2002; 4 in 2003; 8 in 2004; and 3 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 13-7. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	1	100.0	45.4	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	33.3	1.6
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	1	50.0	52.7	1	50.0	2.0	0	0.0	0.0	0	0.0	----	2	28.6	3.2
FEMALE																		
2001	0	0.0	0.0	1	25.0	47.0	3	75.0	6.8	0	0.0	0.0	0	0.0	----	4	100.0	7.0
2002	0	0.0	0.0	0	0.0	0.0	2	100.0	4.3	0	0.0	0.0	0	0.0	----	2	66.7	3.4
2003	0	0.0	0.0	1	25.0	44.8	3	75.0	6.2	0	0.0	0.0	0	0.0	----	4	100.0	6.5
2004	0	0.0	0.0	2	40.0	110.5	2	40.0	5.0	0	0.0	0.0	0	0.0	----	5	100.0	8.1
2005	0	0.0	0.0	0	0.0	0.0	4	80.0	10.4	0	0.0	0.0	0	0.0	----	5	71.4	8.3

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2004; and 1 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 13-8. Reported Gonorrhea Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	1	100.0	11.6	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	9.1	2.3
2002	0	0.0	0.0	1	100.0	11.4	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	4.3	2.2
2003	0	0.0	0.0	0	0.0	0.0	1	50.0	4.6	1	50.0	18.3	0	0.0	----	2	28.6	4.3
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	3	75.0	48.5	0	0.0	0.0	0	0.0	0.0	0	0.0	----	4	50.0	8.9
FEMALE																		
2001	0	0.0	0.0	7	70.0	115.1	0	0.0	0.0	0	0.0	0.0	0	0.0	----	10	90.9	24.4
2002	1	4.5	12.1	13	59.1	168.1	4	18.2	24.2	0	0.0	0.0	1	4.5	----	22	95.7	51.1
2003	0	0.0	0.0	3	60.0	40.9	1	20.0	6.1	0	0.0	0.0	0	0.0	----	5	71.4	11.2
2004	0	0.0	0.0	8	88.9	99.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	9	100.0	20.0
2005	0	0.0	0.0	4	100.0	48.6	0	0.0	0.0	0	0.0	0.0	0	0.0	----	4	50.0	9.2


1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 3 cases reported as unknown race/ethnicity in 2001; 3 in 2002; 1 in 2003; 1 in 2004; and 1 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.

Figure 6. Reported Gonorrhea Rates for 15-19 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 14. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	24	5.7	39.4	205	48.5	821.1	85	20.1	60.1	4	0.9	12.3	0	0.0	----	423	26.7	122.1
2002	22	4.7	33.5	236	50.0	876.3	105	22.2	68.9	8	1.7	22.8	2	0.4	----	472	29.2	132.7
2003	26	4.9	37.4	287	53.8	1,003.9	116	21.8	70.5	7	1.3	18.8	1	0.2	----	533	100.0	145.2
2004	31	4.7	43.3	375	56.6	1,269.7	140	21.1	80.6	2	0.3	5.2	2	0.3	----	663	100.0	174.9
2005	36	4.7	52.2	455	59.2	1,479.9	140	18.2	85.8	6	0.8	17.9	3	0.4	----	769	100.0	215.1
FEMALE																		
2001	59	5.1	99.2	633	54.4	2,487.7	191	16.4	140.6	11	0.9	34.8	3	0.3	----	1,164	73.3	354.1
2002	62	5.4	95.1	602	52.6	2,153.2	247	21.6	166.0	14	1.2	40.9	7	0.6	----	1,144	70.8	336.0
2003	80	6.8	115.3	657	56.3	2,181.2	232	19.9	143.9	13	1.1	35.8	1	0.1	----	1,168	68.7	330.0
2004	82	6.0	114.2	777	56.7	2,469.9	290	21.2	170.0	15	1.1	39.9	1	0.1	----	1,371	67.4	373.0
2005	99	6.3	143.9	882	56.0	2,853.1	339	21.5	211.3	17	1.1	52.6	9	0.6	----	1,575	67.2	458.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2001; 4 in 2002; 1 in 2003; 1 in 2004 and 2 in 2005.

3. Includes 372 cases reported as unknown race/ethnicity in 2001; 311 in 2002; 281 in 2003; 319 in 2004; and 358 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 15. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 1998-2002

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)	86	5.4	299.3	91	5.6	303.9	127	7.5	412.0	148	7.3	466.6	139	5.9	413.9
Male	18	1.1	120.4	28	1.7	181.0	41	2.4	259.4	44	2.2	272.8	45	1.9	258.8
Female	68	4.3	493.4	63	3.9	435.3	86	5.1	572.6	104	5.1	667.1	94	4.0	580.5
East (7)³	112	7.0	105.5	122	7.5	112.9	113	6.6	102.0	109	5.4	95.8	147	6.3	132.5
Male	32	2.0	58.5	39	2.4	70.3	33	1.9	58.1	22	1.1	37.8	45	1.9	78.8
Female	80	5.0	155.5	83	5.1	157.9	80	4.7	148.1	87	4.3	156.5	101	4.3	187.7
Metro (4)³	112	7.0	151.3	137	8.5	178.0	145	8.5	178.5	153	7.5	177.2	156	6.6	215.1
Male	28	1.8	72.6	45	2.8	112.8	62	3.6	147.8	59	2.9	132.5	70	3.0	188.4
Female	84	5.3	236.9	91	5.6	245.5	82	4.8	208.6	94	4.6	224.8	85	3.6	240.3
San Fernando (2)³	92	5.8	64.9	122	7.5	82.6	108	6.3	70.4	142	7.0	89.0	166	7.1	112.5
Male	25	1.6	34.2	40	2.5	52.9	32	1.9	40.9	48	2.4	59.2	52	2.2	68.8
Female	66	4.2	96.1	82	5.1	113.8	76	4.5	101.0	94	4.6	119.9	114	4.9	158.4
San Gabriel (3)	104	6.5	81.2	132	8.1	102.1	137	8.0	104.7	154	7.6	116.7	169	7.2	129.0
Male	30	1.9	45.4	36	2.2	54.1	44	2.6	65.6	60	2.9	89.0	43	1.8	64.5
Female	74	4.7	119.2	96	5.9	152.9	93	5.5	145.9	94	4.6	145.7	126	5.4	195.7
South (6)³	665	41.9	762.8	663	40.9	744.0	757	44.5	822.4	826	40.6	868.9	996	42.5	1067.8
Male	170	10.7	387.4	187	11.5	417.6	225	13.2	486.5	264	13.0	552.3	331	14.1	702.6
Female	494	31.1	1140.8	476	29.4	1073.7	532	31.3	1161.6	562	27.6	1189.2	665	28.3	1440.4
South Bay (8)	271	17.1	372.6	235	14.5	309.9	208	12.2	261.6	283	13.9	341.4	347	14.8	445.9
Male	78	4.9	209.4	65	4.0	168.1	59	3.5	146.2	96	4.7	229.0	120	5.1	302.0
Female	193	12.1	543.8	170	10.5	457.5	149	8.8	380.5	187	9.2	456.5	227	9.7	596.2
West (5)	40	2.5	109.8	36	2.2	91.6	34	2.0	80.1	46	2.3	101.0	37	1.6	107.6
Male	15	0.9	83.3	8	0.5	41.4	7	0.4	33.8	13	0.6	58.8	11	0.5	65.3
Female	25	1.6	135.7	28	1.7	140.0	27	1.6	124.3	33	1.6	140.9	26	1.1	148.1
Unknown	107	6.7	-----	82	5.1	-----	73	4.3	-----	174	8.6	-----	189	8.1	-----
Total	1,589		235.3	1,620		232.7	1,702		236.0	2,035		272.5	2,346		334.6

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 2 cases reported with "unknown/other" gender in 2001; 1 in 2002; 1 in 2003; and 2 in 2005.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-1. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	5.6	24.6	8	44.4	548.9	2	11.1	61.8	0	0.0	0.0	0	0.0	----	18	20.9	120.4
2002	4	14.3	65.0	21	75.0	858.3	1	3.6	19.2	0	0.0	0.0	0	0.0	----	28	30.8	181.0
2003	3	7.3	61.1	27	65.9	1,248.6	1	2.4	22.5	0	0.0	0.0	0	0.0	----	41	32.3	259.4
2004	2	4.5	35.7	26	59.1	955.2	11	25.0	200.6	0	0.0	0.0	0	0.0	----	44	29.7	272.8
2005	1	2.2	16.1	32	71.1	1,386.0	3	6.7	61.1	0	0.0	0.0	0	0.0	----	45	32.4	258.8
FEMALE																		
2001	10	14.7	223.6	33	48.5	2,074.1	5	7.4	150.8	0	0.0	0.0	0	0.0	----	68	79.1	493.4
2002	8	12.7	155.3	35	55.6	1,715.0	7	11.1	169.6	1	1.6	305.8	0	0.0	----	63	69.2	435.3
2003	16	18.6	321.5	44	51.2	2,017.7	8	9.3	185.1	0	0.0	0.0	0	0.0	----	86	67.7	572.6
2004	21	20.2	409.0	52	50.0	2,106.0	13	12.5	266.2	0	0.0	0.0	0	0.0	----	104	70.3	667.1
2005	12	12.8	175.5	63	67.0	2,595.1	11	11.7	213.0	1	1.1	220.1	0	0.0	----	94	67.6	580.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 27 cases reported as unknown race/ethnicity in 2001; 14 in 2002; 28 in 2003; 23 in 2004; and 16 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-2. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	4	16.0	21.5	5	20.0	249.1	7	28.0	33.2	0	0.0	0.0	0	0.0	----	25	27.5	34.2
2002	5	12.5	19.1	13	32.5	455.0	15	37.5	50.5	1	2.5	14.3	1	2.5	----	40	32.8	52.9
2003	8	25.0	34.8	6	18.8	234.2	9	28.1	33.4	1	3.1	16.5	0	0.0	----	32	29.6	40.9
2004	7	14.6	26.4	13	27.1	423.7	20	41.7	61.8	1	2.1	14.1	0	0.0	----	48	33.8	59.2
2005	7	13.5	27.1	16	30.8	692.9	18	34.6	64.7	1	1.9	16.2	1	1.9	----	52	31.3	68.8
FEMALE																		
2001	15	22.7	67.0	13	19.7	519.8	24	36.4	95.0	2	3.0	33.9	0	0.0	----	66	72.5	96.1
2002	11	13.4	47.9	22	26.8	840.4	28	34.1	107.3	4	4.9	66.5	1	1.2	----	82	67.2	113.8
2003	18	23.7	71.6	17	22.4	577.3	28	36.8	94.9	2	2.6	30.4	0	0.0	----	76	70.4	101.0
2004	23	24.5	85.8	15	16.0	462.6	42	44.7	128.7	3	3.2	42.3	1	1.1	----	94	66.2	119.9
2005	16	14.0	60.6	31	27.2	1,292.6	48	42.1	167.6	6	5.3	97.4	0	0.0	----	114	68.7	158.4

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001.

3. Includes 21 cases reported as unknown race/ethnicity in 2001; 21 in 2002; 19 in 2003; 17 in 2004; and 22 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-3. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	4	13.3	42.5	9	30.0	384.6	9	30.0	36.7	0	0.0	0.0	0	0.0	----	30	28.8	45.4
2002	3	8.3	30.1	9	25.0	359.0	16	44.4	58.2	1	2.8	7.4	0	0.0	----	36	27.3	54.1
2003	4	9.1	45.1	6	13.6	271.9	21	47.7	79.9	2	4.5	15.6	0	0.0	----	44	32.1	65.6
2004	5	8.3	52.5	16	26.7	679.6	29	48.3	95.2	0	0.0	0.0	1	1.7	----	60	39.0	89.0
2005	4	9.3	36.3	11	25.6	432.6	21	48.8	68.9	1	2.3	7.6	0	0.0	----	43	25.4	64.5
FEMALE																		
2001	11	14.9	107.5	25	33.8	1,072.3	24	32.4	88.2	2	2.7	15.6	1	1.4	----	74	71.2	119.2
2002	11	11.5	117.5	22	22.9	1,018.1	43	44.8	163.6	0	0.0	0.0	1	1.0	----	96	72.7	152.9
2003	7	7.5	73.5	19	20.4	859.7	51	54.8	179.0	1	1.1	7.6	0	0.0	----	93	67.9	145.9
2004	10	10.6	111.2	20	21.3	949.7	45	47.9	157.2	2	2.1	15.4	0	0.0	----	94	61.0	145.7
2005	15	11.9	135.9	28	22.2	1,166.6	60	47.6	202.9	4	3.2	33.5	1	0.8	----	126	74.6	195.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 19 cases reported as unknown race/ethnicity in 2001; 26 in 2002; 26 in 2003; 26 in 2004; and 24 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 16-4. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	7	25.0	181.4	6	21.4	421.9	8	28.6	40.8	0	0.0	0.0	0	0.0	----	28	25.0	72.6
2002	4	8.9	82.4	8	17.8	505.1	23	51.1	106.5	1	2.2	22.1	1	2.2	----	45	33.1	112.8
2003	1	1.6	17.3	16	25.8	931.2	34	54.8	145.7	2	3.2	40.8	0	0.0	----	62	43.1	147.8
2004	5	8.5	80.5	18	30.5	1,039.7	24	40.7	103.3	1	1.7	20.3	0	0.0	----	59	38.6	132.5
2005	6	8.6	191.6	20	28.6	1,263.3	28	40.0	133.3	2	2.9	51.5	0	0.0	----	70	45.2	188.4
FEMALE																		
2001	8	9.5	205.9	37	44.0	2,600.3	21	25.0	114.2	0	0.0	0.0	0	0.0	----	84	75.0	236.9
2002	5	5.5	106.1	36	39.6	2,279.7	33	36.3	164.8	1	1.1	22.3	1	1.1	----	91	66.9	245.5
2003	6	7.3	110.9	38	46.3	2,276.2	24	29.3	113.2	1	1.2	21.3	0	0.0	----	82	56.9	208.6
2004	4	4.3	63.1	44	46.8	2,413.1	32	34.0	136.3	3	3.2	58.2	0	0.0	----	94	61.4	224.8
2005	7	8.2	218.9	33	38.8	2,330.8	29	34.1	137.6	2	2.4	53.3	0	0.0	----	85	54.8	240.3

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2002; 1 in 2003; and 1 in 2005.

3. Includes 25 cases reported as unknown race/ethnicity in 2001; 23 in 2002; 22 in 2004; 22 in 2004; and 28 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-5. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	6.7	11.9	10	66.7	844.1	2	13.3	55.7	0	0.0	0.0	0	0.0	----	15	37.5	83.3
2002	1	12.5	12.5	3	37.5	280.1	2	25.0	63.0	0	0.0	0.0	0	0.0	----	8	22.2	41.4
2003	3	42.9	30.1	2	28.6	156.2	1	14.3	26.6	0	0.0	0.0	0	0.0	----	7	20.6	33.8
2004	3	23.1	26.1	8	61.5	546.1	1	7.7	23.5	0	0.0	0.0	0	0.0	----	13	28.3	58.8
2005	3	27.3	36.1	4	36.4	340.1	1	9.1	28.1	0	0.0	0.0	2	18.2	----	11	29.7	65.3
FEMALE																		
2001	4	16.0	53.3	12	48.0	1,141.7	3	12.0	103.6	0	0.0	0.0	0	0.0	----	25	62.5	135.7
2002	6	21.4	66.5	12	42.9	976.5	2	7.1	61.1	3	10.7	104.0	0	0.0	----	28	77.8	140.0
2003	11	40.7	94.7	9	33.3	580.1	5	18.5	122.3	1	3.7	27.4	0	0.0	----	27	79.4	124.3
2004	3	9.1	28.1	18	54.5	1,293.4	5	15.2	133.9	1	3.0	29.9	0	0.0	----	33	71.7	140.9
2005	6	23.1	76.6	8	30.8	721.4	6	23.1	189.9	1	3.8	48.5	0	0.0	----	26	70.3	148.1

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 8 cases reported as unknown race/ethnicity in 2001; 7 in 2002; 2 in 2003; 7 in 2004; and 6 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-6. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	2	1.2	170.3	111	65.3	1,133.5	21	12.4	90.4	1	0.6	167.7	0	0.0	----	170	25.6	387.4
2002	2	1.1	173.9	124	66.3	1,274.0	17	9.1	71.8	4	2.1	696.1	0	0.0	----	187	28.2	417.6
2003	1	0.4	73.6	172	76.4	1,541.5	29	12.9	101.9	1	0.4	147.2	0	0.0	----	225	29.7	486.5
2004	0	0.0	0.0	197	74.6	1,868.3	29	11.0	102.5	0	0.0	0.0	0	0.0	----	264	32.0	552.3
2005	7	2.1	815.7	261	78.9	1,911.7	30	9.1	109.3	1	0.3	196.3	0	0.0	----	331	33.2	702.6
FEMALE																		
2001	1	0.2	77.3	344	69.6	3,271.1	51	10.3	228.4	2	0.4	291.7	0	0.0	----	494	74.4	1,140.8
2002	3	0.6	226.5	336	70.6	3,044.1	57	12.0	236.1	1	0.2	141.4	4	0.8	----	476	71.8	1,073.7
2003	4	0.8	286.3	388	72.9	3,361.0	70	13.2	267.1	2	0.4	277.3	0	0.0	----	532	70.3	1,161.6
2004	3	0.5	202.9	433	77.0	3,572.8	75	13.3	262.9	0	0.0	0.0	0	0.0	----	562	68.0	1,189.2
2005	17	2.6	1,949.8	498	74.9	3,585.1	85	12.8	320.5	0	0.0	0.0	3	0.5	----	665	66.8	1,440.4

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001.

3. Includes 131 cases reported as unknown race/ethnicity in 2001; 115 in 2002; 90 in 2003; 89 in 2004; and 94 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-7. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	2	6.3	34.5	4	12.5	242.8	19	59.4	60.1	0	0.0	0.0	0	0.0	----	32	28.6	58.5
2002	0	0.0	0.0	7	17.9	417.0	22	56.4	68.9	1	2.6	28.2	0	0.0	----	39	32.0	70.3
2003	3	9.1	56.3	10	30.3	573.8	13	39.4	38.4	0	0.0	0.0	0	0.0	----	33	29.2	58.1
2004	1	4.5	19.9	4	18.2	227.9	12	54.5	34.8	0	0.0	0.0	0	0.0	----	22	20.2	37.8
2005	3	6.7	46.9	9	20.0	511.4	26	57.8	71.7	0	0.0	0.0	0	0.0	----	45	30.8	78.8
FEMALE																		
2001	5	6.3	88.0	21	26.3	1,559.1	36	45.0	117.8	1	1.3	30.1	1	1.3	----	80	71.4	155.5
2002	10	12.0	167.3	15	18.1	935.7	45	54.2	131.3	2	2.4	55.1	0	0.0	----	83	68.0	157.9
2003	7	8.8	146.0	18	22.5	1,236.4	30	37.5	100.8	3	3.8	99.2	0	0.0	----	80	70.8	148.1
2004	3	3.4	59.1	23	26.4	1,365.3	42	48.3	123.9	1	1.1	30.3	0	0.0	----	87	79.8	156.5
2005	2	2.0	33.8	26	25.7	1,760.5	53	52.5	158.9	1	1.0	30.5	1	1.0	----	101	69.2	187.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2005.

3. Includes 23 cases reported as unknown race/ethnicity in 2001; 20 in 2002; 29 in 2003; 23 in 2004; and 25 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 16-8. Reported Gonorrhea Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	2	2.6	26.3	48	61.5	810.5	15	19.2	105.0	3	3.8	65.7	0	0.0	----	78	28.8	209.4
2002	3	4.6	38.4	46	70.8	745.1	7	10.8	48.0	0	0.0	0.0	0	0.0	----	65	27.7	168.1
2003	3	5.1	38.0	43	72.9	669.3	3	5.1	19.9	1	1.7	21.2	0	0.0	----	59	28.4	146.2
2004	7	7.3	76.7	78	81.3	1,032.8	7	7.3	38.9	0	0.0	0.0	0	0.0	----	96	33.9	229.0
2005	3	2.5	37.7	89	74.2	1,324.9	11	9.2	74.3	0	0.0	0.0	0	0.0	----	120	34.6	302.0
FEMALE																		
2001	4	2.1	58.7	135	69.9	2,240.9	21	10.9	157.8	4	2.1	97.9	1	0.5	----	193	71.2	543.8
2002	8	4.7	108.2	112	65.9	1,687.9	28	16.5	195.4	2	1.2	46.1	0	0.0	----	170	72.3	457.5
2003	3	2.0	38.2	115	77.2	1,574.6	13	8.7	84.3	3	2.0	66.2	0	0.0	----	149	71.6	380.5
2004	5	2.7	60.8	145	77.5	1,851.0	17	9.1	103.5	3	1.6	64.3	0	0.0	----	187	66.1	456.5
2005	10	4.4	123.1	172	75.8	2,360.3	26	11.5	171.3	1	0.4	22.1	1	0.4	----	227	65.4	596.2


1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 38 cases reported as unknown race/ethnicity in 2001; 29 in 2002; 24 in 2003; 21 in 2004; 34 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.

Figure 7. Reported Gonorrhea Rates for 20-24 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 17. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	96	9.8	178.5	426	43.6	2,204.9	196	20.1	136.6	6	0.6	18.7	9	0.9	----	976	48.6	293.1
2002	97	9.9	175.6	400	40.8	1,942.2	233	23.8	158.6	19	1.9	56.1	4	0.4	----	980	49.7	292.2
2003	112	11.7	184.8	393	41.2	1,710.8	249	26.1	158.2	21	2.2	55.8	3	0.3	----	954	100.0	278.3
2004	113	9.1	180.2	589	47.3	2,480.2	288	23.1	183.6	20	1.6	51.4	5	0.4	----	1,246	100.0	358.3
2005	146	11.0	233.5	575	43.4	2,405.8	332	25.1	212.3	21	1.6	59.7	7	0.5	----	1,325	100.0	387.3
FEMALE⁴																		
2001	74	7.2	139.6	504	48.9	2,302.3	188	18.2	141.3	13	1.3	39.6	3	0.3	----	1,031	51.4	323.6
2002	67	6.8	119.0	499	50.4	2,072.0	213	21.5	147.2	21	2.1	57.9	11	1.1	----	990	50.3	308.7
2003	105	9.5	183.7	508	45.8	2,059.6	255	23.0	173.4	34	3.1	90.8	2	0.2	----	1,109	53.8	338.3
2004	108	7.9	189.1	615	45.0	2,508.5	320	23.4	221.6	34	2.5	91.1	7	0.5	----	1,367	52.3	409.9
2005	194	12.6	319.7	647	42.1	2,623.4	384	25.0	263.9	39	2.5	114.4	10	0.7	----	1,535	53.7	478.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2000; 2 in 2001; 1 in 2002; 2 in 2004 and 5 in 2005.

3. Includes 492 cases reported as unknown race/ethnicity in 2001; 406 in 2002; 381 in 2003; 514 in 2004; and 505 in 2005.

4. Includes 1 Transgender: M to F case in 2001.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 18. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)³	48	2.4	250.4	80	4.1	376.9	70	3.4	297.1	132	5.0	511.2	124	4.3	425.8
Male	20	1.0	202.6	46	2.3	423.6	27	1.3	224.8	59	2.3	449.8	56	2.0	368.8
Female	27	1.3	290.4	34	1.7	327.9	43	2.1	372.3	73	2.8	574.6	68	2.4	487.8
East (7)	136	6.8	136.4	161	8.2	159.9	162	7.9	157.5	215	8.2	206.5	255	8.9	245.3
Male	65	3.2	127.9	81	4.1	158.0	75	3.6	143.1	98	3.7	184.8	125	4.4	234.5
Female	71	3.5	145.3	80	4.1	161.8	87	4.2	172.4	117	4.5	229.0	130	4.5	256.6
Metro (4)	239	11.9	270.7	255	12.9	302.0	246	11.9	298.4	299	11.4	369.2	331	11.6	448.1
Male	146	7.3	312.2	164	8.3	365.9	146	7.1	334.4	168	6.4	393.8	184	6.4	471.7
Female ⁴	93	4.6	224.0	91	4.6	229.7	100	4.8	257.9	131	5.0	341.8	147	5.1	421.8
San Fernando (2)	185	9.2	140.6	174	8.8	129.9	197	9.5	141.9	217	8.3	151.5	300	10.5	216.2
Male	98	4.9	144.4	77	3.9	111.2	102	4.9	142.1	99	3.8	134.1	158	5.5	218.6
Female	87	4.3	136.6	97	4.9	150.0	95	4.6	141.7	118	4.5	170.1	142	5.0	213.6
San Gabriel (3)	136	6.8	115.9	161	8.2	133.8	185	9.0	148.8	197	7.5	156.4	228	8.0	176.7
Male	66	3.3	110.0	87	4.4	141.8	85	4.1	134.1	101	3.9	157.1	87	3.0	131.2
Female	70	3.5	122.1	74	3.8	125.6	100	4.8	164.1	96	3.7	155.6	141	4.9	224.9
South (6)³	734	36.5	863.6	700	35.5	815.3	710	34.4	816.4	861	32.9	991.0	864	30.2	1013.0
Male	309	15.4	715.7	300	15.2	690.4	291	14.1	662.4	386	14.8	882.0	395	13.8	911.0
Female	424	21.1	1013.9	399	20.2	940.8	419	20.3	973.5	474	18.1	1099.2	468	16.3	1115.9
South Bay (8)³	305	15.2	473.4	270	13.7	412.9	276	13.4	406.1	369	14.1	524.9	377	13.2	536.3
Male	150	7.5	465.7	140	7.1	425.6	139	6.7	404.9	192	7.3	541.1	175	6.1	483.8
Female	155	7.7	481.2	130	6.6	400.0	137	6.6	407.3	177	6.8	508.4	201	7.0	589.0
West (5)	72	3.6	155.8	70	3.6	158.2	102	4.9	233.3	88	3.4	200.0	89	3.1	274.2
Male	36	1.8	161.4	35	1.8	163.2	53	2.6	248.4	51	2.0	236.0	48	1.7	291.0
Female	36	1.8	150.5	35	1.8	153.5	49	2.4	218.9	37	1.4	165.3	41	1.4	256.8
Unknown	154	7.7	-----	100	5.1	-----	115	5.6	-----	237	9.1	-----	297	10.4	-----
Total	2,009		308.3	1,971		300.4	2,063		307.6	2,615		383.8	2,865		432.3

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 2 cases reported with "unknown/other" gender in 2001; 1 in 2002; 1 in 2004; and 2 in 2005.

4. Includes 1 case reported as Transgender:M to F in 2001.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-1. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	5.0	31.7	11	55.0	993.0	3	15.0	108.1	0	0.0	0.0	0	0.0	----	20	42.6	202.6
2002	5	10.9	148.8	26	56.5	2,070.9	2	4.3	67.4	0	0.0	0.0	1	2.2	----	46	57.5	423.6
2003	3	11.1	70.3	17	63.0	1,005.4	3	11.1	79.5	0	0.0	0.0	0	0.0	----	27	38.6	224.8
2004	8	13.6	190.7	34	57.6	1,905.1	4	6.8	106.4	0	0.0	0.0	0	0.0	----	59	44.7	449.8
2005	5	8.9	97.4	29	51.8	1,816.2	8	14.3	196.7	0	0.0	0.0	0	0.0	----	56	45.2	368.8
FEMALE																		
2001	5	18.5	183.6	11	40.7	1,197.4	2	7.4	85.9	0	0.0	0.0	0	0.0	----	27	57.4	290.4
2002	4	11.8	117.9	17	50.0	1,373.1	4	11.8	133.1	1	2.9	398.7	0	0.0	----	34	42.5	327.9
2003	8	18.6	222.4	22	51.2	1,567.5	2	4.7	61.2	0	0.0	0.0	0	0.0	----	43	61.4	372.3
2004	13	17.8	324.7	32	43.8	1,867.6	11	15.1	290.2	0	0.0	0.0	1	1.4	----	73	55.3	574.6
2005	18	26.5	336.8	28	41.2	1,749.2	9	13.2	226.3	0	0.0	0.0	1	1.5	----	68	54.8	487.8

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2001.

3. Includes 14 cases reported as unknown race/ethnicity in 2001; 20 in 2002; 15 in 2003; 29 in 2004; and 26 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-2. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	21	21.4	115.5	24	24.5	1,038.4	29	29.6	106.2	2	2.0	38.1	1	1.0	----	98	53.0	144.4
2002	16	20.8	101.8	11	14.3	533.7	22	28.6	94.6	2	2.6	42.5	0	0.0	----	77	44.3	111.2
2003	24	23.5	120.8	12	11.8	462.1	41	40.2	144.8	4	3.9	65.6	0	0.0	----	102	51.8	142.1
2004	20	20.2	100.0	25	25.3	959.6	26	26.3	95.8	2	2.0	32.0	3	3.0	----	99	45.6	134.1
2005	47	29.7	206.5	32	20.3	1,662.4	45	28.5	162.6	4	2.5	62.4	1	0.6	----	158	52.7	218.6
FEMALE																		
2001	22	25.3	119.5	19	21.8	814.0	29	33.3	113.7	1	1.1	18.1	0	0.0	----	87	47.0	136.6
2002	11	11.3	66.5	20	20.6	908.7	36	37.1	154.6	4	4.1	77.7	0	0.0	----	97	55.7	150.0
2003	19	20.0	99.9	16	16.8	625.5	33	34.7	124.1	9	9.5	149.7	0	0.0	----	95	48.2	141.7
2004	27	22.9	132.1	14	11.9	508.8	52	44.1	186.8	5	4.2	77.6	0	0.0	----	118	54.4	170.1
2005	42	29.6	191.7	28	19.7	1,492.3	44	31.0	172.6	4	2.8	64.5	1	0.7	----	142	47.3	213.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 37 cases reported as unknown race/ethnicity in 2001; 52 in 2002; 39 in 2003; 43 in 2004; and 52 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-3. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	9	13.6	101.2	12	18.2	717.6	26	39.4	104.3	3	4.5	25.0	3	4.5	----	66	48.5	110.0
2002	7	8.0	85.6	13	14.9	790.9	38	43.7	165.9	5	5.7	43.2	0	0.0	----	87	54.0	141.8
2003	9	10.6	104.6	11	12.9	591.2	40	47.1	164.8	4	4.7	31.2	0	0.0	----	85	45.9	134.1
2004	11	10.9	116.0	20	19.8	921.8	50	49.5	183.0	4	4.0	26.8	0	0.0	----	101	51.3	157.1
2005	7	8.0	63.7	21	24.1	979.5	39	44.8	136.2	6	6.9	43.8	0	0.0	----	87	38.2	131.2
FEMALE																		
2001	14	20.0	165.1	20	28.6	1,192.3	19	27.1	81.1	2	2.9	17.7	0	0.0	----	70	51.5	122.1
2002	13	17.6	141.3	9	12.2	474.6	38	51.4	147.3	2	2.7	15.4	1	1.4	----	74	46.0	125.6
2003	19	19.0	204.0	17	17.0	853.5	44	44.0	163.7	6	6.0	45.6	0	0.0	----	100	54.1	167.0
2004	10	10.4	108.9	23	24.0	1111.7	40	41.7	144.8	10	10.4	74.8	1	1.0	----	96	48.7	160.0
2005	21	14.9	186.1	18	12.8	813.2	77	54.6	267.8	10	7.1	73.4	0	0.0	----	141	61.8	224.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 28 cases reported as unknown race/ethnicity in 2001; 35 in 2002; 35 in 2003; 28 in 2004; and 29 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 19-4. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	38	26.0	671.1	26	17.8	1,480.7	52	35.6	210.1	0	0.0	0.0	0	0.0	----	146	61.1	312.2
2002	38	23.2	734.5	24	14.6	1,436.7	57	34.8	244.8	8	4.9	170.6	1	0.6	----	164	64.3	365.9
2003	33	22.6	599.5	23	15.8	1,311.9	59	40.4	245.2	6	4.1	119.9	1	0.7	----	146	59.3	334.4
2004	31	18.5	514.3	44	26.2	2,485.4	68	40.5	282.2	3	1.8	56.9	1	0.6	----	168	56.2	393.8
2005	35	19.0	1,132.2	32	17.4	2,220.6	71	38.6	327.5	4	2.2	96.4	2	1.1	----	184	55.6	471.7
FEMALE³																		
2001	16	17.2	297.7	26	28.0	1,596.3	28	30.1	136.9	3	3.2	59.4	0	0.0	----	93	38.9	224.0
2002	6	6.6	116.6	32	35.2	1,930.6	30	33.0	143.1	7	7.7	135.6	1	1.1	----	91	35.7	229.7
2003	8	8.0	180.8	31	31.0	2,163.8	30	30.0	164.2	5	5.0	111.5	0	0.0	----	100	40.7	257.9
2004	16	12.2	297.4	49	37.4	2,960.7	43	32.8	205.2	5	3.8	95.4	1	0.8	----	131	43.8	341.8
2005	22	15.0	647.2	42	28.6	2,846.5	55	37.4	262.3	9	6.1	194.5	1	0.7	----	147	44.4	421.8

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 50 cases reported as unknown race/ethnicity in 2001; 51 in 2002; 50 in 2003; 38 in 2004; and 58 in 2005.

3. Includes 1 Transgender: M to F case in 2001.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-5. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	11	30.6	137.2	10	27.8	923.3	4	11.1	105.9	1	2.8	27.4	1	2.8	----	36	50.0	161.4
2002	14	40.0	167.4	8	22.9	673.9	4	11.4	101.7	2	5.7	55.0	0	0.0	----	35	50.0	163.2
2003	26	49.1	267.2	7	13.2	509.9	10	18.9	231.8	4	7.5	104.3	1	1.9	----	53	52.0	248.4
2004	14	27.5	136.8	15	29.4	1,084.4	11	21.6	265.2	5	9.8	136.9	1	2.0	----	51	58.0	236.0
2005	16	33.3	249.2	13	27.1	1,409.0	7	14.6	183.9	1	2.1	55.4	1	2.1	----	48	53.9	291.0
FEMALE																		
2001	8	22.2	83.7	11	30.6	794.6	9	25.0	205.1	2	5.6	39.1	1	2.8	----	36	50.0	150.5
2002	10	28.6	125.6	8	22.9	660.0	5	14.3	132.6	2	5.7	46.7	2	5.7	----	35	50.0	153.5
2003	17	34.7	203.6	9	18.4	717.7	6	12.2	157.6	6	12.2	140.4	1	2.0	----	49	48.0	218.9
2004	8	21.6	82.2	10	27.0	721.2	8	21.6	194.8	6	16.2	129.7	1	2.7	----	37	42.0	165.3
2005	15	36.6	225.0	7	17.1	744.4	8	19.5	230.4	4	9.8	187.5	0	0.0	----	41	46.1	256.8

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 14 cases reported as unknown race/ethnicity in 2001; 15 in 2002; 15 in 2003; 9 in 2004; 17 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-6. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	3	1.0	147.7	218	70.6	2,931.3	25	8.1	104.0	0	0.0	0.0	3	1.0	----	309	42.2	715.7
2002	6	2.0	286.4	203	67.7	2,549.8	43	14.3	169.1	1	0.3	80.8	1	0.3	----	300	42.9	690.4
2003	3	1.0	140.3	219	75.3	2,556.2	36	12.4	132.5	2	0.7	153.3	0	0.0	----	291	41.0	662.4
2004	5	1.3	255.2	289	74.9	3,486.6	39	10.1	148.8	0	0.0	0.0	0	0.0	----	386	44.9	882.0
2005	6	1.5	469.3	295	74.7	2,757.2	49	12.4	187.8	3	0.8	425.5	1	0.3	----	395	45.8	911.0
FEMALE																		
2001	1	0.2	56.8	296	69.8	3,238.9	51	12.0	226.6	2	0.5	163.9	2	0.5	----	424	57.8	1,013.9
2002	7	1.8	365.1	295	73.9	3,015.1	48	12.0	196.7	0	0.0	0.0	4	1.0	----	399	57.1	940.8
2003	3	0.7	155.7	300	71.6	3,002.4	73	17.4	288.7	1	0.2	69.0	0	0.0	----	419	59.0	973.5
2004	8	1.7	432.0	338	71.3	3,438.2	75	15.8	297.2	1	0.2	71.4	0	0.0	----	474	55.1	1,099.2
2005	7	1.5	563.9	350	74.8	3,029.0	61	13.0	249.2	4	0.9	502.1	4	0.9	----	468	54.2	1,115.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 2 cases reported with "unknown/other" gender in 2000; 1 in 2001; 1 in 2002; 1 in 2004; and 1 in 2005.

3. Includes 132 cases reported as unknown race/ethnicity in 2001; 91 in 2002; 73 in 2003; 105 in 2004; and 83 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-7. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	6	9.2	132.2	12	18.5	1,293.4	28	43.1	101.1	0	0.0	0.0	0	0.0	----	65	47.8	127.9
2002	5	6.2	101.6	12	14.8	1,051.0	46	56.8	150.5	0	0.0	0.0	0	0.0	----	81	50.3	158.0
2003	9	12.0	192.4	8	10.7	644.3	39	52.0	130.6	0	0.0	0.0	0	0.0	----	75	46.3	143.1
2004	10	10.2	189.4	17	17.3	1,089.1	55	56.1	159.2	2	2.0	51.3	0	0.0	----	98	45.6	184.8
2005	9	7.2	143.9	18	14.4	1,498.7	78	62.4	229.5	1	0.8	25.2	1	0.8	----	125	49.0	234.5
FEMALE																		
2001	2	2.8	50.3	11	15.5	1,279.9	31	43.7	130.9	1	1.4	42.5	0	0.0	----	71	52.2	145.3
2002	8	10.0	179.9	14	17.5	1,314.6	35	43.8	128.5	0	0.0	0.0	1	1.3	----	80	49.7	161.8
2003	12	13.8	254.4	10	11.5	805.8	44	50.6	146.6	1	1.1	31.1	1	1.1	----	87	53.7	172.4
2004	6	5.1	141.8	17	14.5	1,386.3	60	51.3	214.0	1	0.9	32.7	0	0.0	----	117	54.4	229.0
2005	16	12.3	250.6	32	24.6	2,693.5	66	50.8	198.4	1	0.8	26.2	0	0.0	----	130	51.0	256.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 45 cases reported as unknown race/ethnicity in 2001; 40 in 2002; 38 in 2003; 47 in 2004; and 33 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 19-8. Reported Gonorrhea Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	4	2.7	71.1	104	69.3	2,328.8	24	16.0	165.5	0	0.0	0.0	1	0.7	----	150	49.2	465.7
2002	4	2.9	74.8	97	69.3	2,238.3	13	9.3	95.1	1	0.7	26.4	1	0.7	----	140	51.9	425.6
2003	2	1.4	36.3	89	64.0	1,997.0	18	12.9	134.3	1	0.7	25.5	1	0.7	----	139	50.4	404.9
2004	6	3.1	93.9	129	67.2	2,531.0	28	14.6	191.5	3	1.6	66.9	0	0.0	----	192	52.0	541.1
2005	11	6.3	155.3	116	66.3	2,213.9	26	14.9	173.7	1	0.6	21.6	1	0.6	----	175	46.5	483.8
FEMALE																		
2001	5	3.2	96.8	107	69.0	1,962.5	17	11.0	131.2	2	1.3	56.0	0	0.0	----	155	50.8	481.2
2002	5	3.8	91.0	94	72.3	1,579.7	14	10.8	98.6	5	3.8	125.1	1	0.8	----	130	48.1	400.0
2003	7	5.1	126.7	92	67.2	1,568.8	16	11.7	114.8	5	3.6	123.6	0	0.0	----	137	49.6	407.3
2004	5	2.8	87.3	112	63.3	1,919.2	27	15.3	195.7	4	2.3	96.0	3	1.7	----	177	48.0	508.4
2005	8	4.0	126.2	124	61.7	2,329.3	37	18.4	279.6	3	1.5	71.3	0	0.0	----	201	53.5	589.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Excludes 1 case reported with "unknown/other" gender in 2005.

3. Includes 41 cases reported as unknown race/ethnicity in 2001; 35 in 2002; 45 in 2003; 52 in 2004; and 49 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.





Figure 8. 2005 Reported Gonorrhea Rates Per 100,000 10-24 Year Olds, by Service Planning Area (SPA).

Figure 9. Reported Early Syphilis Rates for 10-14 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 20. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	1	100.0	0.4	0	0.0	0.0	0	0.0	----	1	100.0	0.3
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	1	100.0	2.6	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	50.0	0.2
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	1	33.3	2.6	2	66.7	1.0	0	0.0	0.0	0	0.0	----	3	100.0	0.8
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	1	100.0	2.6	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	50.0	0.3
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 21. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
East (7)	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Metro (4)	0	0.0	0.0	1	100.0	1.2	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	1	100.0	2.3	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
San Fernando (2)	1	33.3	0.6	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	1	33.3	1.3	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
San Gabriel (3)	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
South (6)	1	33.3	1.0	0	0.0	0.0	0	0.0	0.0	1	50.0	0.9	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	1	50.0	1.9	0	0.0	0.0
Female	1	33.3	2.1	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
South Bay (8)	1	33.3	1.2	0	0.0	0.0	0	0.0	0.0	1	50.0	1.1	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	1	33.3	2.4	0	0.0	0.0	0	0.0	0.0	1	50.0	2.2	0	0.0	0.0
West (5)	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Unknown	0	0.0	----	0	0.0	----	0	0.0	----	0	0.0	----	0	0.0	----
Total	3		0.4	1		0.1	0		0.0	2		0.3	0		0.0

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

Note: Cumulative percentages may not equal 100% because of rounding.

**Table 22-1. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity
Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005**

	White			Black			Hispanic			Asian-Pacific			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE															
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
FEMALE															
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 22-2. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	2.8	0	0.0	0.0	0	0.0	----	1	100.0	1.3
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 22-3. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 22-4. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	1	100.0	3.3	0	0.0	0.0	0	0.0	----	1	100.0	2.3
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 22-5. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 22-6. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	1	0.0	7.3	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	100.0	1.9
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	3.1	0	0.0	0.0	0	0.0	----	1	100.0	2.1
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 22-7. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 22-8. Reported Early Syphilis Cases and Rates Per 100,000 10-14 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005


	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	1	100.0	11.5	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	100.0	2.4
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	1	100.0	11.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	100.0	2.2
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.

Figure 10. Reported Early Syphilis Rates for 15-19 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 23. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}			
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Pop	Adj. Rate
MALE																			
2001	2	28.6	2.5	1	14.3	3.0	4	57.1	2.1	0	0.0	0.0	0	0.0	----	7	46.7	346,506	2.0
2002	0	0.0	0.0	3	17.6	8.8	13	76.5	6.7	1	5.9	2.3	0	0.0	----	17	81.0	355,792	4.8
2003	1	9.1	1.2	3	27.3	8.6	7	63.6	3.5	0	0.0	0.0	0	0.0	----	11	100.0	367,144	3.0
2004	1	10.0	1.3	2	20.0	6.2	6	60.0	3.2	0	0.0	0.0	0	0.0	----	10	100.0	379,144	2.6
2005	0	0.0	0.0	5	41.7	13.5	7	58.3	3.6	0	0.0	0.0	0	0.0	----	12	100.0	357,535	3.4
FEMALE³																			
2001	0	0.0	0.0	0	0.0	0.0	7	87.5	4.0	0	0.0	0.0	0	0.0	----	8	53.3	328,681	2.4
2002	0	0.0	0.0	1	25.0	2.9	3	75.0	1.6	0	0.0	0.0	0	0.0	----	4	19.0	340,458	1.2
2003	0	0.0	0.0	1	16.7	2.8	5	83.3	2.6	0	0.0	0.0	0	0.0	----	6	100.0	353,965	1.7
2004	0	0.0	0.0	5	45.5	13.5	5	45.5	2.5	1	9.1	2.3	0	0.0	----	11	100.0	367,521	3.0
2005	0	0.0	0.0	8	47.1	22.1	9	52.9	4.8	0	0.0	0.0	0	0.0	----	17	100.0	343,548	4.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2004.

3. Includes 1 Transgender: M to F case in 2001.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 24. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)	0	0.0	0.0	2	9.5	6.7	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	2	9.5	12.9	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
East (7)	2	14.3	1.9	4	19.0	3.7	1	5.9	0.9	5	23.8	4.4	2	6.9	1.8
Male	2	14.3	3.7	4	19.0	7.2	1	5.9	1.8	1	4.8	1.7	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	4	19.0	7.2	2	6.9	3.7
Metro (4)	6	42.9	8.1	6	28.6	7.8	5	29.4	6.2	2	9.5	2.3	9	31.0	12.4
Male	2	14.3	5.2	5	23.8	12.5	4	23.5	9.5	1	4.8	2.2	6	20.7	16.2
Female ³	4	28.6	11.3	1	4.8	2.7	1	5.9	2.5	1	4.8	2.4	3	10.3	8.5
San Fernando (2)	2	14.3	1.4	1	4.8	0.7	2	11.8	1.3	2	9.5	1.3	1	3.4	0.7
Male	2	14.3	2.7	1	4.8	1.3	1	5.9	1.3	1	4.8	1.2	1	3.4	1.3
Female	0	0.0	0.0	0	0.0	0.0	1	5.9	1.3	1	4.8	1.3	0	0.0	0.0
San Gabriel (3)	1	7.1	0.8	3	14.3	2.3	3	17.6	2.3	3	14.3	2.3	2	6.9	1.5
Male	0	0.0	0.0	3	14.3	4.5	1	5.9	1.5	2	9.5	3.0	1	3.4	1.5
Female	1	7.1	1.6	0	0.0	0.0	2	11.8	3.1	1	4.8	1.6	1	3.4	1.6
South (6)	2	14.3	2.3	2	9.5	2.2	4	23.5	4.3	8	38.1	8.4	10	34.5	10.7
Male	1	7.1	2.3	0	0.0	0.0	2	11.8	4.3	4	19.0	8.4	3	10.3	6.4
Female	1	7.1	2.3	2	9.5	4.5	2	11.8	4.4	4	19.0	8.5	7	24.1	15.2
South Bay (8)	1	7.1	1.4	3	14.3	4.0	1	5.9	1.3	1	4.8	1.2	4	13.8	5.1
Male	0	0.0	0.0	2	9.5	5.2	1	5.9	2.5	1	4.8	2.4	0	0.0	0.0
Female	1	7.1	2.8	1	4.8	2.7	0	0.0	0.0	0	0.0	0.0	4	13.8	10.5
West (5)	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
Unknown	0	0.0	-----	0	0.0	-----	1	5.9	-----	0	0.0	-----	1	3.4	-----
Total	14		2.1	21		3.0	17		2.4	21		2.8	29		4.1

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 1 Transgender:M to F case in 2001.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-1. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	1	50.0	38.0	1	50.0	17.8	0	0.0	0.0	0	0.0	----	2	100.0	12.9
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-2. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	1	50.0	31.9	1	50.0	3.0	0	0.0	0.0	0	0.0	----	2	100.0	2.7
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	1	100.0	12.5	0	0.0	----	1	100.0	1.3
2003	0	0.0	0.0	1	100.0	29.3	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	50.0	1.3
2004	0	0.0	0.0	0	0.0	0.0	1	100.0	2.6	0	0.0	0.0	0	0.0	----	1	50.0	1.2
2005	0	0.0	0.0	1	100.0	35.7	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	100.0	1.3
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	2.9	0	0.0	0.0	0	0.0	----	1	50.0	1.3
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	1	100.0	12.6	0	0.0	----	1	50.0	1.3
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-3. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	3	100.0	8.8	0	0.0	0.0	0	0.0	----	3	100.0	4.5
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	2.9	0	0.0	0.0	0	0.0	----	1	33.3	1.5
2004	0	0.0	0.0	0	0.0	0.0	2	100.0	5.6	0	0.0	0.0	0	0.0	----	2	66.7	3.0
2005	0	0.0	0.0	0	0.0	0.0	1	100.0	2.8	0	0.0	0.0	0	0.0	----	1	50.0	1.5
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	3.1	0	0.0	0.0	0	0.0	----	1	100.0	1.6
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	2	100.0	5.9	0	0.0	0.0	0	0.0	----	2	66.7	3.1
2004	0	0.0	0.0	1	100.0	38.9	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	33.3	1.6
2005	0	0.0	0.0	0	0.0	0.0	1	100.0	2.9	0	0.0	0.0	0	0.0	----	1	50.0	1.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 25-4. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	50.0	19.4	0	0.0	0.0	1	50.0	3.8	0	0.0	0.0	0	0.0	----	2	33.3	5.2
2002	0	0.0	0.0	1	20.0	51.7	4	80.0	15.2	0	0.0	0.0	0	0.0	----	5	83.3	12.5
2003	1	25.0	14.8	0	0.0	0.0	3	75.0	11.0	0	0.0	0.0	0	0.0	----	4	80.0	9.5
2004	0	0.0	0.0	0	0.0	0.0	1	100.0	3.5	0	0.0	0.0	0	0.0	----	1	50.0	2.2
2005	0	0.0	0.0	1	16.7	50.5	5	83.3	19.0	0	0.0	0.0	0	0.0	----	6	66.7	16.2
FEMALE³																		
2001	0	0.0	0.0	0	0.0	0.0	4	100.0	17.1	0	0.0	0.0	0	0.0	----	4	66.7	11.3
2002	0	0.0	0.0	0	0.0	0.0	1	100.0	4.2	0	0.0	0.0	0	0.0	----	1	16.7	2.7
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	4.0	0	0.0	0.0	0	0.0	----	1	20.0	2.5
2004	0	0.0	0.0	1	100.0	48.4	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	50.0	2.4
2005	0	0.0	0.0	1	33.3	59.0	2	66.7	7.9	0	0.0	0.0	0	0.0	----	3	33.3	8.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

3. Includes 1 Transgender: M to F case in 2001.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-5. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-6. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	3.4	0	0.0	0.0	0	0.0	----	1	50.0	2.3
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	1	50.0	8.1	1	50.0	3.2	0	0.0	0.0	0	0.0	----	2	50.0	4.3
2004	0	0.0	0.0	2	50.0	21.7	1	25.0	4.0	0	0.0	0.0	0	0.0	----	4	50.0	8.4
2005	0	0.0	0.0	2	66.7	13.2	1	33.3	3.3	0	0.0	0.0	0	0.0	----	3	30.0	6.4
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	3.6	0	0.0	0.0	0	0.0	----	1	50.0	2.3
2002	0	0.0	0.0	0	0.0	0.0	2	100.0	7.0	0	0.0	0.0	0	0.0	----	2	100.0	4.5
2003	0	0.0	0.0	1	50.0	7.6	1	50.0	3.3	0	0.0	0.0	0	0.0	----	2	50.0	4.4
2004	0	0.0	0.0	2	50.0	15.0	2	50.0	6.4	0	0.0	0.0	0	0.0	----	4	50.0	8.5
2005	0	0.0	0.0	3	42.9	19.6	4	57.1	13.7	0	0.0	0.0	0	0.0	----	7	70.0	15.2

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-7. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	50.0	13.5	0	0.0	0.0	1	50.0	2.5	0	0.0	0.0	0	0.0	----	2	100.0	3.7
2002	0	0.0	0.0	0	0.0	0.0	4	100.0	9.6	0	0.0	0.0	0	0.0	----	4	100.0	7.2
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	2.3	0	0.0	0.0	0	0.0	----	1	100.0	1.8
2004	0	0.0	0.0	0	0.0	0.0	1	100.0	2.2	0	0.0	0.0	0	0.0	----	1	20.0	1.7
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	1	25.0	47.1	3	75.0	7.0	0	0.0	0.0	0	0.0	----	4	80.0	7.2
2005	0	0.0	0.0	0	0.0	0.0	2	100.0	4.9	0	0.0	0.0	0	0.0	----	2	100.0	3.7

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 25-8. Reported Early Syphilis Cases and Rates Per 100,000 15-19 Year Olds by Gender and Race/Ethnicity, South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005


	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	1	50.0	14.0	1	50.0	5.9	0	0.0	0.0	0	0.0	----	2	66.7	5.2
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	5.6	0	0.0	0.0	0	0.0	----	1	100.0	2.5
2004	1	100.0	10.5	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	100.0	2.4
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	6.4	0	0.0	0.0	0	0.0	----	1	100.0	2.8
2002	0	0.0	0.0	1	100.0	13.3	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	33.3	2.7
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	4	100.0	50.8	0	0.0	0.0	0	0.0	0.0	0	0.0	----	4	100.0	10.5

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.

Figure 11. Reported Early Syphilis Rates for 20-24 Year Olds by Year, Los Angeles County (LAC), 2001-2005



(1) LA County Sexually Transmitted Disease Program, Epidemiology Unit, 2006.
Note: Excludes cases from Pasadena and Long Beach.

Table 26. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	2	7.7	2.8	2	7.7	7.8	22	84.6	11.5	0	0.0	0.0	0	0.0	----	26	76.5	7.8
2002	7	12.5	9.9	12	21.4	45.5	34	60.7	18.1	1	1.8	2.3	1	1.8	----	56	76.7	16.7
2003	9	17.0	12.1	6	11.3	21.3	36	67.9	18.6	2	3.8	4.3	0	0.0	----	53	100.0	15.5
2004	20	26.0	26.3	11	14.3	38.2	43	55.8	22.6	2	2.6	4.2	0	0.0	----	77	100.0	22.1
2005	18	18.2	24.2	15	15.2	52.8	55	55.6	29.6	7	7.1	16.7	1	1.0	----	99	100.0	28.9
FEMALE³																		
2001	1	12.5	1.4	0	0.0	0.0	7	87.5	4.0	0	0.0	0.0	0	0.0	----	8	23.5	2.5
2002	1	5.9	1.5	5	29.4	17.0	11	64.7	6.2	0	0.0	0.0	0	0.0	----	17	23.3	5.3
2003	2	14.3	3.1	2	14.3	7.2	8	57.1	4.8	0	0.0	0.0	0	0.0	----	14	100.0	4.3
2004	1	5.6	1.4	8	44.4	25.8	6	33.3	3.3	1	5.6	2.1	1	5.6	----	18	100.0	5.4
2005	4	17.4	5.7	5	21.7	17.6	12	52.2	7.1	1	4.3	2.5	0	0.0	----	23	100.0	7.2

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2002; 1 in 2003; 1 in 2004; and 4 in 2005.

3. Includes 1 Transgender: M to F cases in 2003; 1 in 2004.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena and Long Beach.

Table 27. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Service Planning Area (SPA)¹ and Gender, Los Angeles County, 2001-2005

	2001			2002			2003			2004			2005		
	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate	No.	(%)	Rate
SPA²/Gender															
Antelope Valley (1)	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	1	1.1	3.9	1	0.8	3.4
Male	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	1	1.1	7.6	1	0.8	6.6
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0
East (7)	8	23.5	8.0	4	5.5	4.0	9	13.4	8.7	18	18.9	17.3	12	9.8	11.5
Male	4	11.8	7.9	3	4.1	5.9	8	11.9	15.3	17	17.9	32.1	10	8.2	18.8
Female	4	11.8	8.2	1	1.4	2.0	1	1.5	2.0	1	1.1	2.0	2	1.6	3.9
Metro (4)³	13	38.2	14.7	32	43.8	37.9	19	28.4	18.5	32	33.7	39.5	42	34.4	56.9
Male	13	38.2	27.8	27	37.0	60.2	16	23.9	30.5	29	30.5	68.0	34	27.9	87.2
Female	0	0.0	0.0	5	6.8	12.6	2	3.0	4.0	3	3.2	7.8	8	6.6	23.0
San Fernando (2)	4	11.8	3.0	8	11.0	6.0	15	22.4	10.8	13	13.7	9.1	17	13.9	12.3
Male	3	8.8	4.4	4	5.5	5.8	11	16.4	15.3	11	11.6	14.9	13	10.7	18.0
Female	1	2.9	1.6	4	5.5	6.2	4	6.0	6.0	2	2.1	2.9	4	3.3	6.0
San Gabriel (3)	5	14.7	4.3	5	6.8	4.2	4	6.0	3.2	5	5.3	4.0	9	7.4	7.0
Male	3	8.8	5.0	4	5.5	6.5	2	3.0	3.2	4	4.2	6.2	8	6.6	12.1
Female	2	5.9	3.5	1	1.4	1.7	2	3.0	3.3	1	1.1	1.6	1	0.8	1.6
South (6)³	3	8.8	3.5	7	9.6	8.2	12	17.9	13.8	13	13.7	15.0	19	15.6	22.3
Male	2	5.9	4.6	3	4.1	6.9	9	13.4	20.5	6	6.3	13.7	14	11.5	32.3
Female	1	2.9	2.4	4	5.5	9.4	3	4.5	7.0	6	6.3	13.9	5	4.1	11.9
South Bay (8)	0	0.0	0.0	13	17.8	19.9	4	6.0	5.9	5	5.3	7.1	9	7.4	12.8
Male	0	0.0	0.0	11	15.1	33.4	3	4.5	8.7	3	3.2	8.5	6	4.9	16.6
Female	0	0.0	0.0	2	2.7	6.2	1	1.5	3.0	2	2.1	5.7	3	2.5	8.8
West (5)	1	2.9	2.2	4	5.5	9.0	3	4.5	6.9	6	6.3	13.6	3	2.5	9.2
Male	1	2.9	4.5	4	5.5	18.7	3	4.5	14.1	5	5.3	23.1	3	2.5	18.2
Female	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	1	1.1	4.5	0	0.0	0.0
Unknown	0	0.0	-----	0	0.0	-----	1	1.5	-----	2	2.1	-----	10	8.2	-----
Total	34		5.2	73		11.1	67		9.7	95		13.9	122		18.4

1. Excludes cases from Pasadena (SPA 3) and Long Beach (SPA 8).

2. Main categories.

3. Includes 1 Transgender:M to F case in 2003; and 1 in 2004.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 28-1. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Antelope Valley Service Planning Area 1 (SPA 1), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	1	100.0	20.7	0	0.0	0.0	0	0.0	----	1	100.0	7.6
2005	0	0.0	0.0	0	0.0	0.0	1	100.0	0.0	0	0.0	0.0	0	0.0	----	1	100.0	6.6
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 28-2. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Fernando Service Planning Area 2 (SPA 2), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	33.3	4.3	0	0.0	0.0	2	66.7	5.7	0	0.0	0.0	0	0.0	----	3	75.0	4.4
2002	0	0.0	0.0	1	25.0	32.1	3	75.0	8.5	0	0.0	0.0	0	0.0	----	4	50.0	5.8
2003	3	27.3	0.0	1	9.1	0.0	6	54.5	0.0	1	9.1	0.0	0	0.0	----	11	73.3	15.3
2004	5	45.5	0.0	3	27.3	0.0	2	18.2	0.0	0	0.0	0.0	0	0.0	----	11	84.6	14.9
2005	4	30.8	0.0	0	0.0	0.0	7	53.8	0.0	2	15.4	0.0	0	0.0	----	13	76.5	18.0
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	3.2	0	0.0	0.0	0	0.0	----	1	25.0	1.6
2002	1	25.0	4.4	0	0.0	0.0	3	75.0	9.4	0	0.0	0.0	0	0.0	----	4	50.0	6.2
2003	1	25.0	0.0	1	25.0	0.0	2	50.0	0.0	0	0.0	0.0	0	0.0	----	4	26.7	6.0
2004	0	0.0	0.0	0	0.0	0.0	2	100.0	0.0	0	0.0	0.0	0	0.0	----	2	15.4	2.9
2005	1	25.0	0.0	1	25.0	0.0	2	50.0	0.0	0	0.0	0.0	0	0.0	----	4	23.5	6.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 28-3. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, San Gabriel Service Planning Area 3 (SPA 3), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	3	100.0	9.6	0	0.0	0.0	0	0.0	----	3	60.0	5.0
2002	0	0.0	0.0	0	0.0	0.0	3	75.0	9.5	1	25.0	6.3	0	0.0	----	4	80.0	6.5
2003	0	0.0	0.0	0	0.0	0.0	2	100.0	0.0	0	0.0	0.0	0	0.0	----	2	50.0	3.2
2004	0	0.0	0.0	1	25.0	0.0	3	75.0	0.0	0	0.0	0.0	0	0.0	----	4	80.0	6.2
2005	2	25.0	0.0	1	12.5	0.0	5	62.5	0.0	0	0.0	0.0	0	0.0	----	8	88.9	12.1
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	2	100.0	6.7	0	0.0	0.0	0	0.0	----	2	40.0	3.5
2002	0	0.0	0.0	0	0.0	0.0	1	100.0	3.3	0	0.0	0.0	0	0.0	----	1	20.0	1.7
2003	0	0.0	0.0	0	0.0	0.0	2	100.0	0.0	0	0.0	0.0	0	0.0	----	2	50.0	3.3
2004	0	0.0	0.0	1	100.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	20.0	1.7
2005	0	0.0	0.0	0	0.0	0.0	1	100.0	0.0	0	0.0	0.0	0	0.0	----	1	11.1	1.6

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Pasadena.

Table 28-4. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, Metro Service Planning Area 4 (SPA 4), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	1	7.7	45.2	12	92.3	38.5	0	0.0	0.0	0	0.0	----	13	100.0	27.8
2002	4	14.8	62.7	6	22.2	291.0	15	55.6	52.2	0	0.0	0.0	1	3.7	----	27	84.4	60.2
2003	4	25.0	0.0	1	6.3	0.0	11	68.8	0.0	0	0.0	0.0	0	0.0	----	16	84.2	36.6
2004	8	27.6	0.0	2	6.9	0.0	19	65.5	0.0	0	0.0	0.0	0	0.0	----	29	90.6	68.0
2005	8	23.5	0.0	3	8.8	0.0	19	55.9	0.0	2	5.9	0.0	0	0.0	----	34	81.0	87.2
FEMALE³																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	1	20.0	50.3	4	80.0	15.9	0	0.0	0.0	0	0.0	----	5	15.6	12.6
2003	1	33.3	0.0	0	0.0	0.0	1	33.3	0.0	0	0.0	0.0	0	0.0	----	3	15.8	7.7
2004	0	0.0	0.0	0	0.0	0.0	3	100.0	0.0	0	0.0	0.0	0	0.0	----	3	9.4	7.8
2005	3	37.5	0.0	0	0.0	0.0	5	62.5	0.0	0	0.0	0.0	0	0.0	----	8	19.0	23.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2002.

3. Includes 1 Transgender: M to F case in 2003.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 28-5. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, West Service Planning Area 5 (SPA 5), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	19.7	0	0.0	0.0	0	0.0	----	1	100.0	4.5
2002	1	25.0	9.6	0	0.0	0.0	3	75.0	61.0	0	0.0	0.0	0	0.0	----	4	100.0	18.7
2003	0	0.0	0.0	0	0.0	0.0	2	66.7	0.0	1	33.3	0.0	0	0.0	----	3	100.0	14.1
2004	2	40.0	0.0	2	40.0	0.0	1	20.0	0.0	0	0.0	0.0	0	0.0	----	5	83.3	23.1
2005	1	33.3	0.0	0	0.0	0.0	1	33.3	0.0	1	33.3	0.0	0	0.0	----	3	100.0	18.2
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2004	1	100.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	16.7	4.5
2005	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 28-6. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, South Service Planning Area 6 (SPA 6), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ^{2,3}		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	1	50.0	10.8	1	50.0	3.3	0	0.0	0.0	0	0.0	----	2	66.7	4.6
2002	0	0.0	0.0	0	0.0	0.0	3	100.0	10.0	0	0.0	0.0	0	0.0	----	3	42.9	6.9
2003	0	0.0	0.0	4	44.4	0.0	5	55.6	0.0	0	0.0	0.0	0	0.0	----	9	75.0	20.5
2004	2	33.3	0.0	3	50.0	0.0	1	16.7	0.0	0	0.0	0.0	0	0.0	----	6	46.2	13.7
2005	0	0.0	0.0	7	50.0	0.0	7	50.0	0.0	0	0.0	0.0	0	0.0	----	14	73.7	32.3
FEMALE³																		
2001	0	0.0	0.0	0	0.0	0.0	1	100.0	3.7	0	0.0	0.0	0	0.0	----	1	33.3	2.4
2002	0	0.0	0.0	2	50.0	18.1	2	50.0	7.3	0	0.0	0.0	0	0.0	----	4	57.1	9.4
2003	0	0.0	0.0	1	33.3	0.0	2	66.7	0.0	0	0.0	0.0	0	0.0	----	3	25.0	7.0
2004	0	0.0	0.0	5	71.4	0.0	0	0.0	0.0	0	0.0	0.0	1	14.3	----	7	53.8	16.2
2005	0	0.0	0.0	3	60.0	0.0	1	20.0	0.0	0	0.0	0.0	0	0.0	----	5	26.3	11.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2005.

3. Includes 1 Transgender: M to F in 2004.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

Table 28-7. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity, East Service Planning Area 7 (SPA 7), Los Angeles County, 2001-2005

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	1	25.0	15.6	0	0.0	0.0	3	75.0	7.7	0	0.0	0.0	0	0.0	----	4	50.0	7.9
2002	1	33.3	15.8	0	0.0	0.0	2	66.7	5.1	0	0.0	0.0	0	0.0	----	3	75.0	5.9
2003	1	12.5	0.0	0	0.0	0.0	7	87.5	0.0	0	0.0	0.0	0	0.0	----	8	88.9	15.3
2004	3	17.6	0.0	0	0.0	0.0	12	70.6	0.0	2	11.8	0.0	0	0.0	----	17	94.4	32.1
2005	0	0.0	0.0	0	0.0	0.0	10	100.0	0.0	0	0.0	0.0	0	0.0	----	10	83.3	18.8
FEMALE																		
2001	1	25.0	15.9	0	0.0	0.0	3	75.0	8.0	0	0.0	0.0	0	0.0	----	4	50.0	8.2
2002	0	0.0	0.0	0	0.0	0.0	1	100.0	2.6	0	0.0	0.0	0	0.0	----	1	25.0	2.0
2003	0	0.0	0.0	0	0.0	0.0	1	100.0	0.0	0	0.0	0.0	0	0.0	----	1	11.1	2.0
2004	0	0.0	0.0	0	0.0	0.0	1	100.0	0.0	0	0.0	0.0	0	0.0	----	1	5.6	2.0
2005	0	0.0	0.0	0	0.0	0.0	2	100.0	0.0	0	0.0	0.0	0	0.0	----	2	16.7	3.9

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding.

**Table 28-8. Reported Early Syphilis Cases and Rates Per 100,000 20-24 Year Olds by Gender and Race/Ethnicity
South Bay Service Planning Area 8 (SPA 8), Los Angeles County, 2001-2005**

	White			Black			Hispanic			Asian-Pacific			Other ¹			Total ²		
	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate	No.	(%)	Adj. Rate
MALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	1	9.1	15.5	5	45.5	95.5	5	45.5	30.3	0	0.0	0.0	0	0.0	----	11	84.6	33.4
2003	1	33.3	0.0	0	0.0	0.0	2	66.7	0.0	0	0.0	0.0	0	0.0	----	3	75.0	8.7
2004	0	0.0	0.0	0	0.0	0.0	3	100.0	0.0	0	0.0	0.0	0	0.0	----	3	60.0	8.5
2005	1	16.7	0.0	3	50.0	0.0	1	16.7	0.0	0	0.0	0.0	0	0.0	----	6	66.7	16.6
FEMALE																		
2001	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	0	0.0	0.0
2002	0	0.0	0.0	2	100.0	30.7	0	0.0	0.0	0	0.0	0.0	0	0.0	----	2	15.4	6.2
2003	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	0.0	0	0.0	----	1	25.0	3.0
2004	0	0.0	0.0	1	50.0	0.0	0	0.0	0.0	1	50.0	0.0	0	0.0	----	2	40.0	5.7
2005	0	0.0	0.0	1	33.3	0.0	1	33.3	0.0	1	33.3	0.0	0	0.0	----	3	33.3	8.8

1. Includes data for ethnicity reported as "Other", "Native American" and "Mixed Race".

2. Includes 1 case reported as unknown race/ethnicity in 2003; and 1 in 2005.

These tables should be used only for race/ethnicity comparisons, not for overall totals or gender totals. This is because, if race/ethnicity was not specified, cases were prorated according to the distribution of cases for which race/ethnicity was specified.

Note: Cumulative percentages may not equal 100% because of rounding. Excludes cases from Long Beach.




Figure 12. 2005 Reported Early Syphilis Rates Per 100,000 10-24 Year Olds, by Service Planning Area (SPA).

SPECIAL PROJECTS

School and Community Testing

The STD Program has a long history of collaborating with local schools and community organizations to provide on-site testing for chlamydia and other STDs to youth and other at-risk groups. In 2001, as part of the *Get Tested!* Program sponsored by California DHS, the STD Program developed special protocols and materials for these collaborations, including intake and data collection instruments, and protocols for patient-delivered partner therapy. Using these protocols and materials, on-site screening has been provided throughout the county for continuation schools, community colleges, needle exchange programs and gang diversion programs.

The STD Program continues to collaborate with schools and other community organizations to provide on-site chlamydia testing. STDP offers all elements of chlamydia screening, including counseling and testing, treatment, and partner follow-up, as well as staff training for partnering agencies

Contact Harlan Rotblatt at (213) 744-5903 or Melina Boudov at (213) 744-5956.

Juvenile Hall

Rates of chlamydia and gonorrhea for detainees at LAC Juvenile Halls are among the highest of any group in the County. Incarcerated youth often have an earlier sexual debut, multiple sex partners, a history of sexual and physical abuse, and have engaged in high-risk behaviors, such as trading sex for drugs or money. These behaviors put them at significantly higher risk for acquiring STDs compared to non-incarcerated youth. In 1996, the STD Program initiated a project to address this problem in collaboration with Juvenile Court Health Services, the healthcare provider for adjudicated youth in the County. The project provides STD screening and treatment to females and males in the LAC Juvenile Hall system. In 2005, 21,651 youth were screened for chlamydia and gonorrhea at medical intake to the three LAC Juvenile Halls. There were 1,609 cases of chlamydia, and 379 cases of gonorrhea identified and treated at Juvenile Halls. Among females, 777 (15.6%) were positive for chlamydia and 248 (5.0%) were positive for gonorrhea. Among males, 830 (5.0%) were positive for chlamydia while only 131 (0.8%) were positive for gonorrhea. These rates are consistent with previous years. Chlamydia and gonorrhea screening is currently offered universally at intake to each facility. Excellent STD care, including screening and treatment, for these youth during their incarceration is a top priority for the STD Program.

All detainees that test positive for an STD are treated and receive partner elicitation and notification services. Females who test positive receive enhanced case management which includes individualized risk assessment, health education and risk reduction counseling, referrals for reproductive and other health needs, and help acquiring social and educational services.

Contact Melina Boudov at (213) 744-5956.

APPENDIX A: CASE DEFINITIONS FOR REPORTABLE STDs

California State law (California Administrative Code, Title 17, Public Health, Section 2500, 1996) requires all health care providers to report selected communicable diseases to local health departments. Six of these diseases are reported directly to the County STD Program using the STD Confidential Morbidity Report (STD CMR, Rev. 11/03). These six include: chlamydia, gonorrhea, syphilis, non-gonococcal/non-chlamydial urethritis (NGU), non-gonococcal/non-chlamydial pelvic inflammatory disease, and chancroid. The following case definitions should assist physicians and other health care providers with the reporting process. The clinical description for each disease is brief; a more thorough discussion can be found in Sexually Transmitted Diseases (3rd Edition, 1999) by Holmes et al. STD Program staff is available to answer questions concerning STD case definitions and STD reporting requirements. Three of the six most frequently reported STDs are diagnosed below.

CHLAMYDIAL INFECTION

1. Clinical description

Presence of mucopurulent cervical, urethral, or rectal discharge; epididymitis, salpingitis, or PID in adults; conjunctivitis or pneumonia in infants born to mothers with untreated infection. Asymptomatic infections are common.

2. Criteria for diagnosis

Laboratory identification of *Chlamydia trachomatis* infection by culture, antigen detection, or nucleic acid detection (amplified or non-amplified) methods.

GONORRHEA

1. Clinical description

Presence of purulent cervical, urethral, or rectal discharge; epididymitis, pharyngitis, salpingitis, or PID in adults; conjunctivitis in neonates born to mothers with untreated infection. Asymptomatic infections are common.

2. Criteria for diagnosis

Compatible clinical picture and sexual exposure to a person infected with *N. gonorrhoeae*, OR a, b, or c.

- a. Observation of Gram negative intracellular diplococci in urethral specimen from a male,
- b. Isolation of typical gram-negative, oxidase positive diplococci from a clinical specimen,
- c. Laboratory identification of *N. gonorrhoeae* infection by culture, antigen detection, or nucleic acid detection (amplified or non-amplified) methods.

SYPHILIS

NOTE: Neurosyphilis may co-exist with early stages of syphilis.

PRIMARY SYPHILIS

1. Clinical description

One or more painless indurated ulcers at the site of exposure. Lymphadenopathy is common.

2. Criteria for diagnosis

Compatible clinical picture AND laboratory confirmation by either a or b:

- a. Demonstration of *T. pallidum* in clinical specimen by darkfield, fluorescent antibody or equivalent microscopic methods,
- b. Reactive serologic test for syphilis.

** Nontreponemal tests often do not become reactive until 7-10 days following the lesion onset.

SECONDARY SYPHILIS

1. Clinical description

Clinical manifestations are many, including localized or diffuse often bilateral mucocutaneous lesions and generalized lymphadenopathy. Flu-like symptoms are common.

2. Criteria for diagnosis

Identification of *T. pallidum* from a lesion compatible with secondary syphilis; OR Compatible clinical picture with laboratory confirmation by either a or b:

- a. Reactive nontreponemal test ($> 1:4$) with no prior diagnosis of syphilis,
- b. Four-fold or greater increase in nontreponemal test titer compared with most recent test for individuals with prior history of syphilis

NOTE: Treponemal test (FTA-ABS or MHA-TP) will be reactive.

EARLY LATENT SYPHILIS

1. Clinical description

No clinical signs or symptoms of syphilis.

2. Criteria for diagnosis

Reactive treponemal and nontreponemal test, AND initial infection that has occurred within previous 12 months as demonstrated by either a, b, or c.

- a. Nonreactive or four-fold lower titer nontreponemal test within past 12 months,
- b. History consistent with untreated primary or secondary syphilis in the past 12 months,
- c. Sexual exposure to a partner with primary or secondary syphilis in the past 12 months, or probable early latent syphilis (documented independently as duration < 1 year) and no history of treatment for syphilis following the exposure.

Source: www.lapublichealth.org/std/providers/definitions.htm (as of February 11, 2004).

APPENDIX B: TECHNICAL NOTES

Data Sources: Data on sexually transmitted diseases (STDs) in LAC are obtained through passive and sometimes active surveillance. California law requires that positive laboratory results of STDs and clinically identified suspected cases be reported to local health departments. Suspected and confirmed STD cases are reported confidentially to California Department of Health Services and the Centers for Disease Control and Prevention as required by California Code of Regulations (Section 2500 and Section 2505).

1. Passive surveillance relies on physicians, laboratories, and other health care providers to report STDs to the Department of Health Services (DHS) using the Confidential Morbidity Report (CMR) forms by telephone, by fax, or by internet.
2. Active surveillance entails STD staff regularly contacting hospitals, laboratories, physicians, and other health providers in an effort to identify all cases of a given STD. In addition, STD staff contact schools, hospitals, jails, student health centers, and sentinel physicians and clinics to collect reports of STDs.

Data Limitations: The data tables in the Sexually Transmitted Diseases in Youth Report should be interpreted considering the following notable limitations:

1. Underreporting. The proportion of cases that are not reported varies for each disease. Syphilis surveillance includes both passive and active surveillance, with detailed follow-up of cases and their sexual partners. Thus, underreporting of early syphilis cases is minimized. Due to the acuteness of symptoms for gonorrhea infection, individuals are more likely to seek treatment, and therefore cases are more likely to be reported. On the other hand, chlamydia infections are often asymptomatic and therefore are more likely to be undiagnosed and underreported. In addition, some health care providers may not be aware of their legal requirements to report STDs.

2. Report Delay. Report delay is defined as the time interval between the date an STD diagnosis was made and the date the case was reported to the Health Department. This delay varies by STD, ranging from 1 day to 1 year or more. Therefore, the impact of report delay must be considered when evaluating trends in case numbers and rates over time. Report delay is especially important when evaluating early syphilis data, due to delays in assigning diagnosis while the case is under investigation.

3. Reliability of Rates. All vital statistic rates including morbidity rates are subject to random variation. This variation is inversely related to the number of cases used to calculate rates. A small number of cases can result in highly unstable rates or proportions. To account for this instability, all rates in this report based on less than 20 events are considered “unreliable”. This translates into a relative standard error of the rate of 23% or more, which is the cut-off rate reliably used by the National Center for Health Statistics. Therefore, rates in this report based on less than 20 events should be interpreted with caution.

4. Place of acquisition of infection. Some cases of STDs reported in this report may have been acquired outside of the county. Therefore, most disease rates more accurately reflect the place of diagnosis than the location where an infection was acquired. However, every effort is made in early syphilis cases to determine where the infection actually occurred, in order to more effectively investigate and control the spread of disease. Early syphilis cases determined to have been acquired outside LAC jurisdiction are not included in this report.

5. Jurisdiction. This report does not include cases identified by Long Beach or Pasadena Health Departments. Cases from these jurisdictions are reported directly to the California Department of Health Services and to the CDC.

Rates:

$$\text{Rate per 100,000 population} = (n / \text{population}) \times 100,000$$

APPENDIX C: STD PROGRAM SURVEILLANCE REPORT REQUEST

To continue to receive sexually transmitted disease surveillance reports for Los Angeles County and other information from the STD Program, please fill out this form and fax or mail it in to the phone number and address provided. **This information will only be used by the STD Program to disseminate surveillance reports and other information to interested parties.** Please be sure to include,

- **e-mail address,**
- **name and mailing address,**
- **and, phone number (to help us keep our distribution list updated)**

<input type="checkbox"/> JMD	<input type="checkbox"/> PA/NP	<input type="checkbox"/> RN	<input type="checkbox"/> PhD	<input type="checkbox"/> Clinic Admin.	<input type="checkbox"/> Other:
------------------------------	--------------------------------	-----------------------------	------------------------------	--	---------------------------------

First Name:	Last Name:	Title (i.e., Director, Coordinator, etc.):

Organization Name:	Street Address:

City:	State:	Zip Code:	Phone:

E-mail:	Type of Organization (i.e., CBO, MCO, OPP, etc.):

Please select the surveillance reports you would like to receive by checking off the appropriate boxes:

<input type="checkbox"/>	All Surveillance and Special Reports from the STD Program
<input type="checkbox"/>	Annual STD Morbidity Report
<input type="checkbox"/>	Quarterly STD Morbidity Report (Planned for 2003)
<input type="checkbox"/>	Monthly Early Syphilis Surveillance Summary
<input type="checkbox"/>	Syphilis Elimination Weekly Activity Report (Available only via e-mail)
<input type="checkbox"/>	Special Morbidity and Project Reports (Periodic)
<input type="checkbox"/>	STD Treatment Updates (Periodic)

Fax to (No cover page needed):	OR	Mail to:
(213) 749-9606		Attn: Maria Venzor STD Program 2615 S. Grand Avenue, Rm. 500 Los Angeles, CA 90007

**APPENDIX D: WHERE TO REPORT COMMUNICABLE DISEASES
IN LOS ANGELES COUNTY**

LABORATORY SLIP	CLINICIAN'S CMR	SEND REPORTS TO:
Syphilis Chlamydia Gonorrhea	Syphilis Chlamydia Gonorrhea Chancroid Non-gonococcal Urethritis Pelvic Inflammatory Disease	Sexually Transmitted Disease Program 2615 S. Grand Avenue, Rm. 450 Los Angeles, CA 90007 Telephone: 213-744-3070 Fax: 213-749-9602
AIDS HIV (<i>As of July 1, 2002</i>)	AIDS HIV (<i>As of July 1, 2002</i>)	HIV Epidemiology Program 600 S. Commonwealth Ave., Suite 805 Los Angeles, CA 90005 Telephone: 213-351-8516 Fax: 213-467-4683 www.lapublichealth.org/hiv/hivreporting.htm
Tuberculosis	Tuberculosis	Tuberculosis Control 2615 S. Grand Avenue, Rm. 507 Los Angeles, CA 90007 Telephone: 213-744-6271 Fax: 213-749-0926
Cryptosporidiosis Diphtheria Encephalitis arboviral <i>E. coli</i> O157:H7 Hepatitis A, acute Hepatitis B, acute Listeriosis Malaria Measles (Rubeola) Plague Rabies Typhoid <i>Vibrio</i> Species	List of reportable diseases: www.lapublichealth.org/acd/cdrs.htm	Acute Communicable Diseases Morbidity Unit 313 N. Figueroa St., Rm. 117 Los Angeles, CA 90012 Telephone: 213-240-7821 Call toll free: 888-397-3993 Fax toll free: 888-397-3778

REQUIRED INFORMATION ON THE CMR:

- ✓ **Patient Information:** Name, gender, ethnic group, date of birth, occupation, address, telephone number, social security number.
- ✓ **Diagnosis:** Diagnosis, date of diagnosis, date of onset.
- ✓ **Reporting Person:** Name, address, telephone number.



COUNTY OF LOS ANGELES
Public Health

Sexually Transmitted Disease Program

2615 South Grand Avenue, Room 500
Los Angeles, CA 90007
Tel: (213) 744-3070 Fax: (213) 749-9606

www.lapublichealth.org/std